



Training Manual on Diagnostic of Weevils of Quarantine Importance

Prepared by:



Institute of Weed Science, Entomology and
Plant Pathology, UPLB, Los Baños, Philippines

In Collaboration with:




ASEAN Network on Taxonomy

2017

DIAGNOSTIC OF WEEVILS OF QUARANTINE IMPORTANCE

TRAINING MANUAL



Dr. Barbara L. Caoili
Dr. Pio A. Javier
Dr. Celia dR. Medina
Dr. Sheryl A. Yap
Dr. Hiraku Yoshitake



ASEAN Regional Training Workshop on Diagnostics of Weevils of Quarantine Importance July 10-22, 2017

Institute of Weed Science, Entomology and Plant Pathology
College of Agriculture and Food Science
University of the Philippines Los Baños

Funded by:
Japan – ASEAN Integration Fund
(JAIF)



Japan-ASEAN Cooperation

Through
ASEAN Plant Health Cooperation Network of the ASEANET
(APHCN-ASEANET)



Table of Contents

Project Title	3
Project Duration	3
Training Outline	3
Background	3
Course Description	3
General Objectives	4
Specific Objectives	4
Training Methodology	4
Training Contents	5
Program of the Training Workshop	6
Resource Persons	12
List of Participants	13
Session 2. CLASSIFICATION, BIOLOGY AND ECOLOGY OF WEEVILS	18
Session 3. HABITUS IDENTIFICATION OF WEEVILS	67
Session 4. WEEVILS IN AGRICULTURAL CROPS	118
Session 6. WEEVILS AS STORAGE PESTS AND THIER CONTROL	152
Session 7. DISSECTION OF GENITALIA	216
Session 8. PRESERVATION TECHNIQUES	218
Session 9. DNA SEQUENCING	259
Session 10. QUARANTINE POLICIES	287
Session 11. RESOURCES OF WEEVILS ON THE WEB	336

TRAINING WORKSHOP ON DIAGNOSTICS OF WEEVILS OF QUARANTINE IMPORTANCE

July 10 - 22, 2017

Institute of Weed Science, Entomology and Plant Pathology
Wing A, Biological Sciences Building
College of Agriculture and Food Science
University of the Philippines Los Banos

I. Project Title: Training Workshop on Diagnostics of Weevils of Quarantine Importance

II. Project Duration: Two weeks, from July 10-22, 2017

III. Training Outline

A. Background

Weevils are small to medium insects that belongs to the Order Coleoptera. They are usually dark-colored, brownish to black. Many are considered as pests of crops or stored products. Most of the small weevils can be present in imported agricultural and stored products. Weevils belonging to genus *Sitophilus*, *Callosobruchus*, *Anthonomus*, *Acanthoscelides*, *Merhynchites*, and *Cosmopolites* are the most serious pests of storage products. The damage that they cause is very destructive and expensive.

On the other hand, there are many other species of weevils that attack agricultural crops, for example, *Sternochetus frigidus*, these weevil is responsible in the down-fall of mango export industry in Palawan Island. Currently, there are observation of weevils feeding or attacking crops in some farm areas especially those areas that where converted from forest or shrubland into agricultural lands. These weevils are not previously seen or recorded feeding on agricultural crops. Hence, quick and accurate identification is vital in quarantine processes and converted farm areas. The proper identification could be able to limit or eliminate entry of these destructive pest to the importing countries as well as proper management or control strategies can be administered.

In this workshop, we aim to teach the participants how to identify weevils with economic importance, and the importance of taxonomy and ecology in managing or controlling pests such as weevils. Also, participants will be able to learn techniques in identification based on morphology and DNA sequences.

B. Course description

This workshop is coordinated by Crop Protection Cluster, College of Agriculture, University of the Philippines Los Banos through the ASEAN Plant Health Cooperation Network (APHCN) of ASEANET project on "Taxonomy capacity building to support market access for agricultural trade in the ASEAN region." This project is funded by the Japan-ASEAN Integration Fund (JAIF) that will be implemented for 2017 covering several activities related to training and attachment programs.

This workshop aims to provide the participants basic knowledge, understanding, and practical skills on the importance of taxonomy, biology, and ecology of weevils in the context of pest management. The training course will cover the following topics: basic taxonomy, biology, ecology, identification and collection techniques, and basic DNA barcoding techniques of weevils with quarantine importance. Likewise, participants will also be taught of proper pest management strategies of these weevils.

C. General Objectives

Lecture: At the end of the training, the participants should be able to analyze the importance of controlling or managing weevils, their biology, and ecology.

Laboratory: At the end of the training, the participants should be able to demonstrate basic skills in identifying the species, genera and families of weevils based on morphology and habits, DNA barcoding, collecting, and preservation and labelling of specimens.

D. Specific Objectives

Lecture:

- 1) To describe the taxonomy, biology, and ecology of weevils with quarantine importance.
- 2) To recognize the importance of weevils in quarantine.
- 3) To discuss that weevils can invade one region to another
- 4) To design appropriate pest management strategies in controlling weevils

Laboratory:

- 1) To describe signs of weevil infestation.
- 2) To identify weevil species based on morphology and DNA barcoding.
- 3) To demonstrate basic techniques in identification, collection in the field, preservation of specimens, and DNA sequencing.

E. Training Methodology

The training will be conducted at the Crop Protection Cluster, UPLB. This would consist of lectures on the proper identification of weevils of quarantine importance and how to manage them. There will also be laboratory or hands-on exercises where participants will be exposed to the actual specimens. Likewise, visit(s) to facilities such as

warehouses or agricultural field experiment stations will be conducted so that participants will have an experience on how to collect weevils.

F. Training Contents

a. Course Outline

SESSION 1. OPENING AND INTRODUCTION

- Registration
- Welcome/Opening Program
- Introduction and overview of the training course
- Group photograph

SESSION 2. CLASSIFICATION, BIOLOGY AND ECOLOGY OF WEEVILS

- Lecture: Classification, biology and ecology of weevils
- Lecture: Economic importance of weevils
- Lecture: Taxonomy of Weevils

SESSION 3. HABITUS IDENTIFICATION OF WEEVILS

- Lecture: Morphology and terminology
- Practice: Identification of specimens using keys

SESSION 4. WEEVILS IN AGRICULTURAL CROPS

- Lecture: Important weevils attacking economically important crops
- Lecture: Control strategies and management of weevils

SESSION 5: FIELD COLLECTION IN STORAGE FACILITIES AND AGRICULTURAL AREAS

- Practice: Field observation and sampling
- Practice: Preliminary identification of specimens

SESSION 6. WEEVILS AS STORAGE PESTS AND THEIR CONTROL

- Lecture: Common Storage pests
- Lecture: Control strategies and management of storage pests
- Practice: Identification of storage pests

SESSION 7. DISSECTION OF GENITALIA

- Practice: Demonstration
- Practice: Hands-on work

SESSION 8. PRESERVATION TECHNIQUES

- Lecture: Preservation techniques
- Practice: Demonstration
- Practice: Hands-on work

SESSION 9. DNA SEQUENCING

- Lecture: Proper DNA techniques and application
- Lecture: Extraction of DNA
- Lecture: Detection of DNA
- Lecture: Sequencing and identification of DNA
- Practice: Demonstration of DNA Techniques
- Practice: Extraction of DNA
- Practice: Detection of DNA

- Practice: Sequencing and identification of DNA

SESSION 10. QUARANTINE POLICIES

- Lecture: Quarantine policies and protocols

SESSION 11. RESOURCES OF WEEVILS ON THE WEB

- Browsing of web sites concerning with weevils

SESSION 12. EVALUATION AND CLOSING PROGRAM

- Post-test evaluation

- Closing Program

G. PROGRAM OF THE TRAINING WORKSHOP

Date/Time/Venue	Topic/Activity	Resource Person(s)/Facilitator
Sunday, 09 July 2017		
	Arrival of Participants	Dr. Sheryl A. Yap and Ms. Priscilla M. Barcial, IWEP
Week 1: Monday, 10 July 2017		
Venue: CPC Auditorium	SESSION 1. OPENING AND INTRODUCTION	
08:45 - 09:00	Registration	Ms. Priscilla M. Barcial
09:00 - 09:30	Welcome Address	Dr. Enrico P. Supangco, Dean, College of Agriculture and Food Science
	Opening Remarks	Dr. Teresita U. Dalisay, Director, IWEP
09:30 - 09:40	Workshop Context and Overview	Dr. Sheryl A. Yap
09:40 - 10:00	Introduction of Participants, trainers and training team	
10:00 - 10:15	Group photograph	Ms. Nikka H. Hamor
10:15 - 10:45	Tea/coffee break	
Venue: CPC Auditorium	SESSION 2. CLASSIFICATION, BIOLOGY AND ECOLOGY OF WEEVILS	
10:45 - 12:00	Lecture: Classification, biology and ecology of weevils	Dr. Hiraku Yoshitake, Institute for Agro-Environmental Sciences, NARO
12:00 - 13:00	Lunch break	
13:00 - 14:30	Lecture: Classification, biology and ecology of weevils	

14:30 -15:00	Lecture: Economic importance of weevils	
15:00 - 15:15	Tea/coffee break	All participants
15:15 - 17:00	Country reports	
Venue: Kamayan sa Palaisdaan	DINNER RECEPTION	All participants Resource persons Training team

Week 1: Tuesday, 11 July 2017

Venue: Lab room	SESSION 3. HABITUS IDENTIFICATION OF WEEVILS	
09:00 - 10:30	Lecture: Morphology and Taxonomy	Dr. Hiraku Yoshitake
10:30 -10:45	Tea/Coffee break	
10:45 - 12:00	Practice: Identification of specimens using keys	Dr. Hiraku Yoshitake, Dr. Sheryl A. Yap
12:00 - 13:00	Lunch break	
13:00 - 14:45	Practice: Identification of specimens using keys	Dr. Hiraku Yoshitake, Dr. Sheryl A. Yap
14: 45 - 15:00	Tea/coffee break	
15:00 - 17:00	Practice: Identification of specimens using keys	Dr. Hiraku Yoshitake, Dr. Sheryl A. Yap

Week 1: Wednesday, 12 July 2017

Venue: Lab room	SESSION 4. WEEVILS IN AGRICULTURAL CROPS	
09:00 - 10:30	Lecture: Weevils in agricultural crops	Dr. Celia dR. Medina, IWEP
10:30 -10:45	Tea/Coffee break	
10:45 - 12:00	Lecture: Control strategies and management of storage pests	
12:00 - 13:00	Lunch break	
13:00 - 14:30	Identification of weevils in agricultural crops	Dr. Celia dR. Medina, Dr. Sheryl A. Yap, Dr. Hiraku Yoshitake

14: 30 – 15:00	Tea/coffee break	
15:00 – 17:00	Identification of weevils in agricultural crops	Dr. Celia dR. Medina, Dr. Sheryl A. Yap, Dr. Hiraku Yoshitake
15:45 - 17:00	Practice: Hands-on work	Dr. Hiraku Yoshitake, Dr. Sheryl A. Yap

Week 1: Thursday, 13 July 2017

Venue: Storage facilities (Industry: Raw materials) **SESSION 5. FIELD COLLECTION IN STORAGE FACILITIES AND AGRICULTURAL AREAS**

07:00 - 07:30	Preparation for field trip (Venue: Hotel)	Training team All participants All resource persons
07:30 - 09:30	Travel from hotel to storage facilities	
09:30 - 10:30	Arrival and briefing of the facility	
10:30 -11:30	Observation and sampling	
11:30 - 13:30	Lunch	
13:30 -15:00	Observation and sampling	
15:00	Return to Hotel	
17:00	Arrival at the hotel	

Week 1: Friday, 14 July 2017

Venue: Storage facilities (Agricultural products) **SESSION 5. FIELD COLLECTION IN STORAGE FACILITIES AND AGRICULTURAL AREAS (continued)**

07:00 - 07:30	Preparation for field trip (Venue: Hotel)	Training team All participants All resource persons
07:30 - 09:30	Travel from hotel to storage facilities	
09:30 - 10:30	Arrival and briefing of the facility	

10:30 -11:30	Observation and sampling
11:30 - 13:30	Lunch/Prayer for Moslems
13:30AM -15:00	Observation and sampling
15:00	Return to Hotel
17:00	Arrival at the hotel

Week 1: Saturday, 15 July 2017

Venue: Farm areas **SESSION 5. FIELD COLLECTION IN STORAGE FACILITIES AND AGRICULTURAL AREAS (continued)**

07:00 - 07:30	Preparation for field trip (Venue: Training team Hotel)	All participants All resource persons
---------------	---	--

07:30 - 09:30 Travel from hotel to the farm

09:30 - 10:30 Arrival

10:30 -11:30 Observation and sampling

11:30 - 13:30 Lunch

13:30AM -15:00 Observation and sampling

15:00 Return to Hotel

17:00 Arrival at the hotel

Week 1: Sunday, 16 July 2017

REST DAY

Week 2: Monday, 17 July 2017

Venue: Lab room **SESSION 6. WEEVILS AS STORAGE PESTS AND THEIR CONTROL**

09:00 - 10:30	Lecture: Common Storage pests	Dr. Pio A. Javier
---------------	-------------------------------	-------------------

10:30 - 10:45 Tea/Coffee break

10:45 - 12:00 Lecture: Control strategies and management of storage pests

12:00 - 13:00	Lunch break	
13:00 - 14:30	Identification of storage pests	Dr. Pio A. Javier, Dr. Sheryl A. Yap, Dr. Hiraku Yoshitake
14:30 - 15:00	Tea/Coffee break	
15:00 - 17:00	Identification of storage pests	Dr. Pio A. Javier, Dr. Sheryl A. Yap, Dr. Hiraku Yoshitake

Week 2: Tuesday, 18 July 2017

Venue: Lab room

SESSION 7. DISSECTION OF GENITALIA

09:00 - 10:30	Practice: Demonstration	Dr. Hiraku Yoshitake
10:30 - 10:45	Tea/Coffee break	
10:45 - 12:00	Practice: Hands-on work	Dr. Hiraku Yoshitake, Dr. Sheryl A. Yap
12:00 - 13:00	Lunch break	
	SESSION 8: PRESERVATION TECHNIQUES	
13:00 - 14:30	Lecture: Preservation techniques	Dr. Sheryl A. Yap
14:30 - 15:00	Tea/Coffee break	
15:00 - 17:00	Practice: Demonstration and Hands on work	Dr. Sheryl A. Yap, Dr. Hiraku Yoshitake

Week 2: Wednesday, 19 July 2017

SESSION 9. DNA SEQUENCING

9:00 - 10:30	Lecture: DNA techniques and their application	Dr. Barbara L. Caoili, IWEP
10:30 - 10:45	Tea/Coffee break	
10:45 - 12:00	Lecture: Extraction of DNA	
12:00 - 13:00	Lunch break	

13:00 - 14:30	Lecture: Detection of DNA	Dr. Barbara L. Caoili
14:30 - 15:00	Tea/Coffee break	
15:00 - 17:00	Lecture: Sequencing and identification of DNA	Dr. Barbara L. Caoili

Week 2: Thursday, 20 July 2017

SESSION 9. DNA SEQUENCING (continued)

09:00 - 10:30	Practice: Demonstration of DNA Techniques	Dr. Barbara L. Caoili/Insect Path lab team
10:30 - 10:45	Tea/Coffee break	
10:45 - 12:00	Practice: Extraction of DNA	Dr. Barbara L. Caoili/Insect Path lab team
12:00 - 13:00	Lunch break	
13:00 - 14:30	Practice: Detection of DNA	Dr. Barbara L. Caoili/Insect Path lab team
14:30 - 15:00	Tea/Coffee break	
16:00 - 17:00	Practice Sequencing and identification of DNA	Dr. Barbara L. Caoili/Insect Path lab team

Week 2: Friday, 21 July 2017

SESSION 10. QUARANTINE POLICIES

9:00 - 10:30	Lecture: Quarantine policies and protocols	Quarantine officer
10:30 - 10:45	Tea/coffee break	
10:45 - 12:00	SESSION 11. RESOURCES OF WEEVILS ON THE WEB	
	Browsing of web sites concerning with weevils	Dr. Hiraku Yoshitake, Dr. Sheryl A. Yap
12:00 - 13:00	Lunch break	

SESSION 12. EVALUATION AND CLOSING PROGRAM

13:00 - 16:00	Post-test evaluation	Dr. Sheryl A. Yap
14:00 - 14:30	Tea/coffee break	
14:30 - 16:30	Closing remarks	Dr. Hiraku Yoshitake
	Information related to Attachment Program	ASEAN representative
	Official closing	Dr. Teresita U. Dalisay
	Presentation of certificates	Dr. Hiraku Yoshitake
		Dr. Sheryl A. Yap
		Dr. Teresita U. Dalisay
	Response from participants	Participant

Week 2: Saturday, 22 July, 2017

DEPARTURE

H. RESOURCE PERSONS

1. Dr. Barbara L. Caoili, Professor, Institute of Weed Science, Entomology and Plant Pathology, College of Agriculture and Food Science, University of the Philippines Los Baños, Los Baños, Laguna, Philippines 4031; E-mail: blcaoili@up.edu.ph; Tel: +6349-536-1315
2. Dr. Pio A. Javier, Research Professor, Institute of Weed Science, Entomology and Plant Pathology, College of Agriculture and Food Science, University of the Philippines Los Baños, Los Baños, Laguna, Philippines 4031; E-mail: pio.javier@yahoo.com.ph; Tel: +6349-536-1315
3. Dr. Celia dR. Medina, Associate Professor, Institute of Weed Science, Entomology and Plant Pathology, College of Agriculture and Food Science, University of the Philippines Los Baños, Los Baños, Laguna, Philippines 4031; E-mail: cdrmedina@yahoo.com; Tel: +6349-536-1315
4. Dr. Sheryl A. Yap, Assistant Professor, Institute of Weed Science, Entomology and Plant Pathology, College of Agriculture and Food Science, University of the Philippines Los Baños, Los Baños, Laguna, Philippines 4031; E-mail: sayap3@up.edu.ph; Tel: +6349-536-2617
5. Dr. YOSHITAKE Hiraku, Research Scientist for Insect Systematics, Institute for Agro-Environmental Sciences, NARO, Kannondai 3-1-3, Tsukuba, Ibaraki 305-8604, Japan; E-mail: zoumushi@affrc.go.jp; Tel: 029-838-8348; Fax: 029-838-8354

I. ASEANET-APHCN TEAM

1. Dr. Lum Keng Yeang, Chairperson, ASEANET-APHCN, c/o CABI SEA, P.O. Box 210, UPM Post, 43400 Serdang, Selangor, Malaysia, E-mail: ky.lum@cabi.org; Telp.: +60-3-89432921.
2. Dr. Soetikno S. Sastroutomo, ASEANET-APHCN, c/o CABI SEA, P.O. Box 210, UPM Post, 43400 Serdang, Selangor, Malaysia, E-mail: s.soetikno@cabi.org; Telp.: +60-3-89432921.

J. LIST OF PARTICIPANTS

BRUNEI

Ms. Norkhadijah Binti Haji Latip

Senior Agriculture Asisstant
 Department of Agriculture and Agrifood of Brunei Darussalam
 Crop Protection Unit, Brunei Agricultural Research Centre, Department of Agriculture and Agrifood, Kilanas, BF2520
 Tel: +673 8 813 533
 E-mail: crop.protection@agriculture.gov.bn

Ms. Helwa Mazlan

Assistant Agricultural Officer
 Department of Agriculture and Agrifood,
 Biosecurity Division, Jalan Menteri Besar, Bandar Seri Begawan, BB3910, Negara Brunei Darussalam
 Tel: +673 7130584
 E-mail: plantbios@gmail.com

CAMBODIA

Mr. Chheng Rasey

Technical Officer
 Department of Plant Protection Sanitary and Phytosanitary, General Directorate of Agriculture, MAFF
 54B/49F, St 395/656, Sangkat Teuk La Ork3, Khan Toul Tork, Phnom Penh, Cambodia
 Tel: 089 839487
 E-mail: raseygda@gmail.com

Ms. Roeun Vanna

Technical Officer
 Department of Plant Protection Sanitary and Phytosanitary, General Directorate of Agriculture, MAFF
 54B/49F, St 395/656, Sangkat Teuk La Ork3, Khan Toul Tork, Phnom Penh, Cambodia
 Tel: 017 869281
 Fax:
 E-mail: Roeunvanna9@gmail.com

INDONESIA**Mr. Kemas Usman**

Technical Officer
Indonesian Agricultural Quarantine Agency
Ministry of Agriculture Republic of Indonesia
Jalan Harsono RM. No. 3 Ragunan,
Pasar Minggu, Jakarta, 12550
Indonesia
Tel : 62 21 7821367
Fax : 62 21 7821367
E-mail: usman.kiemas@gmail.com

Mr. Joni Hidayat

Technical Officer
Applied Research Institute of Agricultural Quarantine (ARIAQ),
JL Raya Kampung Utan – Setu, Mekar Wangi, Cikarang Barat, Bekasi West Java 17520
Indonesia
Tel: 02182618923
Fax: 02182618923
E-mail: joni@buttmkp.org

LAOS**Ms. Sounaly Sommany**

Entomologist, Plant health section
Plant Quarantine Division
Department of Agriculture
Ministry of Agriculture and forestry
Lane Xang Avenue, Patuxay Square
P.O. Box 811 Vientiane Capital
Tel: + 856 20 28268045
Fax: + 856 21 452649
E-mail: nanasounaly@gmail.com

Mr. Touy Bounvilayvong

Technical Officer
Plant Protection Centre, Department of Agriculture, Ministry of Agriculture and Forestry
Tel: + 85621812164
Fax: + 85621812164
E-mail: Bounvilayvongtouy@yahoo.com

MALAYSIA**Mohd Nazri bin Mohd Joha**

Assistant Agriculture Officer
Entomology Unit, Plant Biosecurity Division, Department of Agriculture
Jalan Gallagher, 50632 Kuala Lumpur
Malaysia
Tel: + 6012 3397945
Fax: + 03 26977205
E-mail: nazrijoha@gmail.com

Siti Aisah Binti Jumaat

Assistant Director
Plant Biosecurity Division
Department of Agriculture Jalan Gallagher
50480 Kuala Lumpur,
Malaysia
Tel: + 603 26977141
Fax: + 603 26977205
E-mail: rwo.aisah@gmail.com

MYANMAR**Dr. Ko Ko**

Staff Officer
Plant Protection Division, Baying Naung Rd, West Gyogon, Insein P.O. Box 11011,
Yangon Myanmar
Tel: + 95-9-797342006
Fax: + 95-1640975
E-mail: kokosharp@gmail.com

Mr. Lang Khaw Htang

Deputy Staff Officer
Department of Agriculture,
Falam District, Chin State
Tel: + 95-9-768262852
Fax: + 95-1-640975
E-mail: kokosharp@gmail.com

PHILIPPINES**Ms. Jacqueline R. Ramos**

Agriculturist II
Bureau of Plant Industry –National Plant Quarantine Services Division
Bureau of Plant Industry 692 San andres St. Malate, Manila
Tel: + 02 527 4446 / 09 278596845
E-mail: jacquelin Ramos02@yahoo.com

Dr. Peter Magdaraog

Senior Agriculturist
Bureau of Plant Industry
Diosdado Macapagal Government Centre
Bgy.Maimpis, City of San Fernando, Pampanga
Tel: + 045 455 1755
Fax: + 455 455 1755
E-mail: petermagdaraog@yahoo.com

SINGAPORE**Mr. Chu Si Hao**

Scientist
Agri-Food & Veterinary Authority of Singapore
6 Perahu Road Singapore 718827
Tel: +65 63165194

Fax: +65 63161090
E-mail : [Chua Si Hao@ava.gov.sg](mailto:Chua_Si_Hao@ava.gov.sg)

THAILAND

Mr. Ittipon Bannakan

Professional Entomologist
Department of Agriculture Ladyao Chatuchak
50 Phaholythin Rd.
Bangkok, Thailand 10900
Tel: + 662 9406304
Fax: + 662 9405396
E-mail: bannakan@hotmail.com

Mr. Danai Chaireunkaew

Agricultural officer, Practitioner Level
Plant Quarantine Research Level
50 Phaholythin Rd.
Department of Agriculture Chatuchak
Bangkok, Thailand 10900
Tel: + 66 0 2561 2145 ext 110
Fax: + 66 0 2561 2146
E-mail: musixnoi@hotmail.com

VIETNAM

Dr. Hoang Kim Thoa

Head of Entomology group and Deputy Head Plant Pest and Disease Diagnosis and Identification Division
Plant Quarantine Diagnostic Centre
Plant Protection Department
149 Ho Dac Di Street, Dong Da District,
Ha Noi Capital Vietnam
Tel: + 84 38573424
Fax: + 04 38513746
E-mail: kimthoappd@gmail.com

Mr. Bui Duc Hieu

Technical Officer
Plant Quarantine Sub-department region 1
No 2F –Tran Quang Khai Street – Hong Bang district-Hai Phong City, Vietnam
Tel: 08 0982259679E-mail: Hieukdtv1@yahoo.com.vn



ASEAN Regional Training Workshop on Diagnostics of Weevils of Quarantine Importance

July 10-22, 2017

**Institute of Weed Science, Entomology and Plant Pathology
College of Agriculture and Food Science
University of the Philippines Los Baños**

SESSION 2

CLASSIFICATION, BIOLOGY AND ECOLOGY OF WEEVILS

Lecture Notes by:

DR. HIRAKU YOSHITAKE
Institute for Agro-Environmental Sciences
NARO, Tsukuba, Japan

CLASSIFICATION, BIOLOGY AND ECOLOGY OF WEEVILS

1. Classification of Weevils

Weevils mean beetles which belong to the superfamily Curculionoidea having a worldwide distribution.

Kingdom Animalia

Phylum Arthropoda

Subphylum Hexapoda

Class Insecta

Order Coleoptera

Suborder Polyphaga

Infraorder Cucujiformia

Superfamily Curculionoidea

- Weevils are most diverse in tropical and subtropical latitudes.
- Currently comprise 62,000 described species in ca 5,800 described genera (Oberprieler *et al.*, 2007).
- The total number of existing weevil species has been estimated at ca. 220,000 or higher (Oberprieler *et al.*, 2007), due to the presence of numerous unknown species.
- Virtually all taxa of seed plants in both terrestrial and freshwater habitats, and nearly all plant parts, are utilized as food by weevils (Oberprieler *et al.*, 2014).

The higher classification system of Curculionoidea has not been settled yet. A varying number of weevil families are recognized from 7 to over 20 in different systems.

In modern phylogenetic classifications, the superfamily Curculionoidea comprises 7 families (e.g., Oberprieler *et al.*, 2014):


- 1) **Family Nemonychidae**
- 2) **Family Anthribidae**
- 3) **Family Belidae**
- 4) **Family Attelabidae** (incl. Rhynchitidae)
- 5) **Family Caridae**
- 6) **Family Brentidae** (incl. Apionidae, Nanophyidae)
- 7) **Family Curculionidae** (incl. Brachyceridae, Dryophthoridae, Platypodidae, Scolytidae)

However, this system is somewhat hard to use especially in morphological identification of major groups placed in Curculionidae, the largest family in Curculionoidea. In this lecture, therefore, we fundamentally follow this system, but recognize the following 11 families as major weevil groups from the ASEAN region for convenience in the subsequent lecture "HABITUS IDENTIFICATION OF WEEVILS".

- 1) Family Anthribidae
- 2) Family Attelabidae
- 3) Family Rhynchitidae
- 4) Family Brentidae
- 5) Family Apionidae
- 6) Family Nanophyidae
- 7) Family Curculionidae
- 8) Family Brachyceridae
- 9) Family Dryophthoridae
- 10) Family Platypodidae
- 11) Family Scolytidae

Bibliography

- 1) Leschen R. A. B. & Beutel R. G. (eds), 2014. Handbook of Zoology, Arthropoda: Insecta; Coleoptera, Beetles, Volume 3: Morphology and Systematics (Phytophaga). Walter de Gruyter, Berlin/Boston. 675 pp.
- 2) Morimoto, K., H. Kojima & S. Miyakawa, 2006. The Insects of Japan, Volume 3: Curculionoidea: General Introduction and Curculionidae: Entiminae (Part 1). Phytobiini, Polydrusini and Cyphicerini (Coleoptera). 406 pp. Touka Shobo, Fukuoka.
- 3) Oberprieler, R.G., A.E. Marvaldi & R.S. Anderson, 2007. Weevils, weevils, weevils everywhere. Zootaxa, 1668: 491-520.



Various weevils

2. Biology and Ecology of Weevils

Family Nemonychidae (based mainly on Anderson *et al.*, 2014)

This is a small weevil family comprising 78 described species in 26 genera.




Fig. 3.1.1 *Cimberis compta* (LeConte), adult, dorsal habitus (© Agriculture Canada; illustration by C. Paquette).




Fig. 3.1.2 *Atopomacrus iter* Kuschel adult, dorsal habitus, scale line = 1 mm (© Landcare Research, New Zealand; illustration by D. W. Helmore).




Fig. 3.1.3 *Brarus mystei* Kuschel, adult, dorsal habitus, scale line = 1 mm (© Landcare Research, New Zealand; illustration by D. W. Helmore).




Fig. 3.1.5 *Notomacrus araucariae* Kuschel, adult, dorsal habitus, scale line = 1 mm (© Landcare Research, New Zealand; illustration by D. W. Helmore).




Fig. 3.1.6 *Rhinorhynchus rufatus* (Broun), adult, dorsal habitus, scale line = 1 mm (© Landcare Research, New Zealand; illustration by D. W. Helmore).




Fig. 3.1.4 *Bunyacus monteithi* Kuschel, adult, dorsal habitus, scale line = 1 mm (© Landcare Research, New Zealand; illustration by D. W. Helmore).

Fig. 3.1.8 *Zimmittellus fronto* Kuschel, adult, lateral view (© Landcare Research, New Zealand; illustration by B. Rhode).



Nemonychidae species

Distribution:

- Nemonychidae are distributed in the Australian, Nearctic, Neotropical and Palearctic regions.
- No species are known from the Afrotropical and Oriental regions.
- Nemonychids are most diverse in the southern-temperate Neotropical region (Chile, Argentina and Brazil) and the eastern Australian one (including New Guinea, New Caledonia and New Zealand).

Biology and Ecology:

- Nemonychidae are largely associated with conifers of the families Araucariaceae, Pinaceae and Podocarpaceae, only three genera living on angiosperms: two on Nothofagaceae in the southern-temperate Neotropical region and one on Ranunculaceae in the Palearctic region.

- Adults of all Nemonychidae appear to feed primarily if not exclusively on pollen of their hostplants.
- The eggs of the conifer-associated nemonychids are placed against the pollen-sacs in male cones before shedding the pollen. Larvae feed mainly on pollen and tender sporophylls. The full-grown larvae drop to the ground and then pupate in the soil.
- The larvae of an angiosperm-associated nemonychid, *Nemonyx lepturoides*, are consistently found inside the follicles of *Consolida regalis* (Ranunculaceae). They feed on the large soft seeds and fall to the ground when the fruit capsule splits, and then pupate in a spherical cocoon.



Araucaria araucana pollen cones
<http://powo.science.kew.org/taxon/urn:lsid:ipni.org:names:261681-1>



General appearance:

- Adults: body approximately 1.5–6.0 mm in length, moderately elongate and slightly convex, setose with both decumbent and erect hairs but without any broad scales; cuticle shining or matte, rarely slightly metallic, not otherwise brightly colored.
- Larvae: body usually less than 10 mm long, cylindrical, soft, white; legs very small; vestiture consisting of scattered long and short setae.

Family Anthribidae (based mainly on Mermudes & Leschen, 2014)

Currently including 3,861 species in 3 subfamilies: Anthribinae (28 tribes, 308 genera, 3148 species), Choraginae (5 tribes, 62 genera, 630 species), and Urodontinae (8 genera, 83 species).




Fig. 3.1.9 *Rhinorhynchus rufulus* (Broun), larva, lateral view (© Landcare Research, New Zealand; illustration by B. M. May).




Fig. 3.2.1 Anthribidae. A, *Notochoragus crassus* (Sharp), adult; B, *Araecerus palmaris* (Pascoe) male with female leg (insert); C, *Helmoreus sharpi* Holloway, adult; D, *Arcopais spectabilis* (Broun), larva (© Landcare Research, New Zealand; illustration by D. W. Helmore), scale line = 1 mm.

Distribution:

- Anthribinae and Choraginae are distributed worldwide.
- Urodontinae occur in the Afrotropical and Palaeartic regions.

Biology and Ecology:

- Anthribinae contain herbivores, fungivores, and carnivores.
Herbivores: e.g., larvae of *Ptychoderes* species feed on stems or recently fallen tree trunks.
Fungivores: e.g., *Euparius* species feed on polypore fungi.
Carnivores: e.g., larvae of *Anthribus* species are predators Sternorrhyncha such as scale insects.
- The majority of Choraginae are fungivores as adults and larvae, developing in fungus-infected wood of mainly angiosperms.
- Urodontinae weevils are herbivores, and larvae of most species develop in seeds.



living

of

General appearance:


- Adults: body excluding rostrum 1.0–35.0 mm in length, compact, short to elongate, and convex or depressed; surfaces matte, subglabrous, or rarely glabrous; color black, brown, reddish-brown to gray, rarely green, and often with stripes or maculae.
- Larvae: body cylindrical, c-shaped, circular in cross-section (slightly flattened in), soft, white or pallid; legs usually present; vestiture consisting of scattered setae.



Anthribus nebulosus
<http://www.colpolon.biol.uni.wroc.pl/brachytarsus%20nebulosus.htm>

Family Belidae (based mainly on Marvaldi and Ferrer, 2014)

Containing 38 currently recognized genera and approximately 350 described species. The family is classified into 2 subfamilies, Belinae (ca. 145 species in 25 genera and 3 tribes) and Oxycoryninae (ca. 200 species in 13 genera and 3 tribes).



Bruchela rufipes
<http://www.coleo-net.de/coleo/texte/bruchela.htm>




Fig. 3.3.4 A–D, Oxycoryninae, adults. A, *Aglycyderes setifer* Westwood (Aglycyderini). B, C, *Aralius wollastoni* (Sharp) (Aglycyderini). B, female habitus; C, male head and pronotal anterior margin. D, *Metrioxena* sp. (Metrioxenini: Metrioxenina). E, *Afrocorynus turbatus* Marshall (Metrioxenini: Afrocorynina). Scale lines = 1 mm (A–D, F, © CSIRO, Australia; E, © A. E. Marvaldi).

Distribution:

- Belidae are distributed worldwide, with the greatest diversity in the southern hemisphere.
- Belinae are restricted to the southern continents, with the highest diversity in the Australian region (128 species in 21 genera) and a few also in South America (17 species in four genera).
- Oxycoryninae have broader but fragmented ranges, with the three tribes showing an American, Asian-African, and Pacific-Atlantic island distribution, respectively.

Biology and Ecology:


- The eggs are deposited inside plant tissues and the larvae develop endophytically.
- The larvae of Belinae are borers in stems and branches of shrubs and trees. Those of many Pachyurini and Agnesiotidini develop in gymnosperms (conifers of Araucariaceae, Podocarpaceae and Cupressaceae), and the Belini have mostly angiosperm hosts (especially acacias in Australia).
- Oxycoryninae develop mostly in reproductive plant organs, of gymnosperms (araucarias and cycads) as well as of angiosperms, but also in stems or under bark. The American species develop in reproductive organs, such as female cones of araucaria, male strobili of cycads, root-parasitic angiosperms, fleshy inflorescences of Balanophoraceae, and flowers and fruiting bodies of Hydnoraceae. Aglycyderini are associated with a wide variety of plants (mostly eudicotyledons), with the larvae growing in dead branches or twigs (usually under bark), mining leaves, or boring in freshly dead or dying branches.

General appearance:

- Adults: body 1.5–20 mm in length, flattened to moderately convex and elongate; cuticle dark to reddish-brown or brownish-yellow, with a metallic sheen in a few species; subglabrous or with vestiture, including hair-like setae ranging from sparse to dense.
- Larvae: body form variable, elongate, slightly flattened and parallel-sided to robust and strongly curved, of even width to widest at middle of abdomen.

Family Attelabidae (based mainly on Riedel, 2014)

Composed of ca. 2,500 described species in roughly 150 genera belonging to 2 subfamilies, Attelabinae and Rhynchitinae.



Larvae of Belidae weevils (Marvaldi & Ferrer, 2014)





Fig. 3.4.1 Attelabidae, habitus of adults, scale bar = 1 mm. A, *Minurus seniculus* (Chile); B, *Auletobius consimilis* (North India); C, *Eugnampthus bimaculatus* (North India); D, *Rhynchites auratus* (Europe); E, *Scolocnemus wallacei* (Sumatra); F, *Pterocolus ovatus* (Florida); G, *Aspidobyctiscus pavei* (North India); H, *Attelabus variolosus* (Morocco); I, *Pilolabus viridans* (Mexico); J, *Euscelophilus* sp. (China: Kunming); K, *Lamprolabus bispinosus* (Sumatra); L, *Euscelus scutellatus* (Dominican Republic); M, *Phymatopsinus pustula* (Zimbabwe); N, *Euops indicus* (India); O, *Hoplapoderus hystrix* (South India); P, *Allapoderus manaliensis* (North India); Q, *Paratrachelophorus brachmanus* (Vietnam).

Distribution:

- Attelabidae are found in all major zoogeographic regions but are most diverse in the tropics and absent from New Zealand and Pacific islands. There is one species known from New Caledonia.

Biology and Ecology:

- Their host range at the family level is broad.
- Most attelabids are associated with dicotyledons, such as Rosaceae, Fagaceae, Betulaceae and Salicaceae in the Holarctic region, Myrtaceae in Australia, and Anacardiaceae in America.
- A few Rhynchitinae are associated with gymnosperms, such as Pinaceae and Cupressaceae.
- Oviposition behavior among Attelabidae is diverse, with many shoot-cutting, leaf-cutting, and leaf-rolling species. There are some fruits and seed eaters.



C, Funnel-shaped nidus by *Deporaus betulae*.

D, Longitudinal roll by *Apoderus coryli*.

E, Transverse roll by *Attelabus nitens*.

Nidus construction by the female from a leaf of a food plant (Riedel, 2014).

General appearance:

- Adults: body 2.0 mm to 25.0 mm in length, subglabrous, setose or pilose, but without any broad scales; sensory setae present on elytral sides of most Rhynchitinae, sparse or absent in Attelabinae and Bytiscini; cuticle often metallic or otherwise brightly colored.

- Larvae: body usually C-shaped and subcircular in cross-section, but orthosomatic and depressed in leaf-miners, with a sparse to moderately dense vestiture of setae and moderately densely arranged asperities, which are indistinct in Attelabinae, but distinct in Rhynchitinae.




Fig. 3.4.7 Larva and pupa of *Attelabus nitens*; from Fenili (1952), with friendly permission from the publisher; A, fully grown larva, lateral view; B, abdominal spiracle, ventral (above) and dorsal view (below); C, head of larva, dorsal view; D, larval maxillae and labium, ventral view; E, details of labium; F, maxilla; G, pupa, lateral view; H, pupa, dorsal view; I, pupa, ventral view.

Family

Caridae (based on Oberprieler, 2014)

The smallest weevil family containing only 4 described genera and 6 described species, but several further species occur in Australia.




Fig. 3.5.1 Caridae, adults, dorsal habitus. A, *Car condensatus* Blackburn; B, *Carodes relevatus* Zimmerman; C, *Car pini* Lea; D, *Caenominurus topali* Voss (© Landcare Research, New Zealand; illustration by B. Rhode). Scale bars: 1 mm.

Distribution:

- This family is restricted to Australia, New Guinea, and South America.
- The genus *Car* occurs throughout Australia and another genus *Carodes* is known only from northern Queensland, Australia.
- An undescribed genus with several species is present in New Guinea
- The American genera *Caenominurus* and *Chilecar* are mostly known from southern Chile, South America.

Biology and Ecology:

- All Caridae are associated with conifers, in particular Cupressaceae.
- Much of the biology of the family remains to be discovered.
- In southeastern Australia, the females of a large *Car* species oviposit in young, closed female strobili (cones) of *Callitris endlicheri*.



Callitris endlicheri

http://chabg.gov.au/images/photo_cd/cupressaceae/

- The larva generally feeds in the thick basal parts of the sporophylls, around the woody stem of the strobilus.
- The full-grown larva bores out of the strobilus and drops to the ground, and then pupates in a small cell in the soil.




Fig. 3.5.3 Caridae, larva. A–B, *Car condensatus* Blackburn, C–H, *Car* sp. (*Car condensatus* group) tunneling in female strobilus of *Callitris endlicheri* (larva indicated by arrow). A, larval habitus, lateral view (from May 1994); B, antennal area of head capsule (from Oberprieler 2000) (ant, antenna; f.s., frontal suture; man, mandibular base; st, stemma); scale bar: 1 mm. C, Oviposition hole in strobilus with resin plug; D, larva in base of sporophyll; E, larva tunneling in bases of sporophylls around woody stem of strobilus; F–G, larva feeding in base of sporophyll near stem of strobilus; H, strobilus with two larvae, one in base of sporophylls, the other in seed chamber.

General appearance:

- Adults: body 2.0–5.0 mm in length, shortly elongate, often subparallel in dorsal view, moderately coarsely setose, without scales; cuticle pale testaceous to black.
- Larvae: Body <10.0 mm in length, robust and cylindrical, slightly curved; cuticle sparsely covered with short, fine setae.

Family Brentidae (based on Oberprieler, 2014 and other authors)

The family Brentidae in the wide sense includes the two large subfamilies Apioninae and Brentinae and 4 smaller groups, the Eurhynchinae, Ithycerinae, Microcerinae, and Nanophyinae. In this composition, the Brentidae comprise ca. 540 genera and 4,400 described species.

Distribution:

- The Brentidae are cosmopolitan in distribution, but the different lineages generally have more restricted ranges.
- Eurhynchinae (3 genera & 29 species) is endemic to Australia and New Guinea, the monotypic Ithycerinae to North America, and Microcerinae (3 genera & 67 species) to Africa.
- Apioninae (ca. 205 genera and subgenera & <2,200 species and subspecies), Brentinae (291 genera & ca. 1,760 species) and Nanophyinae (33 genera & ca. 310 species) are more widely distributed and most diverse in tropical regions.

Biology and Ecology:

- The larvae of Eurhynchinae, Ithycerinae and Microcerinae are stem- or root-borers.
- The larvae of Brentinae have adapted to tunneling in dead or dying wood (and also adopted more specialized lifestyles of mycetophagy, predation and myrmecophily).
- The larvae of Apioninae and Nanophyinae have adapted to developing in the softer, living tissues of stems, leaves, flowers, fruits and seeds (including the induction of galls).
- The hosts of Brentidae are predominantly dicotyledonous angiosperms.
- Only a few Apioninae develop on conifers or cycads.

General appearance:

See the figure plates in the following pages. The family Brentidae in the broad sense is too diverse to be grouped by appearance.




Fig. 3.6.1.1 Eurhynchinae. A, *Eurhynchus quadrituberculatus* (Boheman), ♂, dorsal habitus; B, *Ctenaphides maculatus* (Pascoe), ♂, dorsal habitus; C, *Aporhina pulchra* Oberprieler, ♂, dorsal habitus; D, *Eurhynchus laevior* (Kirby), larva, lateral view; E, *Eurhynchus laevior* (Kirby), larva, mesothoracic leg, lateroventral view; F, *Eurhynchus laevior* (Kirby), larva, head, frontal view; G, *Aporhina australis* (Heller), larva, head, frontal view; H, *Aporhina australis* (Heller), larva, maxillae and labium, ventral view; I, *Aporhina australis* (Heller), larva, abdominal segments I and II, showing dorsal carinulae, lateral view; J, *Eurhynchus laevior* (Kirby), larva, left antenna and mandibular attachment, frontal view. Scale bars: A–D, 1 mm; E, 0.1 mm; F, G, I, 0.5 mm. (D–G, I–J after Oberprieler 2000.)




Fig. 3.6.1.2 Eurhynchinae, life history. A, *Eurhynchus laevior* (Kirby), ♀ on leaf of *Persoonia levis* (Proteaceae); B, *Eurhynchus laevior* (Kirby), larva on tunnel in stem of *Persoonia levis*; C, *Aporhina australis* (Heller), ♀ on stem of *Litsea lecfiana* (Lauraceae); D, *Aporhina australis* (Heller), larval galls in stem of *Litsea lecfiana*; E, *Aporhina australis* (Heller), larva in stem gall on *Litsea lecfiana*; F, *Aporhina australis* (Heller), pupa in upper part of stem gall on *Litsea lecfiana*. (C, D, F courtesy of Jack Hasenpusch, Innsfail.)




Fig. 3.6.1.3 Ithycerinae. A, *Ithycerus noveboracensis* (Forster), ♂, dorsal habitus; B, same, larva, left antenna and mandibular attachment, frontal view; C, same, larva, lateral view; D, same, head, frontal view. Scale bars: 1mm. (B after Oberprieler 2000; C-D after May 1993).




Fig. 3.6.1.4 Microcerinae. A, *Microcerus latipennis* Fähræus, ♀, dorsal habitus; B, *Episus gibbosus* Aurivillius, ♂, dorsal habitus; C, *Episus westermanni* Aurivillius, ♂, dorsal habitus; D–F, *Gyllenhalia crinita* Marshall, ♀, lateral habitus; G, *Episus fictus* Gyllenhal, ♂, head in lateral view; H, *Episus fictus* Gyllenhal, ♂, eye; I, *Microcerus latipennis* Fähræus, ♀, eye; J, *Gyllenhalia crinita* Marshall, ♀, left antenna, dorsal view; K, *Episus fictus* Gyllenhal, ♂, left antenna, dorsal view; L, *Microcerus latipennis* Fähræus, ♀, left antenna, dorsal view. M, *Microcerus latipennis* Fähræus, ♂, apex of left hind tibia showing mucro and single spur; N, *Microcerus costalis* Fähræus, metendosternite. Scale bars: A–I, 1mm; J–M, 0.5 mm.




Fig. 3.6.2.1 Brentinae. A, *Lasioryhynchus barbicornis* (Fabricius), adult; B, *Anomobrentus kuscheli* Damoiseau, adult; C, *L. barbicornis*, larva (© Landcare Research, New Zealand).




Fig. 3.6.2.2 Brentinae. A, *Cylas* sp. (South Africa) (© CSIRO, Canberra); B, *Cyphagogus* sp. (Malaysia) (© Landcare Research, New Zealand).




Fig. 3.6.3.1 Apioninae, adults. A, type species of *Apion*, *Apion frumentarium* (Linnaeus); B, *Lepanomus crinalis* Balfour-Browne (by A. Stojczew); C, *Pterapion wagneri* Heller (by J. Kania); D, *Megatracheloides chloris* (Faust) (by J. Świętojańska); E, *Myrmacielus* sp.; F, *Cybebus dimidiatus* (Olivier); G, *Tanaos interstitialis* Fähræus; H, *Antliarhis peglerae* Péringuey. Not at the same scale (Fig. A, © M. Russell; Fig. B, C, © M. Wanat; Fig. D from Wanat 2001, © Mantis, Olsztyn and M. Wanat).

Family




Fig. 3.6.3.6 *Pseudaspidapion botanicum* Alonso-Zarazaga & Wang, third instar larva. A, habitus, lateral; B, head, dorsal; C, mandible, dorsal; D, antenna, dorsal; E, labium, ventral; F, left maxilla, ventral; G, clypeus and labrum, dorsal; H, epipharyngeal lining, ventral. Not at the same scale (all © Wang Zh.-L.).




Fig. 3.6.4.1 Brentidae Nanophyinae, adults, dorsal. A, *Nanophyes brevis* Boheman (Nanophyini), length = 1.8 mm; B, *Corimalia* nr. *aliena* (Faust) (Corimaliini), length = 3.0 mm; C, *Lyalia curvata* Alonso-Zarazaga & Perrin (Nanophyini), length = 4.58 mm; all © M. A. Alonso-Zarazaga.

Curculionidae (based on Oberprieler, 2014 and other authors)

The family Curculionidae in the wide sense includes 10 major subfamilies: Brachycerinae (ca. 8 tribes, 95 genera & 1,350 species), Dryophthorinae (152 genera & 1,200 species), Platypodinae (<1,400 species in 34 genera), Cyclominae (ca. 8 tribes, 148 genera & 1,550 species), Entiminae (ca. 1,370 genera & <12,000 species),

Curculioninae (ca. 350 genera & 4,500 species), Molytinae (ca. 989 genera & 8,700 species), Conoderinae (7,571 species in 940 genera), Cossoninae (ca. 1,700 species in 275 genera), and Scolytinae (ca. 250 genera & 6,000 species) (Oberprieler et al. 2007). In the wide sense, the family Curculionidae, which is the largest among Curculionoidea families, comprises ca. 4,600 genera & 51,000 species.

Distribution:

- The family Curculionidae in the wide sense is cosmopolitan in distribution, occurring in all kinds of habitats from seashores to high mountains and from deserts to rainforests to lakes and rivers.
- All the major subfamilies are also cosmopolitan, although Dryophthorinae and Platypodinae are largely limited to tropical regions.

Biology and Ecology:

- Curculionid larvae generally live endophytically in all parts of plants, from roots to trunks and stems to leaves, flowers, fruits and seeds.
- Many also feed ectophytically on roots in the soil or under water.
- Curculionidae are associated with all groups of plants, apparently ancestrally with angiosperms but secondarily with conifers and cycads, pteridophytes (ferns), bryophytes (mosses) and even algae.
- They have also formed extensive and specialized associations with monocotyledons (especially Brachycerinae and Dryophthorinae but also several groups of Curculioninae, Conoderinae and Molytinae).

-

General appearance:

See the figure plates in the following pages. The family Curculionidae in the broad sense is too diverse to be grouped by appearance.




Fig. 3.7.1.1 Brachycerinae, dorsal habitus. A–D. Brachycerini: A, *Brachycerus albotectus* Péringuey; B, *Euretus aurivillii* Péringuey; C, *Byrsops deformis* (Boheman); D, *Synthocus nigropictus* Pascoe; E, Cryptolaryngini: *Cryptolarynx* sp.; F–L. Erirhinini: F, *Aonychus* sp.; G, *Desmidophorus cumingi* Boheman; H, *Ocladius* sp.; I, *Tadius erirhinoides* Pascoe; J, *Trichocaulus longipilis* Fairmaire; K, *Notaris scirpi* (Fabricius); L, *Echinocnemus* sp.; M–O. Tanysphyrini: M, *Bacosomus avidus* (Oke); N, *Stenopelmus rufinatus* Gyllenhal; O, *Tanysphyrus lemnae* Paykull; P, Myrtonymymini: *Myrtonymus* sp.; Q, Raymondionymini: *Ubychia leonhardi* Reitter. Scale bars : 1 mm except 0.5 mm for P and Q.




Fig. 3.7.1.4 Brachycerinae, immature stages. A, *Brachycerus monachus* Fåhraeus, larva, lateral view; B, *Brachycerus monachus* Fåhraeus, larva, head, anterior view; C, *Brachycerus monachus* Fåhraeus, larva, thoracic spiracle, lateral view; D, *Brachycerus ornatus* (Drury), pupa, ventral view; E, *Desmidophorus crassus* Hubenthal, larva, lateral view; F, *Desmidophorus crassus* Hubenthal, larva, head, anterior view; G, *Ocladius dianthi* Marshall, larva, lateral view; H, *Ocladius dianthi* Marshall, larva, head, anterior view; I, *Ocladius dianthi* Marshall, pupa, ventral view; J, *Echinocnemus* sp., larva, lateral view; K, *Echinocnemus* sp., larva, head, anterior view; L, *Echinocnemus* sp., larva, spiracles of abdominal segments VII and VIII; M, *Lissorhopterus oryzophilus* Kuschel, larva, lateral view; N, *Lissorhopterus oryzophilus* Kuschel, larva, head, anterior view; O, *Lissorhopterus oryzophilus* Kuschel, larva, spiracle of abdominal segment VIII, lateral view; P, *Neochetina eichhorniae* Warner, larva, lateral view; Q, *Neochetina eichhorniae* Warner, larva, head, anterior view; R, *Neochetina eichhorniae* Warner, larva, spiracle of A8. Scale bars: A, B, D, P, 2 mm; E–J, M, 1 mm; C, K, L, N, Q 0.2 mm; O, R, 0.1 mm. (A–C, M–O after May 1994; D after Louw 1990; E–F after Morimoto & Kojima 2006; G–I after Marvaldi 2000; J–L, P–R after May 1994.)




Fig. 3.7.2.1 Bagoini. A, *Bagous alismatis* (Marsham), dorsal habitus; B, *B. luteitarsis* Hustache, dorsal habitus; C, *B. contrarius* O'Brien, dorsal habitus; D, *B. longirostris* Vitale, dorsal habitus; E, *B. longulus* Gyllenhal, dorsal habitus; F, *B. utriculariae* O'Brien, dorsal habitus; G, *B. tubulus* Caldara & O'Brien, dorsal habitus; H, *B. elegans* (Fabricius), dorsal habitus; I, *Bagous* sp. near *interruptus* Faust, dorsal habitus; J, *B. trapae* Parshad, dorsal habitus; K, *B. setosus* LeConte, dorsal habitus; L, *B. tempestivus* (Herbst), prosternal canal (arrow), ventral view; M, *B. olceseii* Tournier, penis, dorsal view; N, *B. peregrinus* Gratshev, penis, dorsal view; O, *B. geniculatus* (Hochhuth), penis, lateral view, sb = setal brush, fdp = "false" dorsal process; P, *B. glabrirostris* (Herbst), spiculum ventrale; Q, *B. freti* Caldara & O'Brien, spiculum ventrale; R, *B. brevipennis* Kirsch, spiculum ventrale; ap = arm, ar = apodeme. Scale bars 1 mm except 0.25 mm for M-R.




Fig. 3.7.2.2 Gonipterini. A, *Bryachus squamicollis* Pascoe, dorsal habitus; B, *Gonipterus platensis* (Marelli), dorsal habitus; C, *Oxyops* sp., dorsal habitus; D, *Iptergonus aberrans* (Lea), dorsal habitus; E, *Pantoreites vittatus* Pascoe, dorsal habitus; F, *Prophaesia cretata* Pascoe, dorsal habitus; G, *Syarbis* sp., dorsal habitus; H, genus from New Caledonia, lateral habitus; I, *Gonipterus* sp., penis in dorsal view; J, *Gonipterus* sp., penis with endophallic sclerite evaginated, lateral view; K, *Gonipterus* sp., endophallic sclerite; L, *Gonipterus platensis* (Marelli), endophallic sclerite; M, *Gonipterus pulverulentus* Lea, endophallic sclerite; N, *Gonipterus lepidotus* Gyllenhal, proventricular armature; O, *Oxyops excavatus* Boisduval, proventricular armature; P, *Iptergonus aberrans* (Lea), proventricular armature; Q, *Syarbis* sp., proventricular armature. Scale bars 1 mm, except K–M, 0.5 mm.




Fig. 3.7.2.3 Hyperini. A, *Donus nidensis* Mazur & Petryszak, dorsal habitus; B, *Hypera arator* (Linnaeus), dorsal habitus; C, *Herpes porcellus* (Lacordaire), dorsal habitus; D, *Coniatus tamarisci* (Fabricius), dorsal habitus; E, *Agriochaeta crinita* Pascoe, dorsal habitus; F, *Gerynassa nodulosa* Pascoe, dorsal habitus; G, *Lycosura bispinosa* Pascoe, dorsal habitus; H, *Nothyperus cionodes* Marshall, dorsal habitus; I, *Cepurus torridus* Olivier, dorsal habitus; J, *Choropholus africanus* Marshall, dorsal habitus; K, *Frontodes brevicornis* Marshall, dorsal habitus; L, *Phelypera distigma* (Boheman), dorsal habitus; M, *Hypera arundinis* (Paykull) larvae feeding on stem of *Sium latifolium* (Apiaceae); N, *Gerynassa* sp., larva, lateral view; O, *Frontodes brevicornis* Marshall with cocoon. Scale bars 1 mm.




Fig. 3.7.2.4 Immature stages. A, *Bagous lutulentus* (Gyllenhal), larva, lateral view; B, *Bagous lutulentus* (Gyllenhal), head capsule, anterior view; C, *Bagous lutulentus* (Gyllenhal), pupa, ventral view; D, *Gonipterus platensis* (Marelli), larva, lateral view; E, *Oxyops concretus* Pascoe, larva, lateral view; F, *Bryachus squamicollis* Pascoe, head capsule, anterior view; G, *Bryachus squamicollis* Pascoe, pupa, ventral view; H, *Hypera postica* (Gyllenhal), larva, lateral view; I, *Hypera postica* (Gyllenhal), larva, thoracic segments, lateral view; J, *Hypera rumicis* (Gyllenhal), head capsule, anterior view; K, *Hypera arundinis* (Paykull), pupa, ventral view (A–C, after Gosik 2009; D–G, after May 1994; H, after Skuhrovec 2006 a; I, after Skuhrovec 2005 a; J, after Skuhrovec 2006 b; K, after Gosik 2008 b).




Fig. 3.7.3.1 Dryophthorinae, adults. A, *Sphenophorus australis* Chittenden (Rhynchophorini), habitus, dorsal, length = 12.0 mm. B, *Sphenophorus cariosus* (Olivier) (Rhynchophorini), habitus, dorsal, length = 10.0 mm. C, *Rhodobaenus tredecimpunctatus* (Illiger) (Rhynchophorini), habitus, dorsal, length = 10.0 mm. D, *Sitophilus granarius* (Linnaeus) (Rhynchophorini), habitus, dorsal, length = 4.0 mm. (From Dillon & Dillon 1972 © Dover Publications; from Bousquet 1990 © Agriculture and Agri-Food Canada.)




Fig. 3.7.3.3 Dryophthorinae, *Sphenophorus brunnipennis* (Germar), (Rhynchophorini), larva. A, habitus, lateral, length = 9.0 mm. B, Maxilla and labium, ventral. C, Maxilla, dorsal (note branched setae). (From May 1993: FNZ, Figs. 1062, 1068a, 1068b, © Landcare Research, New Zealand.)




Fig. 3.7.4.1 Cyclominae, dorsal habitus. A, *Acantholophus niveovittatus* Blackburn (Amycterini); B, *Gagatophorus draco* (Macleay) (Amycterini); C, *Bronchus verrucosus* (Linnaeus) (Hipporhinini); D, *Gronops lunatus* (Fabricius) (Hipporhinini); E, *Epichthonius coronatus* (Boheman) (Cyclomini); F, *Somatodes misumenus* Gyllenhal (Cyclomini); G, *Rhythirrinus inaequalis* (Fabricius) (Rhythirrinini); H, *Aparete palpebrosa* Pascoe (Rhythirrinini); I, *Aades cultratus* (Fabricius) (Aterpini); J, *Chrysolopus spectabilis* (Fabricius) (Aterpini); K, *Euthyphasis acuta* Pascoe (Aterpini); L, *Dichotrachelus sulcipennis* Stierlin (Dichotrachelini); M, *Listroderes difficilis* Germain (Listroderini); N, *Steriphus major* (Blackburn) (Listroderini); O, *Notiomimetes pascoei* Wollaston (Notiomimetini); P, *Aphela helopoides* Pascoe (Notiomimetini). Scale bars 1 mm.




Fig. 3.7.5.2 A, B, Adults, general habitus. A, *Entimus sastrei* Viana; B–D, *Naupactus xanthographus* (Germar). B, Dorsal; C, lateral; D, ventral. Scale bar 5 mm.




Fig. 3.7.5.1 Life cycle of Entiminae. A, *Naupactus leucoloma* Boheman, mature larva in soil; B, *N. leucoloma*, pupa in earth cell; C, *N. cervinus* Boheman, adult feeding; D, *Atrichonotus taeniatus* Berg, posture.




Fig. 3.7.5.3 A–F, Adult morphology, head and mouthparts. A–C, *Platyaspistes argentinensis* Kuschel. A, head, dorsal; B, head and antenna, lateral; C, rostrum and mouthparts, ventral, with scar and process; D–F, *Prostomus scutellaris* Schoenherr, mandibles. D, female, dorsal; E, male, dorsal; F, male, ventral.




Fig. 3.7.7.1 Molytinae adults, dorsal: A, *Cryptorhynchus lapathi* Dejean (Cryptorhynchini), length = 8.2 mm; B, *Gasterocercus longipes* Kôno (Gasterocercini), length = 11.2 mm; C, *Torneuma sardoum* Desbrochers (Torneumatiini), length = 2.8 mm; D, *Psepholax sulcatus* White (Psepholacini), length = 7.8 mm; E, *Mechistocerus nipponicus* Kôno (Aedemonini), length = 9.6 mm; F, *Sophrorhinus inseperata* Faust (Sophrorhinini), length = 5.9 mm; G, *Hylobius abietis* (Linnaeus) (Molytini: Hylobiina), length = 12.7 mm; H, *Liparus germanus* (Linnaeus) (Molytini: Molytina), length = 15.3 mm; I, *Plinthus megerlei* (Panzer) (Molytini: Plinthina), length = 11.3 mm; (lengths given from base of rostrum to posterior of abdomen) © Natural History Museum, UK.




Fig. 3.7.7.2 Molytinae adults, dorsal: A, *Typoderus machadoi* Marshall (Typoderini), length = 4 mm; B, *Amorphocerus talpa* Schoenherr (Amorphocerini), length = 5.7 mm; C, *Dreuxetes carinirostris* (Champion) (Anchonini), length = 6 mm; D, *Cleogonus rubetra* (Fabricius) (Cleogonini), length = 9.9 mm; E, *Cycloteres audouini* Boheman (Cycloterini), length = 11.7 mm; F, *Sclerocardius africanus* Boheman (Sclerocardiina), length = 15.2 mm; G, *Theogenete campbelli* Anderson (Lymantini), length = 2.4 mm; H, *Euderus lineicollis* (Wiedemann) (Euderini), length = 8.8 mm; I, *Ithyporus setulosus* Hustache (Ithyporini), length = 8.8 mm; J, *Lepyrus pinguis* Casey (Lepyrini), length = 13.2 mm; K, *Phoenicobates vittatus* Champion (Phoenicobatini), length = 4.3 mm; (lengths given from base of rostrum to posterior of abdomen)
© Natural History Museum, UK.




Fig. 3.7.7.3 Molytinae adults, dorsal: A, *Alcidodes senex* Sahlberg (Mecysolobini), length = 4.3 mm; B, *Phrynixus intricatus* Broun (Phrynixini), length = 4.6 mm; C, *Pissodes pini* (Linnaeus) (Pissodini), length = 7.9 mm; D, *Juanorhinus ruficeps* (Aurivillius) (Juanorhinini), length = 3.3 mm; E, *Orthorhinus cylindrirostris* (Fabricius) (Orthorhinini), length = 18.6 mm; F, *Trachodes hispidus* (Linnaeus) (Trachodini), length = 3.7 mm; G, *Aorus anthracinus* Brancsik (Amalactini), length = 9.2 mm; H, *Petalochilus gemellus* (Gyllenhal) (Petalochilini), length = 10.3 mm; I, *Omophorus stomachosus* Boheman (Metatygini), length = 10.2 mm; J, *Peribleptus scalptus* Boheman (Paipalesomini), length = 14.1 mm; K, *Trigonocolus vitticollis* Marshall (Trigonocolini), length = 4.0 mm; (lengths given from base of rostrum to posterior of abdomen) © Natural History Museum, UK.




Fig. 3.7.7.4 Molytinae adults, dorsal: A, *Cholus albicinctus* Germar (Cholini), length = 14.4 mm; B, *Pacholenus canescens* Vanin & Reichardt (Pacholenini), length = 9.6 mm; C, *Sternechus cicatricosus* Fiedler (Sternechini), length = 8.2 mm; D, *Guioperus griseus* Perty (Guioperini), length = 14.3 mm; E, *Dinomorphus pimeloides* Perty (Dinomorphini), length = 22 mm; F, *Brachyceropsis tuberculosus* (Gyllenhal) (Brachyceropseini), length = 17.4 mm; G, *Lithinus superciliosus* Klug (Lithinini), length = 22.4 mm; H, *Styanax scrobiculatus* Roelofs (Styanacini), length = 8.3 mm; (lengths given from base of rostrum to posterior of abdomen) © Natural History Museum, UK.




Fig. 3.7.8.1 Mesoptiliinae adults, dorsal. A, *Carcilia strigicollis* Roelofs (Carciliini), length = 10 mm; B, *Magdalis memnonia* Gyllenhal (Magdalidini), length = 6.5 mm; C, *Apocnemidophorus blandus* Hustache (Mesoptiliini), length = 5.5 mm; D, *Laemosaccus ruficornis* Champion (Laemosaccini (North America), length = 4.5 mm; E, *Neolaemosaccus notatus* (Pascoe) (Laemosaccini, Australia), length = 5.6 mm (lengths given from base of rostrum to posterior of abdomen).




Fig. 3.7.9.1 Curculionidae, Conoderinae: dorsal habitus of Conoderitae (A–D), Bariditae (E–H), Ceutorhynchitae (I–K), and Oorbitiditae (L); length without rostrum. A, *Cratosomus vaginalis* (Linnaeus), length = 25 mm; B, *Corysomerus capucinus* (Beck), length = 2.9 mm; C, *Campyloscelus westermanni* Boheman, length = 10.8 mm; D, *Copturus curvaturatus* (Heller), length = 7.0 mm; E, *Baris artemisiae* (Panzer), length = 4.2 mm; F, *Eisonyx crassipes* LeConte, length = 4.9 mm (from Pakaluk & Carlow 1994, © E. P. Roberts); G, *Apostasimerus serrirostris* Boheman, length = 6.5 mm; H, *Diorycentrinus tuberculatus* Hustache, length = 4.6 mm; I, *Craponius inaequalis* (Say), length = 3.3 mm; J, *Eubrychius velutus* (Beck), length = 2.0 mm; K, *Hadroplontus trimaculatus* (Fabricius), length = 3.9 mm; L, *Oorbitis cyanea* (Linnaeus), length = 2.4 mm.




Fig. 3.7.10.6 Habitus, adults. A, *Acalyptus carpini* (Schoenherr) (Acalyptini); B, *Eudelodes bicolor* (Faust) (Acalyptini); C, *Acentrus histrio* (Schoenherr) (Acentrusini); D, *Anoplus roboris* Suffrian (Anoplini); E, *Anthonomus rectirostris* (Linnaeus) (Anthonomini); F, *Camarotus* sp. (Camarotini); G, *Cionus olivieri* Rosenschöld (Cionini); H, *Cerocranus extremus* Kuschel (Cranopoeini); I, *Sigastus fuscodorsalis* (Heller) (Cryptoplini); J, *Curculio glandium* Marsham (Curculionini); K, undescribed genus from Australia with horizontal mandibles, seven-segmented funicles, divaricate claws and trochanteral setae (Curculionini); L, *Derelomus piriformis* (Hoffmann) (Derelomini). Scale bars 1 mm.




Fig. 3.7.10.7 Habitus, adults. A, *Aphanonyx latesetosus* (Hustache) (Diabathrariini); B, *Dorytomus longimanus* (Forster) (Ellescini); C, *Ellescus bipunctatus* (Linnaeus) (Ellescini); D, *Ludovix fasciatus* (Gyllenhal) (Erodiscini); E, *Ita chyanoni* Meregalli & Borovec (Itini); F, *Gymnetron beccabungae* (Linnaeus) (Mecinini); G, *Mecinus janthinus* Germar (Mecinini); H, *Galloisia inflata* Hustache (Microstylini); I, *Microstylus rufus* Schoenherr (Microstylini); J, *Nerthops guttatus* Schoenherr (Nerthopini); K, *Acallopius* sp. (Nerthopini); L, *Teinomphasus robustus* Oberprieler (Ochyromerini). Scale bars 1 mm.




Fig. 3.7.11.1 A, B, *Mesites fusiformis* Wollaston, tribus Cossonini (length 4.0 mm); C, D, genus near *Stenoscelodes*, tribus Onycholipini (length 4.0 mm).




Fig. 3.7.12.1 A, B, Scolytinae – *Scolytoplatypus tycon* Blandford (female), tribe Scolytoplatypodini (length 4.2 mm); C, D, *Hylesinus crenatus* (F.), tribe Hylesinini (length 4.5 mm).




Fig. 3.7.13.1 A, B, *Dinoplatypus pseudocupulatus* Schedl (male, length 3.5 mm); C, D, *Schedlarius mexicanus* (Chapuis) (length 5.6 mm).

3. Economic Importance of Weevils

Due to their broad host range at the superfamily level, Curculionoidea contains a large number of agricultural, forestry, and storage pests.

Agricultural pests:

- Rice water weevil, *Lissorhoptrus oryzophilus* (Curculionidae)
Serious pest in rice production. Native to N America. Spread to other regions including Asia.
- Sweet potato weevil, *Cylas formicarius* (Brentidae)
Most serious pest of sweet potato. Widely distributed in tropical regions.
- Red palm weevil, *Rhynchophorus ferrugineus* (Dryophthoridae)
The most damaging pest of palms. Native to SE Asia but introduced to many other regions.
- Mango palm weevil, *Sternochetus frigidus* (Curculionidae)
Serious pest of the flesh of mango. Probably originated from Malaysia.

... and many other examples

Forestry pests:

- Bark beetles (Scolytidae spp.)
Containing many injurious pests in wood production.
- Oak ambrosia beetle, *Platypus quecivorus* (Platypodidae)
Injurious pest of oaks and chinquapin trees. Widely distributed in Asia, including temperate, subtropical and tropical zones.

... and many other examples

Storage pests:

- Rice Weevil, *Sitophilus oryzae* (Dryophthoridae)
Is distributed worldwide.

- Maize weevil, *Sitophilus zeamais* (Dryophthoridae)
Is distributed worldwide, but less common in temperate zones.

Invasion risks of pest weevils:

- An invasive species tends to cause serious damages in “new ground”.
- Invasion risks of pest species are increasing due to the accelerated international trades by globalization.
- Importance of plant quarantine is greatly increasing.
- Quick and accurate species identification is important for monitoring invasive species and taking prompt measures against them.
- Generally, species identification of weevils is very difficult due to the delay in weevil taxonomy. A new pest weevil can be an undescribed species, which is new to science.

Bibliography

- 1) Genka, M. & H. Yoshitake, 2014. Chronological change of taxonomic composition of exotic weevils (Coleoptera, Curculionoidea) intercepted by Japanese plant quarantine. Research bulletin of the Plant Protection Service Japan, (50): 17–46. (In Japanese, with English summary.)
- 2) Yoshitake, H., S. Miyahara, M. Nishino & K. Suzuki, 2012. *Metapocyrtus* (*Trachycyrtus*) *hederaeophilus* sp. nov. (Coleoptera, Curculionidae, Entiminae), a pest of the English ivy cultivated in Mie Prefecture, Honshu, Japan. Japanese Journal of Systematic Entomology, 18: 261-267.
- 3) Yoshitake, H., T. Fujisawa & J. Goto, 2011. A new *Ceutorhynchus* species (Coleoptera, Curculionidae) attacking field-cultivated wasabi (Brassicaceae) in Northern Japan. Japanese Journal of Systematic Entomology, 17: 293-299.



ASEAN Regional Training Workshop on Diagnostics of Weevils of Quarantine Importance

July 10-22, 2017

**Institute of Weed Science, Entomology and Plant Pathology
College of Agriculture and Food Science
University of the Philippines Los Baños**

SESSION 3 & 4


HABITUS IDENTIFICATION OF WEEVILS

Lecture Notes by:



DR. HIRAKU YOSHITAKE
Institute for Agro-Environmental Sciences
NARO, Tsukuba, Japan

HABITUS IDENTIFICATION OF WEEVILS

1. Morphology and Terminology – Adults: General



1. Morphology and Terminology – Adults: Genitalia



Female genitalia, in lateral view

2. Morphology and Terminology – Larvae (Oberprieler et al. 2014)




Fig. 3.5 Structure and chaetotaxy of generalized weevil larva (modified after Marvaldi 1999). A, lateral view; B, ventral view; C, caudal view.

3. Morphology and Terminology – Pupae (Oberprieler et al. 2014)




Fig. 3.6 Structure and chaetotaxy of weevil pupa (*Tranes* Schoenherr, Curculionidae) (modified after May 1994). A, ventral view; B, dorsal view (*aps*, apical pronotal seta; *bps*, basal pronotal seta; *brs*, basirostral seta; *dps*, discal pronotal seta; *drs*, distirostral seta; *fes*, femoral seta; *lps*, lateral pronotal seta; *msns*, mesonotal seta; *mtns*, metanotal seta; *os*, orbital seta; *pds*, postdorsal seta; *pos*, postorbital seta; *sos*, supraorbital seta, *pls*, pleural seta; *ts*, tergal seta; *vs*, vertical seta).

Weevil pupae are exarate, with free appendices and distinct body regions (head, thorax, abdomen). The integument is usually creamy white. Setae vary in shape, number and disposition in different taxa. The head is deflexed beneath the prothorax and the rostrum onto the legs.

HABITUS IDENTIFICATION OF WEEVILS

2. Identification of specimens using keys

In this training workshop, a key to major families of the superfamily Curculionoidea and a key to major subfamilies of the family Curculionidae mainly from the ASEAN region are provided in order to learn how to sort weevil specimens based on morphological traits.

In addition, an illustrated key to *Caulophilus oryzae* (Curculionidae) and *Sitophilus* species (Dryophthridae), all of which are important storage pests, is given in order to learn species identification of weevils practically based on morphological traits.

See **Appendix: Glossary of Weevil Characters** when necessary.

Major Curculionoidea families from ASEAN Region



1. Anthribidae



2. Scolytidae

(<http://www.ffpri-kys.affrc.go.jp>) (<http://www.ffpri-kys.affrc.go.jp>)



3. Platypodidae



4. Attelabidae



5. Rhyncitidae



6. Dryophthridae



7. Apionidae



8. Nanophyidae



9. Brentidae





10. Brachyceridae





11. Curculionidae


11 major families are known from the ASEAN region

An Illustrated Key to Major Curculionidea Families from the ASEAN Region






(II)



Ventrite similarly articulate from 1st to 4th. 5th ventrite deeply articulate with 4th basally.

1st and 2nd ventrites fused together. 3rd to 5th deeply and similarly articulate to each other basally.




Next (III)

Next (IV)

(III)




Tibiae mucronate in both sexes, uncinata in males. Inner margins of fore tibiae serrate.



Attelabidae


Tibiae simple apically. Inner margins of fore tibiae simple.



Rhynchitidae


(II)

Ventrite similarly articulate from 1st to 4th. 5th ventrite deeply articulate with 4th basally.



Next (III)


1st and 2nd ventrites fused together. 3rd to 5th deeply and similarly articulate to each other basally.



Next (IV)


(IV)

The first segment of antennal club smooth, sparsely setiferous.







Dryophthoridae

Antennal club entirely pubescent.




Next (V)






(VII)

Antennae orthocerous.



Brentidae

Antennae with scape long, geniculate.



Curculionidae


Brachyceridae



Desmidophorus crassus

- Distributed mainly in Africa
- Several species from the ASEAN region
- Containing some pests of *Hybiscus*

An Illustrated Key to Major Subfamilies of the Family Curculionidae from the ASEAN region



Entiminae

Exceptional taxa



Pachyrhynchini

- Mandibles neither with cusp nor its scar.
- Many species from the Philippines




Sitonini


- Mandibles neither with cusp nor its scar.
- Several species from the Philippines



Euphyllobiomorphus

- Rostrum slender, cylindrical
- Known only from the Ryukyus, Japan
- Just great if you find it in the Philippines (possibly in high mountains)





Orobittidae

- Only 1 sp. of *Orobittis* is known from East Asia
- Feeding on *Violaceae* plants
- Just great if you find it in the ASEAN region (possibly in high mountains)




Orobittis species



A larva of *Orobittis* species
in a seed capsule of *Viola* sp.


①

1st ventrite strongly narrowed behind hind coxae and nearly subdivided into three parts by coxal cavities




Orobittidinae

1st ventrite normal, not subdivided by coxal cavities



Next


Mesepimera almost as large as mesepisterna, strongly ascended and visible dorsally between bases of pronotum and elytra.





Ceutorhynchinae

Only a few species have been known from the Philippines, but there are many undescribed species


Mesepimera not ascended, invisible dorsally





Next




An Illustrated Key to Major Subfamilies of the Family Curculionidae from the ASEAN region





Labial palpi 1- or 2-segmented, rigid, inserted in a hole at latero-ventral corners of prementum behind apex, or often obliterate. Prementum truncate, vertical and thick at apex.


apex of prementum



Lixinae

Labial palpi 3-segmented, inserted on Apical margin of prementum, which is more or less rounded and not forming a vertical flat surface at apex.

apex of prementum




Apex of rostrum
in ventral view

③

③

Tarsal grooves of fore legs oblique to tibial axis, trough-like for receiving tarsi when contacted and bordered laterally by setal comb oblique to axis of tibia.




setal comb



Next

Tarsal grooves of fore legs broadly flattened and bare behind tarsal sockets occupying major exterior surface behind apex. Inner setal comb translocated to ventral margin.

outer side







inner side



setal comb



Cossoninae







An Illustrated Key to Species feeding on Storage seeds
 (modified from Yoshida, Watanabe & Sonoda, 1989)

1A Antennal club pubescent at apex (Fig. 1A-a). Antennal funicle six-segmented (Fig. 1A-b). Pygidium exposed. ----- 2

1B Antennal club entirely pubescent (Fig. 1B-a). Antennal funicle seven-segmented (Fig. 1B-b). Pygidium not exposed. Body dark to blackish brown, about 3.0 mm in length.----- ***Caulophilus oryzae* (F14-1)**

Central American species which extends to the West Indies and the far south of the USA. Known as the broad-nosed grain weevil, an important pest of storage seeds.

2A Rostrum (Fig. 2A-a) more than twice as long as head, with antennal scrobes which do not touch eyes posteriorly (Fig. 2A-b). ----- 3

2B Rostrum (Fig. 2B-b) nearly twice as long as head; upper margins of antennal scrobes touching eyes posteriorly. Body dark to blackish brown, 3.0–5.0 mm in length. --
----- ***Sitophilus linearis* (F15-1)**

Pan-tropical. Known as the tamarind weevil, an important pest of tamarind and other Caesalpinioideae.

3A Pronotum with elliptical punctures (Fig. 3A). Elytral striae deep but weakly punctured; each interval nearly as wide as stria. Body entirely dark brown, shiny, 2.5–4.0 mm in length. ----- ***Sitophilus granarius* (F15-2)**

Subcosmopolitan. Known as the wheat weevil, grain weevil or granary weevil, a serious pest of stored cereal grains.

3B Pronotum with subovate punctures (Fig. 3B). Elytra strongly striate-punctured; each interval narrower than stria. Body dark to blackish brown; elytra usually with two pairs of light-colored patches. ----- 4


4A Scutellum with lateral margin (Fig. A-a) shorter than basal margin (Fig. A-b). -----
----- ***Sitophilus zeamais* (F15-3)**


Cosmopolitan. Known as the maize weevil or as the greater rice weevil, an injurious pest of cereal crops.

4 B Scutellum with lateral margin (Fig. B-a) as long as basal margin (Fig. B-b).

----- ***Sitophilus oryzae* (F15-4)**

Cosmopolitan. Known as the rice weevil, an injurious stored product pest attacking cereal crops, including wheat, rice, and maize.





HABITUS IDENTIFICATION OF WEEVILS

Appendix: Glossary of Weevil Characters
 (<http://weevil.info/glossary-weevil-characters>)

Accessory glands (female genitalia) – glands associated with the female genitalia, which in at least some cases contain symbiotic organisms. Probably not homologous throughout Curculionoidea, and appear in different forms and attached to different parts of the genitalia. See Kuschel (1987).

Adelognatha (informal group of weevils) - a group of Curculionidae characterised by the adelognathous character state (q.v.), broadly equivalent to the Entiminae sensu Thompson (1992). They are thus also characterised by a short, broad rostrum, the deciduous mandibular process present, larval antennae lacking a projecting apical part; larvae living free in soil. Some Entiminae have phanerognathous mouthparts, however (q.v.). The group has no formal status. See also 'Phanerognatha'.

Adelognathous (head, mouthparts) – refers to the maxillae being concealed by the prementum. Thompson (1992) notes that in some cases the stipites are visible at the posterior angles of the prementum, and that in fact the degree to which the maxillae are concealed by the prementum is very variable. The contrasting term is 'Phanerognathous' (q.v.)

Adelphic – female mating with more than one male – not limited to weevils.

Aedeagal apodemes (male genitalia) – see Penis apodemes

Aedeagus (male genitalia) – Combined structure comprising the tegmen and penis. See also 'pedal' and 'pedotectal' as types of aedeagus.

Aggonoporium (male genitalia) – Term coined by Arzanov (2003) for sclerite or complex of sclerites in the endophallus around the junction with the ductus ejaculatorius. The sclerites may participate in attaching the endophallus to the bursa copulatrix during copulation. See Transfer apparatus.

Airtube (larva) – respiratory appendage (May, 1994)

Annulated – segmented; in rings – sometimes used of antennal club.

Anapleural suture (metathorax) – see metanepisternal suture (preferred term).

Anepisternum (thorax) - anterior sclerite of the peluron, discernable on the mesothorax and metathorax in weevils (mesanepisternum, metanepisternum). Synonym: episternum. (definition changed May 2012)

Anophthalmic (head) – lacking eyes.

Antenna (head) - sensory appendage on either side of the rostrum, comprising 11 articles: scape, 7-segmented funicle and 3-segmented club) (see under 'Antennal segment' for discussion of terminology of the antennal 'segments'). Although sometimes referred to as 12-segmented, this depends on the interpretation of the club components (q.v.) and is apparently a misapprehension. (see states: geniculate antenna, subgeniculate antenna) See also Antennal segment, desmomere, antenomere. (definition changed 1/06/2012)

Antennal article (head) – see Antennal segment (preferred term). (definition changed 23/11/2011)

Antennal club (head) - see under Club.

Antennal insertion (head) – point of attachment of the antenna. Antennal insertions are considered to be exposed when at least some portion of each antennal socket is visible from above the long axis of the rostrum. (After Lawrence, Beutel, Leschen & Slipinski, 2010).

Antennal segment (head, antenna) – individual elements of the antenna. The terms used suffer from a difference in meaning between morphological correctness and in practice. Morphologically the antenna, like other insect antennae, comprises three true 'segments': a scape, pedicel and flagellum. The flagellum may be secondarily subdivided into a number of antennomeres, which are not true segments (in terms of holding muscle insertions). However, the traditional division in weevils is into scape, funicle and club, the funicle comprising the pedicel and up to 6 antennomeres of the flagellum (rarely seven if an antennomere more usually part of the club is separate), and the club a further three flagellum antennomeres. The pedicel is sometimes referred to as the first antennomere or first segment of the funicle. The terms pedicel and flagellum are not used in weevil taxonomy. This provides confusion over terminology, with the different articles being termed segments, antennomeres, desmomeres, joints etc. The elements of the club are rarely termed antennomeres but the term 'clubomere' has been used (not recommended here). The term 'segment' is adopted here for all articles, recognizing the morphological inappropriateness of this. Synonyms: antennal article, antennomere, desmomere (in part), clubomere (in part), joint. (definition changed 1/06/2012).

Antennomere (head, antenna) – see Antennal segment (preferred term). The issues surrounding the use of this and other terms for the structure are discussed under 'Antennal segment' above. (definition changed 01/06/2012).

Anterior tentorial pit (head) – the invagination of the head capsule corresponding to the position of the anterior arms of the tentorium inside the head. The Anterior Tentorial Pits lie at the anterior end of the subgenal sulcus in weevils, near the mandibular articulation (Lyal, 1995).

Apical membranous lobe (male genitalia) – unsclerotised posteriorly-directed extensions of parameroid lobes on tegmen (Wanat, 2007).

Apical setal comb (leg, tibia) – row of strong setae around the apex of the tibia and projecting distad. There is often a gap dorsally between the comb on the posterior and anterior faces, to allow the tarsus to be reflexed. The anterior comb may extend along the dorsal face of the tibia (e.g. some Molytinae), in which case it is termed 'ascending'. (Thompson, 1992). See also 'secondary comb', 'corbel'. Synonyms: distal comb, tibial comb, setose fringe, fringing setae.

Apodeme - chitinous ingrown of the exoskeleton to which muscles are attached; rod-like projection from any of several structures (e.g. penis, tegmen).

Apodemes of penis (male genitalia) – see Penis apodemes.

Apodeme of tegmen (male genitalia) – See Tegminal apodeme

Appendiculate claw (leg, tarsus) - See Tarsal Claw.

Apterous – lacking wings

Baculum (pl – baculi) (female genitalia) – Sclerotised bars or struts which serve to support a membranous ovipositor.

Basal arms (of spiculum gastrale) (male genitalia) – the posterior fork of the spiculum gastrale, on which various muscles attach. Synonym; furcal arms.

Basal piece (male genitalia) – see Tegminal ring (sometimes also tegminal apodeme).

Basal pocket (elytra, submarginal fold) - the region of the sublarginal fold that diverges away from the outer wall of the elytra to form a more or less deep pocket to accommodate the pleurite of the basal ventrites and the metathoracic lateral sclerites. See also 'Submarginal fold'.

Basal sclerites (male genitalia) – paired (usually) sclerites lying near the junction of the endophallus with the ductus ejaculatorius. Of variable form. See Transfer apparatus.

Basisternum (thorax) - the anterior part of each sternum, meeting the sternellum posteriorly. Sometimes used incorrectly for the sternellum (q.v.) on the prothorax, but can be distinguished from this because the sternellum lies between and behind the coxae, whilst the basisternum is before the fore coxae.

Beak see rostrum (preferred term).

Bevel (leg, tibia) – see discussion under Corbel

Bifid claw (leg, tarsus) see "Tarsal Claw"

Buccal cavity (head, mouthparts) – mouth cavity

Bursa (female genitalia) – large closed membranous lobe of female genitalia branching from vagina generally at same point as common oviduct and spermathecal duct. Synonyms: bursa copulatrix; copulatory pouch; vaginal bursa. [emended June 2012]

Bursa copulatrix (female genitalia) – see Bursa

Bursal sclerite (female genitalia) – sclerite lying in the vaginal and bursal floor around the junction with the common oviduct and spermathecal duct.

Cap-piece (male genitalia) – see Parameroid plate.

Cardo (head, mouthparts) – basal piece of the maxilla

Cervical sclerites (head) - small trapezoidal triangular sclerite or pair of sclerites at the posterior end of the gular suture(s) (Lyal, 1995). Synonym: postgula.

Claw (leg, tarsus) - see 'Tarsal claw' (preferred term).

Club (head, antenna) - the apical antennal segments. The club generally has three segments (see under 'Antennal segment' for discussion of terminology of the antennal 'segments'), either separate as in some basal groups or combined as in most Curculionoidea. In many weevils there appear to be 4 club segments, each delimited by a suture and each bearing a ring of elongate setae. However, the form of the groove

between the apical 'segment' and the others differs from that between the true antennomeres (Wanat, 2001). The apparent 4-segmented club has been termed 'pseudotetramerous' by Wanat, 2001. The elements may be termed 'club segments', 'club antennomeres' or 'antennomeres of the club', the first of these being least clumsy (although inaccurate); the term 'clubomere' has also been used but is linguistically dubious. Key character states: loose: articulation between segments and segment 'neck' visible; compact: articulation and basal constrictions not seen, but borders between segments clearly visible; connate: borders or sutures between segments partly or completely lost. (Wanat, 2001) Synonym: Antennal club. (definition changed 1/06/2012)

Clubomeres (head, antennal club) – the antennomeres that make up the club of the antenna. – see Antennal segment (preferred term). (definition changed 23/11/2011)

Collar (prothorax) – anterior of prothorax when delimited, at least laterally, by a groove or simply an angle, from the rest of the prothorax to enclose the base of the head capsule; often developed into postocular lobes.

Collum (female genitalia, spermatheca) – the part of the spermatheca to which the spermathecal duct is connected (see duct lobe). [Emended June 2015].

Common oviduct (female genitalia) – see median oviduct.

Complex apparatus (male genitalia, endophallus) – see transfer apparatus.

Connate - of two segments or other articulating parts of the body. Not freely moving against one another. There may or may not be a line between them marking the original suture. (emended June 2012)

Copulatory pouch (female genitalia) – see Bursa.

Corbel (leg, tibial apex) – area at apex of tibia defined by setal combs and tibial ridges. The terminology has been and continues to be confused, and relevant terms are summarised here to facilitate understanding. Thompson (1992) recommends the term 'corbel' be abandoned in favour of 'outer bevel' (q.v.) and 'inner flange' (q.v.), an argument based partially on an argument on Latin derivation of the term, since challenged by Oberprieler (in press). Synonyms: Körbichen, corbulae,

Apical setal comb - ring of stout setae around the tibial apex directed distad (q.v.). Anterior and posterior sides of the ring may not be joined, and ring may be incomplete.

Bevel – a slanted area between the tibial apex and the anterior face of the tibia. An ambiguous term.

Inner flange – anterior / distal margin of the tibial socket when raised into a carina or ridge. It may be raised only near the tibial insertion or extend from dorsal to ventral margins of the tibia; it lacks setae. Often it is associated with the uncus, which may have the flange as its base. It can occur on all three tibiae (Thompson, 1992)

Open corbel - apex of tibia with apical setal comb (anterior and posterior) but no secondary comb and no inner flange. The space delimited by the inner flange and the anterior aical setal comb lacks setae or scales. Synonym: simple [tibial apex]; [tibial apex] without corbel.

Enclosed corbel: see True corbel

False Corbel - tibial apex with inner flange raised distal to setal comb (Thompson, 1992). Synonym: Semi-enclosed corbel.

True corbel – more or less flat or concave area at an angle between the tibial apex and the anterior face of the (particularly hind) tibia, demarcated distally by the anterior apical setal comb and proximally by a secondary setal comb (closed or enclosed corbel) or lacking a proximal setal row. The surface may be lacking scales or with scales. Synonym: outer bevel.

Cornu (female genitalia, spermatheca) – the apical, generally curved, more or less acuminate part of the spermatheca.

Coxite (female genitalia) - see Gonocoxite (preferred term)

Coxite-stylus (female genitalia) - term applied to the structure where the stylus and gonocoxite are believed to have fused. Coined by Howden (1992) in discussion of eyeless weevils in the Nearctic and Neotropics. Synonym: vaginal palp.

Deciduous mandibular process (head, mandible) – horn- or blade-like processes on the mandibles of Entiminae some Brachycerinae that almost always fall off in early adult life, leaving a distinct scar. The deciduous process may sit on a pedicel, which is present after the process has fallen off. See extended discussion in Thompson, 1992. Synonyms: pupal mandibles, mandibular appendages, mandibular cusp, deciduous pieces, false mandibles, provisional mandibles, Supplementzähne.

Declivity (elytra) – the posterior part of the elytra that slopes ventrad. Synonym: elytral declivity.

Desmomere (antenna, funicle) – the pedicel and antennomeres that make up the funicle of the antenna (Alonso-Zarazaga, 1989) – see discussion under Antennal segment (preferred term). Synonyms: antennomere, funicle segments, antennal segment. (definition changed 1/06/2012)

Disc (thorax, pronotum) – dorsal portion of the pronotum.

Discrimen (thorax, metathorax) - Median longitudinal depressed line sometimes present on the metaventricle, corresponding with a median ridge internally representing the invagination of the true metasternum. The ridge is usually connected with the metendosternite. May also be visible on the mesoventrite.

Dorsal Adventitious Tooth (leg, tibia) - tooth on apex of tibia dorsally in addition to uncus.

Dorsal Apical Projection (leg, tibia) - dorsal projection on hind tibia lying at the basal end of the apical setal comb when this is ascending.

Dorsal plate (male genitalia) – see Parameroid plate.

Duct-lobe (female genitalia, spermatheca) – the part of the spermatheca from which the spermathecal duct arises. Sometimes extended into a lobe or a tube. Synonym: collum.

Ductus ejaculatorius (male genitalia) – duct between the testes and endophallus. Synonym: ejaculatory duct.

Ejaculatory duct (male genitalia) – see Ductus ejaculatorius.

Elbowed antenna – see geniculate antenna

Elytral declivity see declivity.

Elytral file – see file.

Elytral strigil – see file.

Elytral stria (pl - Elytral striae) (elytra) – longitudinal groove or row of punctures along elytra. There are usually 10, with striae III and VIII often joining posteriorly. The first stria is the one closest to the sutural margin of the elytron. Synonyms: stria.

Elytral submarginal fold (Elytra) – the projecting longitudinal ridge on the ventral surface of the elytra near the costal margin. In the basal third it may be more or less strongly curved away from the elytron inner surface, this character having been used for the Barididae of Zherikhin & Egorov (1990). The fold is likely to function in locking the elytron to the abdomen and metathorax, the fold width and depth basally being related to the size of the metepisternite. Discussed by Davis (2009). Synonyms: epipleural fold, inferolateral flange.

Elytral sutural flange (elytra) - flange on the sutural margin of each elytron ventral to the margin itself. The flange on the right hand elytron fits in the groove between the dorsal surface of the left hand elytron and the flange below it. The margin of the left-hand elytron is wider than that of the right-hand elytron in 'higher' weevils, but about the same in 'lower' weevils. In higher weevils the extension of the flange allows for elytral locking when closed. In cases where the elytra are 'fused' the 'fusion' may be due to the flanges being curled around each other and thus immovable.

Elytro-tergal stridulation system (elytra and abdomen) – the organ responsible for sound production in many higher weevils. It comprises a file – a closely-set patch of parallel ridges on the inner surface of the elytra near the apex or along the sutural margin, and a plectrum – usually a patch or row of tubercles or a single tubercle each side of the mid-line on abdominal tergite 7. The plectrum may be a ridge or pair of ridges in some taxa. In Ithyporina and some Cryptorhynchina the female has the file on tergite 7, derived from the wing-binding patch. The organization of the tubercles and the shape and position of the file can provide useful characters. (Lyal & King, 1996).

Emarginate – notched at the margin. May be used to indicate a concave (in plan) part of a margin.

Enclosed corbel (leg, tibia) – see discussion under corbel.

Endophallus (male genitalia) – membranous sac lying within penis contiguous with ductus ejaculatorius and penis. Sometimes armed with a variety of sclerites (see Transfer apparatus). Everts during copulation. Arzanov (2003) recognized and named a number of lobes of the endophallus. Synonym: internal sac.

Epimeron (thorax) – posterior sclerite of the pleuron discernable on the mesothorax and metathorax in weevils (mesepimeron, metepimeron)

Epipleural bridge (prothorax) – see Hypomerall lobe (preferred term)

Epipleural fold - see elytral submarginal fold (preferred term)

Epipleuron (thorax, elytra) – the inflexed portion of the elytron laterally when the elytra are closed (Torre-Bueno, 1989).

Episternum (thorax) – see Anepisternum (preferred term)

Epistome (head, mouth) – sclerite immediately posterior to the labrum; since the labrum is lost / fused to the clypeus in most adult weevils other than Nemonychidae and Anthribidae the epistome is often used to discuss the sclerite immediately above the buccal cavity without having to decide if it is clypeus or clypeus + labrum fused. It may be fairly clearly demarcated from the dorsal part of the rostrum lying posterior to it (part of the frons) by colour and a ridge or depression (epistomal ridge and epistomal suture respectively) or a transverse row of setae. Laterally there may be a depression (termed the paraclypeal depression by Morimoto & Kojima, 2003), which often has a small tuft of setae (termed the paraclypeal setae by Morimoto & Kojima, 2003). Synonyms: epistoma, clypeus.

Eye-flaps see postocular lobes (preferred term).

False spiculum (male genitalia) – see Spiculum relictum (preferred term)

False strut (male genitalia) – see Spiculum relictum (preferred term)

Fenestrae (male genitalia) – unsclerotised area of the parameroid plate (q.v.) bounded by sclerites (see suprafenestral sclerite and subfenestral sclerite). Homology across Curculionoidea is unlikely. (see Wanat, 2007).

Fifth tarsomere (leg, tarsus) - the distal tarsomere, bearing the tarsal claws. Synonyms: unguitractor, onychium.

File (elytra – elytro-tergal stridulation system) – patch of fine, parallel ridges on the internal surface of the elytra, generally near the apex, which form part of the elytro-tergal stridulation system. Synonyms: elytral file, elytral strigil, pars stridens, stridulatory rasp, stridulatory file.

First connecting membrane (male genitalia) - see Post-tegmina membrane. The First connecting membrane of Clark (1977) is the Pre-tegmina membrane.

Flagellum (male genitalia) – long tubular sclerite sometimes present within the endophallus with the gonopore at its apex. In some cases (more rarely) the flagellum is one or a pair of rods within the ejaculatory duct but outside the endophallus. The two structures are not homologous. See also ‘Transfer apparatus’. See Wanat, 2007.

Forceps (male terminalia) – externally-visible processes from 8th sternite in the amycterine *Phalidura* (Thompson, 1992).

Frena (male genitalia, endophallus) – paired sclerites in endophallus, situated sub-basally when endophallus everted, comma-shaped or hook-shaped. Kuschel (1995) suggested their presence is a plesiomorphic character state in Curculionoidea. (Wanat, 2001, 2007).

Frons (head, head capsule) – The area dorsally on the head anterior to the antennal insertions and posterior to the epistoma, where visible. The term is sometimes used for the area between the eyes, which might better be termed ‘forehead’. (Davis 2011)

suggested in Sitophilus that it extends ventrally and surrounds the scrobe. See discussion under head. [Emended June 2015]

Funicle (head, antenna) - the middle part of the antenna, lying between the scape and the club, comprising the pedicel and up to 6 'desmomeres' (antennomeres) ('pedicel' and 'flagellum' are terms rarely if ever used in weevil taxonomy).

Furcal arms (male genitalia) – see basal arms (of spiculum gastrale).

Furcasternum - see Sternellum (preferred term).

Gena (head) – the part of the head capsule on each side below the eye, bordered dorsally by the frontal suture, posteriorly by the occipital suture, ventrally by the subgenal suture. See also Head.

Geniculate antenna (head, antenna) - the state in higher weevils where the pedicel (basal segment of the funicle) inserts subapically on the scape, causing the funicle to be held at an angle to the scape. cf. subgeniculate antenna. Synonym: elbowed antenna.

Genital chamber (male genitalia) – see Genital pocket

Genital pocket (male genitalia) – membranous invagination between tergum VIII and sternum VIII and to which the spiculum gastrale is attached. It includes both genital opening and rectum. Wanat (2007) considers the pocket to extend from SVII, since SVIII is withdrawn dorsal to SVII at rest. Wanat also places the anterior end of the genital pocket at the base of the apodemes of the penis. Synonyms: genital chamber, genital sheath of Wanat 2007. [Emended June 2015]

Genital sheath (male genitalia) – see Genital pocket

Genital spiculum (female genitalia) - see Spiculum ventrale

Genital tract (female genitalia) – the vagina and bursa.

Gland-lobe (female genitalia, spermatheca) - see discussion under Spermatheca (ramus)

Gonatoceri (informal group term) – name applied to a grade of Curculionoidea with a geniculate antenna and male genitalia lacking a separate tectum (q.v.), broadly equivalent to the Curculionidae of Thompson (1992) which are characterised by this state. See also Gonatoceros, Orthoceros, Orthoceri, Heteroceri.

Gonatoceros (antenna) – see geniculate (preferred term). Term now also used to describe a group of weevils (see below) and as a 'shorthand' to describe typical character states (e.g. 'gonatoceros genitalia').

Gonatoceros (male genitalia) – see pedal genitalia (preferred term).

Gonocoxite (female genitalia) – one of a pair of conical sclerites apically on ovipositor, associated with segment IX and between which the female vulva opens. See also 'coxite-stylus'. Synonyms: hemisternite, coxite. May be proximal and distal elements – see Lanteri & del Rio, 2008

Gonopore (female genitalia) – the opening of the common oviduct into the vagina.

Gonopore (male genitalia) – the opening through which the sperm are transferred to the female genitalia on eversion of the endophallus. This may simply be the junction of the ductus ejaculatorius with the endophallus, at the tip of the flagellum, where present. Synonym: phallotreme.

Gula (head, head capsule) - the sclerite lying between the gular sutures, where these are separate (Lyal, 1995)

Gular cavity (head) – see mouth cavity.

Gular suture (head, head capsule – rostrum) - suture or pair of sutures extending from the posterior margin of the head capsule to the posterior tentorial pit(s). The sutures are often fused to a single suture in weevils. (Lyal, 1995)

Head (adult) – The head of the adult weevil has a more or less bulbous head capsule (q.v.) which accommodates the muscles of the mouthparts, and a variably elongate rostrum (q.v.) bearing the antennae and mouthparts and through which the tendons of the mandibles and maxillae pass. The various sclerites of the head are fused together and, particularly dorsally, sutures obscured; there are few reliable landmarks. Dorsal structures: The Vertex is the posterior part of the head, extending from the occiput to the level of the antennal insertions. Behind the eyes this may be called the Temple, between the eyes the Forehead, and between the anterior margins of the eyes and the level of the antennal insertions it may be called the Epifrons. Anterior to the epifrons the Frons may be separated by a groove, or the junction between the two simply be marked by some longer setae, at least laterally; the frons itself may lack setae or scales. Anterior to the frons the Epistome may be slightly lower or at a different angle to the frons, and may be fringed by a sparse row of setae posteriorly. The epistome and frons may be united as a distinct Nasal plate (=nasal plaque, plaga).

Lateral and Ventral structures: The ventral structures were discussed extensively by (Lyal 1995). See also Gena, Gula, Gular suture, Cervical sclerites, Hypostoma, Interocular pit, Submentum, Venter

Head capsule (head, adult) – the head apart from the rostrum. See more detailed discussion under Head.

Hemisternites (female genitalia) – see Gonocoxite (preferred term).

Hemisternite (male terminalia) – one of a pair of sclerites on sternum VIII

Heterocerri (informal group term) – name applied to a grade of Curculionoidea characterised by possession of geniculate antennae (q.v.) and orthocerous genitalia (q.v.). Comprises Dryophthoridae, Eirrhinidae, Nanophyidae, Brachyceridae. See also Gonatocerri, Orthocerri.

Horn sheath (prothorax) – simple or bifurcate tube-like invagination between prothoracic horns of some Baridinae and Conderinae. (Davis, 2009). Synonym: sheath.

Humeral angle – see humerus

Humeral callus (mesothorax, elytra) – projection on humerus; may be used as simply another term for humerus.

Humerus (mesothorax, elytra) – anterolateral angle of elytron, sometimes produced. Tends to be rounded in apterous species. If there is a projection some authors term this a 'humeral callus' or a 'humeral umbone' (for a knob-like projection). Synonyms: humeral angle, humeral callus.

Hypomerale lobe (prothorax) - the sclerotised area behind the procoxae formed from pleural (hypomerale) projections, meeting in the midline and abutting anteriorly medially on the sternellum. Synonyms: pleural process; post-coxal bridge; post-coxal process; epipleural bridge; hypomeron.

Hypomeron - see Hypomerale lobe (preferred term).

Hypostoma (head) – ventral part of the head bearing the posterior articulation of the mandible (postcoila). In most weevils (not Scolytinae, Platypodinae) the hypostoma projects anteriorly as a hypostomal process (q.v.)

Hypostomal process (head) – Anterior process of the hypostoma bearing the lower (posterior) mandible articulation and thus projecting below the mandible and partially enclosing the pleurostomal sinus (q.v.). Synonyms: hypostomal tooth, postgenal arm

Hypostomal sinus (head) – The notch between the hypostoma (q.v.) and the labium, in which the maxillae can be seen. (Morimoto & Kojima, 2003). [Emended June 2015]

Hypostomal sulcus (head, rostrum) – part of subgenal sulcus posterior to the posterior mandibular articulation.

Hypostomal tooth – see hypostomal process.

Inferolateral flange (elytra) - see elytral submarginal fold (preferred term)

Inner flange (leg, tibia) – anterior / distal margin of the tibial socket when raised into a carina or ridge. It may be raised only near the tibial insertion or extend from dorsal to ventral margins of the tibia; it lacks setae. Often it is associated with the uncus, which may have the flange as its base. It can occur on all three tibiae (Thompson, 1992). See discussion on the Corbel. [emended June 2015]

Integument - exoskeleton, particularly outer covering of entire insect.

Intercoxal process (abdomen) – portion of ventrite 1 extending anteriorly between hind coxae.

Intercoxal process (prothorax) – posterior extension of prothoracic sternum (basisternum) between fore coxae, meeting the sternellum if coxae sufficiently separated.

Interocular pit (head) – abrupt depression on dorsum of head, generally between eyes but sometimes more distally placed on the rostrum. This marks the junction of the dorsal tentorial arms with the head capsule (Davis 2011).

Internal sac (male genitalia) – see endophallus (preferred term).

Interstice see Interstria (preferred term).

Interstria (elytra) – raised longitudinal strip on elytra between striae. The interstria nearest the sutural margin is interstria I (Lawrence & Britton, 1991). Some authors (e.g. Vanin & Reichardt, 1976) numbered the interstriae in a different way, for example the first being called the sutural interval, the second as interstice I, the third as interstice II, and so on (Vanin, 2008). Synonyms: interval, interstice.

Interval see Interstria (preferred term)

Introse - with teeth on the apical margin of the mandible and the mandibles not opening beyond the lateral margins of the rostrum.

Joints of funicle see antennal segment.

Labium (head, mouthparts) – plesiomorphically in Coeloptera the labium comprises a tripartite prementum (bearing the labial palps), mentum and submentum. Curculionoidea have a prementum but the mentum and submentum are not visible as separate sclerites. The single sclerite may be known as the Postmentum (e.g. (Morimoto and Kojima 2003) or the submentum (e.g. Lyal, 1955). Davis (2011) refers to the projecting portion of this sclerite supporting the prementum as the postmentum and the region between this and the posterior tentorial pits as the submentum; earlier authors termed this projecting part the peduncle.

Labral rods (larva, mouthparts) see tormae (preferred term)

Lateral lobes (male genitalia) – see parameroid lobes.

Latero-ventral sulci (head, rostrum) - (Wanat, 2001)

Lenticular – lens-shaped

Ligula (head, mouthparts) –lobe of the labium, projecting anteriorly of the mentum. Plesiomorphically this comprises the glossae and paraglossae, although as Crowson (1981) states these are absent in beetles the homology of the structure is unknown. [Emended June 2015]

Ligula (male genitalia) – see Ostiolar sclerite.

Mandibles (head, mouthparts) - In various forms across the suborder. Terms used may include: Extrose - with teeth on the external face, the mandibles rotating away from the midline and opening away from the midline of the rostrum and not meeting apically. Introse: Teeth on the anterior/interior face, meeting when mandibles closed (the more common situation). Plurisetose: numerous setae, often of similar length on the outside face of the mandible. Paucisetose: only a few setae present on the mandible. [emended June 2015]

Mandibular cusp (head, mouthparts) – see Deciduous mandibular process

Mandibular scar (head, mouthparts) – circular or oval scar left on mandible after loss of the deciduous mandibular process (q.v.).

Manubrium (male genitalia) – See Tegminal apodeme.

Median lobe – see Penis (preferred term)

Median Lobe Apodemes (male genitalia) – see Penis Apodemes (preferred term).

Median notch (male genitalia) – term applied to gap between parameroid lobes (Wanat, 2007)

Median oviduct (female genitalia) – the ectodermal oviduct opening into the bursa-vagina complex, often near the spermathecal duct. Synonym: common oviduct.

Median struts (male genitalia) - see Penis Apodemes (preferred term).

Mentum (head) – middle plate of the labium, not visible as a separate plate in adult Curculionoidea.

Mesanepisternum (thorax, mesothorax) – anterior pleural sclerite of the mesothorax. Synonym: Mesepisternum. (definition changed June 2012).

Mesendosternite (thorax) - A pair of internal apodemes formed by invaginations within the mesocoxal cavities, probably homologous with the furca.

Mesepimeron (thorax, mesothorax) - Posterior pleural sclerite of the mesothorax.

Mesepisternum (thorax, mesothorax) – see Mesanepisternum (preferred term)

Mesonotum (thorax, mesothorax) – the dorsal sclerite of the mesothorax.

Mesorostrum (head, rostrum) - the section of the rostrum widened above the antennal insertion. If the broadening is absent, the term refers to the level of the antennal insertion. (Wanat, 2001). Purely a descriptive term lacking any morphological significance.

Mesosternal canal see Mesoventral canal.

Mesoventral canal (mesothorax) – channel on the mesosternum to receive the rostrum when the head is rotated down and the rostrum lies along the ventral surface of the thorax. Found in some groups currently within the Cryptorhynchinae and Molytinae. (cf mesoventral receptacle). Synonyms: mesosternal channel; mesosternal canal. [Emended June 2015]

Mesosternal channel see Mesoventral canal.

Mesosternal cup see Mesoventral receptacle.

Mesosternum see mesoventrite.

Mesothoracic receptacle see Mesoventral receptacle.

Mesoventral receptacle (thorax, Mesoventrite) – cup-shaped depression of the mesothorax that accepts the tip of the rostrum if this is bent between the front legs. It may be in the form of a full cup with lateral walls and a posterior wall, or with lateral walls only. Not homologous across Curculionoidea, although often used as an apomorphy for

Cryptorhynchinae. Synonyms: mesosternal cup; mesothoracic cup; mesothoracic receptacle; pectoral receptacle. [Emended June 2015]

Mesoventrite (thorax, mesothorax) - ventral sclerite lying in front of and between mesocoxal cavities, comprised of the fused preepisternum and basisternum (according to Wanat, 2001) or preepisterna and katepisterna (according to Lawrence et al, 2010). Generally referred to as the mesosternum in weevil taxonomic literature but the true mesosternum is invaginated. Synonym: Mesosternum.

Metanepisternal suture (metathorax) - suture between metaventrite and metepisternum. In cross-section metaventral part of suture is cup-shaped, holding the metepisternum and allowing it to flex during flight. In some higher weevils sclerolepidia are present along the suture, arising from the metaventrite. The suture may be lost in flightless weevils. (Lyal et al., 2006). Synonyms: anapleural suture, metepisternal suture, metathoracic pleurosternal suture.(term changed March 2012).

Metanepisternum (metathorax) – anterior pleural sclerite of the metathorax, lying laterad to the metaventrite and mesoventrad to the metepimeron. Often partially and sometimes completely hidden by elytra. Synonym: metepisternum. [Emended June 2015].

Metanotum (thorax, metathorax) – the dorsal sclerite of the metathorax.

Metarostrium (head, rostrum) - basal part of rostrum, from base to widening above antennal insertion (Wanat, 2001). Purely a descriptive term lacking any morphological significance.

Metasternal canal (metathorax) - see Metaventral canal

Metasternum see metaventrite.

Metathoracic pleurosternal suture see metanepisternal suture.

Metaventral canal (metathorax) - channel on the metasternum to receive the rostrum when the head is rotated down and the rostrum lies along the ventral surface of the thorax. Found in Aedemonini and Sophrorhinini. Synonym: metasternal canal. [Emended June 2015]

Metaventrite (thorax) - ventral sclerite lying behind and between mesocoxal cavities, comprised of the fused preepisterna and katepisterna (Lawrence et al, 2010). Generally referred to as the mesosternum in weevil literature but the true mesosternum is invaginated along the midline (see Discrimen). Synonym: metasternum.

Metendosternite (metathorax) – internal sclerite of the metathorax. It usually consists of a median stalk, two short to long lateral ‘furcal arms’ and a ‘longitudinal flange’ anterior to and between the arms from which a pair of tendons arise. There are often phylogenetically-important characters associated with this.

Metepimeron (thorax) – Posterior pleural sclerite of the metathorax, positioned laterad of and above the metanepisternum and mostly or completely concealed by the elytra. It may be fused to the metanepisternum.

Metepisternal suture see Metanepisternal suture (preferred term).

Metepisternum see Metanepisternum (preferred term)

Mouth cavity (head) – opening at the front of the head accommodating the mandibles, maxilla and labium. Synonyms; Buccal cavity, gular cavity.

Mucro (pl - mucrones) (leg, tibia) – tooth-like process on the inner (ventral) apical angle of the tibia, but not linked to the inner flange (q.v.). Proposed by Thompson (1992) to be homologous with the uncus (q.v.) following fusion with the inner flange. See also 'pre-mucro'. In many Apionidae and Nanophyidae it is present in males but not females. (Emended December 2012)

Mycangium (pl - mycangia) – structures on the body used for the transport of symbiotic fungi. Found in a number of weevils, particularly Scolytinae, and have clearly evolved more than once.

Nasal plaque – see Nasal plate.

Nasal plate (head) – plate formed from the fused frons and epistome, where demarcated. See discussion under head. Synonyms: nasal plaque, plaga.

Nodulus (female genitalia, spermatheca) – the basal part of the spermatheca, between the cornu, ramus and collum. [Emended June 2015]

Noto-sternal suture (prothorax) – suture, often obscured, between the notal and sternal components of the prothorax, extending anteriorly from the fore coxae externally. Synonym: pleuro-sternal suture.

Ocular lobe (prothorax) – see postocular lobe (preferred term).

Onychium – see fifth tarsomere

Open corbel (leg, tibia) – see discussion under Corbel

Orthoceri (informal group term) – name applied to a grade of Curculionoidea with Orthocerous antennae (q.v.) and male genitalia of the plesiomorphic 'orthocerous' type (q.v.). These are the basal weevils on the generally-accepted cladograms. See also Heteroceri, Gonatoceri.

Orthocerous (antenna) – straight, lacking an 'elbow' between the scape and funicle. cf 'geniculate antenna'.

Orthocerous genitalia (male genitalia) – see pedotectal genitalia (preferred term)

Ostiolar sclerites (male genitalia) – pair of sclerites on dorsal wall of penis adjacent to ostium. Synonym: ligula.

Ostium (male genitalia) – opening of penis through which the endophallus everts.

Outer bevel (leg, tibia) – The slope on the anterior face of the tibia between the anterior apical margin (as defined by the anterior apical setal comb) and the unmodified anterior face, delimited by a transverse swelling or ridge just basal to the tibial apex bearing with a secondary setal comb. Almost always on the hind tibia only (Entiminae, Brachycerinae). See discussion on the Corbel.

Ovipositor (female genitalia) – the organ by which the eggs are laid, including gonocoxites, styli and vulva. (changed June 2012).

Pappolepida (scales) – scales which divide multiple times at apex to give them a velvety or feathery appearance (Brown, 2015 (PhD); George et al, 2015)

Paracoila (head, mouthparts) – pouch for receiving the cardo of the maxilla near the base of the hypostomal sinus (Morimoto & Kojima, 2003).

Parameres see parameroid lobes (preferred term).

Parameral lobes (male genitalia) – see parameroid lobes.

Parameroid lobes (male genitalia) - lobes attached to the posterodorsal part of the tegminal ring. May have posteriad unsclerotised extensions termed ‘apical membranous lobes’. The parameroid lobes are not unequivocally homologous with the parameres of other Coleoptera, hence the difference in terminology. See also parameroid plate. Synonyms: parameres, parameral lobes.

Parameroid plate (male genitalia) - dorsal section of tegmen extending posteriorly. Generally comprising the parameroid lobes or a single lobe, sometimes with additional sclerites or showing membranous areas surrounded by sclerite. It is a two-layered structure, and sometimes has different sclerites on the dorsal and ventral layers. The parameroid lobes are not unequivocally homologous with the parameres of other Coleoptera, hence the difference in terminology. See parameroid lobes, fenestrae, suprafenestral sclerites, subfenestral sclerites, prostegium. Synonyms: dorsal plate, tegminal plate, cap-piece.

Pars stridens see File.

Pectoral canal see Rostral canal

Pectoral receptacle - see Mesothoracic receptacle

Pedal genitalia (male genitalia) - the state found in majority of ‘Gonatoceri’, where the tectum (q.v.) is not visible, the penis body and its apodemes describe two separate curves when viewed laterally and are articulated with each other, the tegmen has two more or less distinct parameroid lobes. Compare with ‘pedotectal genitalia’ (q.v.). Synonym: gonatoceros genitalia.

Pedicel (head, antenna) - second of the three segments of the basic insect antenna (the other two being the scape and flagellum). ‘Pedicel’ and ‘flagellum’ are terms rarely if ever used in weevil taxonomy.

Pedon (male genitalia) – ventral plate of penis. See also pedotectal genitalia and tectum.

Pedotectal genitalia (male genitalia) – term describing the type of penis found in basal ‘orthoceros’ weevils. The dorsal and ventral plates of the penis body (‘tectum’ and ‘pedon’ respectively) are separate for most of their length, united by a membrane, and fused at the base (c.f. tectum not visible), the apodemes and the penis body describe a smooth curve in profile (cf a double curve). The parameroid lobes are more or less fused together (cf separate). Synonym: orthoceros genitalia.

Peduncle (head) – the part of the submentum (q.v.) projecting between the maxillae and bearing the prementum (q.v.). See discussion under labium.

Penis (male genitalia) – the part of the male genitalia comprising the unpaired intromittent organ with its apodemes, and containing the eversible endophallus and the opening of the ductus ejaculatorius (gonopore). It is enclosed by the tegmen. The ‘body’ and apodemes (= temones, aedeagal apodemes) are sometimes compared in length as a taxonomic character. Synonyms: median lobe; ‘aedeagus’ of some authors, including Wanat 2007.

Penis apodemes (male genitalia) – a pair of apodemes extending anteriorly from the base of the penis. In more detail: invaginations of the post-tegmenal membrane (first connecting membrane) lying anterior to the penis body, and generally connected to it by sclerotization. The apices of the apodemes support a pair of longitudinal muscles arising dorsolaterally on the tegmenal ring and a pair arising on the apex of the tegmenal apodeme; these serve as protractors of the penis (Burke, 1959, Cerezke, 1964), and possibly also retractors (Cerezke, 1964). Synonyms: aedeagal apodemes, median lobe apodemes, median struts, temones (in some cases the term ‘temones’ is applied to tegmenal apodemes). [emended September 2015]

Penis body (male genitalia) – the part of the penis not including the apodemes.

Pennon (male genitalia) – fine filamentous sclerotized process within the genital pocket, fringing semi-permanent folds (visible when stained with chlorazol black). (Wanat, 2001, 2007).

Phallobase (male genitalia) – Anterior or basal part of the aedeagus from which the parameroid lobes or parameroid plate arise. See Tegmen.

Phallotreme (male genitalia) – see Gonopore. Sometimes applied to ostium, incorrectly.

Phanerognatha (informal group of weevils) – Curculionidae characterised by the phanerognathous (qv) character state, often also known as ‘long-nosed’ weevils, and also characterised broadly by a long slender rostrum; the mandibles lack a deciduous process, the larval antenna has a more or less conical and projecting apex, and the larvae are concealed within plants. The group is not monophyletic, and has no formal status. See also ‘Adelognatha’.

Phanerognathous (head, mouthparts) – maxillae not covered by the prementum. (cf ‘adelognathous mouthparts’. Characteristic of the ‘Phanerognatha’ (qv).

Plaga – see Nasal plate.

Plectral tubercles (abdomen, tergite VII) – two or more raised tubercles on tergite VII, each with a small seta posteriorly, that act against the stridulatory file in the elytra-tergal stridulatory system (q.v.) (Lyal & King, 1996)

Pleural process - see Hypomerall lobe (preferred term).

Pleuro-sternal suture (prothorax) – see Noto-sternal suture

Pleurostomal sinus (head) – emargination of buccal cavity margin between precoila and postcoila to accommodate the base of the mandible and allow it to rotate outside the margins of the rostrum. Shallow in Attelabidae, Scolytinae and Platypodinae, deeper in other weevils (Morimoto & Kojima, 2003).

Pleurostomal sulcus (head, rostrum) – part of subgenal sulcus dorsal to the mandible.

Postcolia (head) – the socket for the ventral (posterior) articulation of the mandibles (postartis), born on the hypostoma (q.v.). (Morimoto & Kojima, 2003) [Emended June 2015]

Post-coxal bridge - see Hypomerale lobe (preferred term).

Post-coxal lamellae (prothorax) – blade-like projections behind fore-coxae, extending between coxa and posterior margin of prothorax, as a part of the sternal canal for the rostrum. Apomorphic for Curculionidae: Cryptorhynchinae: Aedemonini, and also for Aonychus (Erihrhini). Similar structures, anteriorly abutting the coxa and posteriorly meeting medially are found in the Camptorhinini (currently in the Cryptorhynchinae but being transferred to the Molytinae, and itself polyphyletic). Synonyms: post-coxal flanges, post-coxal walls of the pectoral canal.

Post-coxal process (prothorax) – see Hypomerale lobe (preferred term).

Posterior declivity (elytra) – distal part of elytra where, at rest, sutural margin curves ventrad towards lateral margin.

Posterior tentorial pit (head) – invaginations of the cuticle on the ventral surface of the head corresponding to the positions of the posterior tentorial arms internally. The pits mark the junction of the gular suture and the subgenal suture. They may be very difficult to see in weevils, and may be fused to a single pit or extend effectively along a single gular suture. See Lyal (1995) for a discussion.

Postgenal arm (head) – see hypostomal process.

Postgula (head) – see cervical sclerites (preferred term).

Postmentum (head) – see submentum (preferred term). Davis (2011) limits the postmentum to the projecting part of the labium supporting the prementum (peduncle, q.v.), terming the posterior part the submentum. See discussion under labium q.v. [Emended June 2015]

Postocular lobe (prothorax) – lobulate projection on the anterior margin of the pronotum, lying just behind the eye and covering it when the head is rotated ventrad. Often fringed with setae, probably having as an eye-cleaning function. Synonyms: ocular lobe; eye-flap. [Emended June 2015]

Post-tegmina membrane (male genitalia) – membrane between tegmen and base of penis (Wanat, 2007). Synonym: first connecting membrane.

Praemucro (leg) - see Premucro (preferred term)

Precoila (head, mouthparts) – the socket for the dorsal (anterior) articulation of the mandibles (preartis), born on the clypeus. The clypeus in weevils is not separated from the frons by a suture, so cannot be distinguished.

Pregula (head) – see Submentum

Prementum (head, mouthparts) – the anterior plate of the mentum, bearing the palpi. See discussion under Labium.

Premucro / pre-mucro (leg, tibia) – small preapical tooth arising ventrally (occasionally distally) at apex of tibia (Kuschel, 1951, Thompson, 1992), basal to mucro/uncus. It can co-exist with a mucro or uncus (i.e. is homologous with neither). It is marked by a paired tuft of long curved setae arising either side of its base and curving more or less over it. In some groups (e.g. many Lixinae, some Molytinae) it is more strongly-developed in females than males, and may be absent or very small in the latter. Synonyms: praemucro, pre-mucro, secondary mucro. (Emended December 2012)

Pretarsus (leg) – terminal segment of the leg, consisting of paired or, more rarely, single claws. See Tarsal Claw. (definition changed June 2012).

Pre-tegmina membrane (male genitalia) – membrane between 9th segment and the tegmen in males (Wanat, 2001, 2007) Synonym: second connecting membrane.

Pronotum (thorax, prothorax) – the dorsal sclerite of the prothorax.

Prorostrum (head, rostrum) - the part of the rostrum from the anterior end of the mesosrostrum (broadening above the antennal insertion) to the apex of the rostrum (Wanat, 2001). Purely a descriptive term lacking any morphological significance.

Prostegium (male genitalia) – sclerite in ventral layer of parameroid plate to which the arms between the tegmina apodeme and the parameroid lobes attach. Present on 'orthocerous' weevils and forming part of tegmina ring in more derived groups. (Wanat, 2007; Kuschel, 1989). Synonym: supra-annular sclerite.

Prosternal canal (prothorax) – channel on the prosternum (basisternum and sternellum) to receive the rostrum when the head is rotated down and the rostrum lies along the ventral surface of the thorax. It may extend between the separated coxae onto the sternellum or stop at the fore coxae, thus lying along the basisternum only. See also Rostral canal.

Prosternal process (prothorax) – the part of the prosternum lying between the fore coxae (if these are separated) and extending to the hind margin of the prosternum unless interrupted by the hypomerall lobes. Comprising an anterior basisternum and a posterior sternellum which, if they meet, are separated by the sternacostal suture.

Prosternum (prothorax) – the sternal element of the prothorax, comprising an anterior basisternum and a posterior sternellum which are either separated by contiguous procxae or, if they meet, are separated by the sternacostal suture.

Prothoracic horns (prothorax) – pair of short or long projections generally arising just anterior to fore coxae in some Baridinae and Conoderinae, used by males in fighting.

Proventriculus (alimentary canal) – posterior part of the foregut comprising eight longitudinal folds, variously sclerotised in some groups. See Kissinger, 1963 and Calder, 1989

Proximal comb (leg, tibia) – see secondary comb.

Pterygium (head, rostrum) - One of the two lateral expansions on the rostrum lying above and partly concealing the scrobes in which the antennae are concealed.

Pygidium (abdomen) – One or more of the apical abdominal tergites which are exposed by the elytra at rest and are more highly sclerotised and strengthened than the other tergites. Thompson (1992) uses a ‘tergal formula’ (TF) in which tergites concealed by the elytra are given in normal type ‘1-6’, those forming the pygidium in bold type ‘7’ and those concealed in the genital chamber in parentheses ‘(8)’.

Ramus (female genitalia, spermatheca) – the part of the spermatheca to which the spermathecal gland is attached. See Gland-lobe. [Emended June 2015]

Rectal loop – oblique sclerotized band surrounding the rectum, functioning as part of the cryptonephridial system (Lyal & Favreau, 2015)

Rectal ring – transverse sclerotized ring surrounding the rectum, functioning as part of the cryptonephridial system (Lyal & Favreau, 2015)

Rectal valve – sclerotized ring or loop surrounding the rectum, functioning as part of the cryptonephridial system (Lyal & Favreau, 2015)

Rostral canal (thorax) - channel on the thoracic sterna and ventrites to receive the rostrum when the head is rotated down and the rostrum lies along the ventral surface of the thorax. On the prothorax it is generally delimited by longitudinal carinae, probably marking the pleuro-sternal suture. Synonym: pectoral canal, rostral furrow, prosternal canal. [Emended July 2015]

Rostral furrow see Rostral canal (preferred term).

Rostrum (head) – the anterior part of the head, bearing the mouthparts, extended into a tube in most weevils, but may be short or absent (e.g. Scolytinae). See more detailed discussion under Head. Wanat (2001) considers the rostrum as comprising a ‘Metarostrium’ (q.v.), ‘Mesorostrum’ (q.v.) and ‘Prorostrum’ (q.v.) (base, middle and distal) but these terms are purely descriptive and do not relate strictly to morphological homologies. Synonyms: beak, snout. [Emended June 2015]

Scape (head, antenna) – the basal segment of the antenna.

Sclerolepidia (metathorax) – specialized scales arising from metaventricle along the metanepisternal suture (Lyal & King, 1996) (Emended June 2015)

Scrobe (head, rostrum) - The linear depression extending generally from the antennal insertion towards the head capsule (although sometimes deflected ventrad and occasionally with an anteriad element) into which the antennal scape fits when withdrawn against the rostrum. If shallow they are sometimes termed ‘open’. [Emended June 2015]

Scutellar shield (mesothorax) – the part of the mesoscutellum which is exposed between the bases of the elytra. In most weevil literature this is just referred to as ‘scutellum’ (q.v.).
Synonym: scutellum (in part).

Scutellum (mesothorax) – Posterior part of mesotergum. The term is often used to refer only to the portion of the scutellum that is visible between the bases of the elytra (see ‘Scutellar shield’). Appropriate terminology is ‘visible’ vs ‘concealed’, not ‘present’ vs ‘absent’.

Second connecting membrane (male genitalia) – see Pretegmenal membrane.

Secondary mucro – see premucro.

Secondary (tibial) comb (leg, tibia) – row of stout setae subapically on dorsal surface of the tibia just basal to the apical comb. It may join the apical comb to produce the ‘corbel’ (q.v.)

Sheath – see Horn Sheath.

Spermatheca (female genitalia) – Sac-like structure, usually sclerotised in weevils, attached by the Spermathecal Duct to the genital tract and with a Spermathecal Gland. The spermatheca may be considered as having several different areas: Gland-lobe (= ramus) from which the Spermathecal Gland arises; Duct-lobe (= collum), where the Spermathecal Duct attaches, nodulus (basal part of the spermatheca on which the gland-lobe and duct-lobe arise, cornu, the apical often curved and acuminate part. A spermathecal compressor muscle may extend between the cornu and nodulus.[Emended January 2016]

Spermathecal duct (female genitalia) - duct between the Spermatheca and the Genital Tract, usually meeting it at the junction between the Common Oviduct and the Bursa. Variation in the position of this attachment can be useful systematically.

Spermathecal Gland (female genitalia) – gland attached to the spermatheca. After maceration the gland is largely lost, but the sac-like lumen of the gland remains, and is usually termed the ‘spermathecal gland’ in taxonomic descriptions.

Spicule plates (male genitalia) – sclerites in the second connecting membrane of the genital pocket adjacent to the base of the spiculum gastrale. In at least some cases likely to be homologous to the basal plate of the spiculum gastrale, although do not always have muscle attachments (see (Clark 1977))

Spiculum gastrale (male genitalia) – Internal apodeme of male sternite IX. Comprises a long apodeme (or manubrium) and a posterior sclerotized plate or pair of arms (basal arms). Very variable through Curculionoidea. The basal arms are fused to the genital pocket. Although derived from Sternite IX, it may not be wholly homologous with that sternite. Synonyms: urosternite, sternite IX. (edited 25/11/2011).

Spiculum relictum (male terminalia) - small pouch or apodeme sometimes invaginated between hemisternites on male sternum VIII (Thompson, 1992). Possibly serially homologous with the apodeme of sternum IX. Synonyms: false strut, false spiculum.

Spiculum ventrale (female terminalia) – internal apodeme of sternum VIII. Comprises a pair of basal arms, sometimes united to form a basal plate, and an apodeme, sometimes

subsumed by the basal arms to give a V-shaped or U-shaped structure. Erbey et al (2010) termed the apodeme the 'genital spiculum'. Synonym: genital spiculum. (modified June 2012)

Spinasternum (prosternum) see Sternellum.

Sternacostal suture (thorax) – suture between basisternum and sternellum. (Davis, 2009)

Sternellum (thorax) - the second sclerite of the ventral part of each thoracic segment. In weevils it is generally distinct on the prothorax, forming for example the posterior part of the canal in Cryptorhynchinae and a small clear sclerite just behind and between the fore coxae in most others. On the mesothorax it is rarely if ever distinct, but would lie between the mid coxae. On the metathorax it is small and lies between the hind coxae, if discernible. Wood (2007-14), after Hopkins, 1909) figures the sternellum for each thoracic segment. The sclerite on the prothorax has also been called the 'furcasternum' and 'spinasternum'; homologising sclerites in this complex area is difficult. See the discussion on Characters and character-states: "Sternellum or Prosternal process" for further information and a terminological issue. Synonyms: 'furcasternum', 'spinasternum'.

Sternite IX (male genitalia) – see spiculum gastrale

Sternite VIII (female genitalia) – see spiculum ventrale

Striae – see elytral striae.

Stridulatory File – see file.

Stridulatory rasp – see file

Stylus (female genitalia) - a distal segment borne by the Gonocoxite (q.v.) in most species. Generally bears setae, and rarely 'lost' or fused to the Gonocoxite (see 'coxite-stylus').

Subfenestral sclerites (male genitalia) – sclerotised transverse sclerite adjacent to parameroid lobe anterior to an unsclerotised area (fenestrae (q.v.) of the parameroid plate (q.v.). Homology across Curculionoidea is unlikely. (see Wanat, 2007).

Subgenal sulcus (head, rostrum) – sulcus on the ventral surface of the adult weevil head running between the anterior tentorial pits to the posterior tentorial pits. Synonyms: subgenal suture; pleurostomal sulcus + hypostomal sulcus.

Subgenal suture (head, rostrum) – see Subgenal sulcus (preferred term).

Subgeniculate antenna – state in some Apionidae where the apical part of the scape is sharply curved, causing the funicle to lie at an angle to the long axis of the scape, giving the impression of the geniculate state (q.v.) (Wanat, 2001)

Submarginal fold (elytra) – longitudinal furrow and carina internally on elytron close to costal margin, widening basally. The deeper wider "Basal pocket" of the fold fits over the raised sides of ventrite 1 + 2, and the metepisternite, the posterior part locks over the other abdominal ventrites. The particularly broad basal part has been used as a character of the Baridinae + Conoderinae + Ceutorhynchinae.

Submentum (head, rostrum) – the posterior plate of the labium, posteriorly meeting the gula at the level of the posterior tentorial pits (q.v.) and anteriorly meeting the prementum. Within the Curculionoidea there is no separate mentum (median plate of the labium) but the submentum cannot be unequivocally synonymised with the submentum of other Coleoptera (where the mentum is present) or the more plesiomorphic postmentum of other insects. Davis (2011) distinguishes a postmentum (the anteriorly projecting part of the fused mentum + submentum) and the submentum, which lies between that and the posterior tentorial pits. Discussed by Lyal (1995), Morimoto & Kojima (2003) and Davis (2011). See discussion under Labium. Synonyms: postmentum; pregula.

Supra-annular sclerite (male genitalia) – see Prostegium.

Suprafenestral sclerites (male genitalia) – sclerotised transverse sclerite adjacent to parameroid lobe posterior to an unsclerotised area (fenestrae (q.v.) of the parameroid plate (q.v.). Homology across Curculionoidea is unlikely. (see Wanat, 2007).

Tarsal groove (leg, tibia) – furrow on the apico-ventral surface to receive the tarsus if folded back along the leg.

Tarsal Claw (leg, tarsus) - paired claws at the apex of the tarsus. They may be simple (more or less gradually tapering from base to apex and lacking teeth or other projections), appendiculate (with a ventral tooth or other projection arising in the basal third, often arising from a process apparently splitting off from the claw, and sometimes directed between the claws), toothed or bifid (with the apex 'doubled' - split from the middle of the claw or nearer the apex to provide two teeth for each claw, one generally shorter than the other). Synonym: pretarsus.

Tarsomere (leg, tarsus) - the individual portion of a tarsus. There are five tarsomeres in the weevil tarsus, although the fourth is very small and often hidden between the lobes of the third, hence the description 'pseudotetramerous'. The only exception is the Raymondionymidae, where there are four tarsomeres only. See also fifth tarsomere; the fifth tarsomere is sometimes termed the onychium, although as this term is sometimes also used for the claws themselves, it is best avoided. Synonym: tarsal segment. [Emended June 2015]

Tectum (male genitalia) – dorsal plate of penis, when present. See also pedon and pedotectal genitalia.

Tegmen (male genitalia) - Anterior or basal part of the aedeagus, comprising a more or less complex ring or sheath surrounding the aedeagus (q.v.), a ventral apodeme directed anteriorly (see 'apodeme of tegmen') and dorsal parameroid lobes (q.v.) or parameroid plate (q.v.). The shape of the tegmen, and particularly of the parameroid lobes / plate differs between the pedal and pedotectal type of aedeagus. Synonym: phallobase (in part). (emended June 2012)

Tegmenal (in conjunction with ring, apodeme etc) - see Tegminal.

Tegminal apodeme (male genitalia) - apodeme extending anteriorly from base of tegmen ventrally. Wanat (2007) refers to the 'basal piece' as comprising the apodeme and arms curving round the side of the aedeagus and meeting a dorsal plate (tegminal plate), and an equivalent concept is used by Lawrence et al (2010). Synonyms: Apodeme of tegmen, manubrium, tegminal strut, basal piece (in some cases). [Emended September 2015]

Tegminal plate (male genitalia) – see Parameroid plate.

Tegminal ring (male genitalia) – the part of the tegmen that forms a more or less complete ring around the penis, from which arise the parameroid lobes or plate and the tegminal apodeme. Synonym: basal piece (in some cases).

Tegminal strut (male genitalia) – see Tegminal apodeme, Manubrium.

Temones (male genitalia) – see penis apodemes. Term used by Alonso-Zarazaga, 1989, 1990).

Temple (head, head capsule) - lateral part of the head between the posterior margin of the eye and the margin of the pronotum (Wanat, 2001). See discussion under Head.

Tentorium (head) – internal skeleton of the head, comprising plesiomorphically a pair of anterior arms joined to a pair of posterior arms, these united by a tentorial bridge. The anterior and posterior arms are apodemal invaginations arising at the anterior tentorial pits and posterior tentorial pits respectively. Discussed by Lyal (1995).

Tergal formula (abdomen) – see under Pygidium.

Thoracic horns (prothorax) - anteriorly-directed horns arising from the prothorax ventrally, just anterior to the coxal cavity. Often with a median more or less forked sheath between them. Image below is of a thorax viewed from the ventro-posterior aspect, with the sheath pointing towards the viewer and the horns pointing away.

Tormae (larva, mouthparts) - sclerotized rods in the labrum. Their shape can provide taxonomically-useful characters. Synonym: labral rods.

Transfer apparatus (male genitalia, endophallus) – more or less complex sclerotised elements in endophallus around the gonopore. Velázquez de Castro et al. (2007) distinguished three different sclerites in *Sitonini*: pinna, cucullus, and hamulus. cf flagellum. Synonyms: aggonoporium; basal sclerite; complex apparatus; armature sclerites; anterior endophallic sclerite; sclerotized structure of internal sac.

True corbel – see discussion under corbel

Uncus (leg, tibia) - Apical tooth of the tibia associated with the Inner Flange (q.v.). The traditional distinction between the uncus and the mucro (q.v.) is that the mucro arises from the ventral (inner) apical angle and is not continuous with the dorsal (outer) margin), while the uncus arises from, or is continuous with, the dorsal margin (Marvaldi & Lanteri, 2005). Thompson (1992) notes that the uncus is always associated with the inner flange, so that a mucro, once associated with the inner flange, is termed an uncus. There are cases where the flange and the mucro do not smoothly connect, even where both are present.

Unguitractor - see Fifth tarsomere (preferred term)

Urosternite - see spiculum gastrale (preferred term)

Vagina (female genitalia) – tubular part of the ovipositor between the junction with the common oviduct and the gonocoxites, where it opens through the vulva.

Vaginal palp see 'coxite-stylus'

Venter (abdomen) – the visible ventral surface of the abdomen.

Venter (head, head capsule) - area between the eyes ventrally, limited anteriorly by the base of the rostrum (level of the anterior margin of the eyes) and posteriorly by the posterior tentorial pit. Includes part of the submentum sensu Lyal, 1995 and part of the postgena. (Wanat, 2001). There is potential to confuse this use of the term with the more common use as applied to the ventral surface of the abdomen, and it is not in general use.

Ventral sulci (head, rostrum) - (Wanat, 2001)

Ventrite (abdomen) – visible (exposed) abdominal sternites. The first two sternites are concealed, forming part of the coxal cavity, so that the first ventrite is the third sternite. There are five ventrites visible in weevils. The nature of the sutures between the ventrites is discussed by Thompson (1992).

Vertex (head, head capsule) - posterior part of the head, extending from the occiput to the level of the antennal insertions. It, and its subdivisions, are discussed under the entry on Head. Some authors limit the term to the area behind the eyes dorsally, not including the part of the head capsule normally covered by the pronotum (i.e. lacking the sculpture and setation of the more normally exposed part of the head) [Emended June 2015]

Vulva (female genitalia) – opening of vagina between gonocoxites.

Wing-binding patch (abdomen) – patch of very fine spines generally found paired on abdominal sterna, used by the weevil in folding the wings under the elytra.



ASEAN Regional Training Workshop on Diagnostics of Weevils of Quarantine Importance

July 10-22, 2017

**Institute of Weed Science, Entomology and Plant Pathology
College of Agriculture and Food Science
University of the Philippines Los Baños**

SESSION 4

WEEVILS IN AGRICULTURAL CROPS

Lecture Notes by:

DR. CELIA DR. MEDINA

**Institute of Weed Science, Entomology and Plant Pathology
College of Agriculture and Food Science
University of the Philippines Los Baños**

Weevils Attacking Economically Important Crops



Celia DR. Medina, PhD

Institute of Weed Science, Entomology & Plant Pathology
College of Agriculture and Food Science
University of the Philippines Los Baños

Weevils Attacking Mango

(*Sternochetus frigidus*, *S. mangiferae*, & *S. olivieri*)

Sternochetus mangiferae (Fabricius, 1775)

Other Scientific Names

Acryptorhynchus mangiferae (Fabricius)

Cryptorhynchus mangiferae (Fabricius)

Curculio mangiferae

Sternochetus ineffectus (Walker)

International Common Names

English: mango seed weevil, mango stone weevil; mango weevil; Spanish: gorgojo de la semilla del mango; picudo de la semilla del mango, French: charançon de la graine du manguier; charançon de la mangue

EPPO code

CRYPMA (*Sternochetus mangiferae*)

Distribution

Asia - Bangladesh, British Indian Ocean Territory, India, Indonesia, Myanmar, Nepal, Oman, Sri Lanka, Yemen (CABI as of August 2016)

(Invalid previous record: Bhutan, China, Hongkong, Malaysia, Pakistan, Philippines, Thailand, UAE, Vietnam)

Africa - Central African Republic, Gabon, Ghana, Guinea, Kenya, Liberia, Madagascar, Malawi, Mauritius, Mozambique, Reunion, Seychelles, Tanzania, Zambia (Invalid previous record: Nigeria)

Not America - Hawaii

Central America & Caribbean - Barbados, British Virgin Islands, Dominica, Grenada, Guadeloupe, Martinique, Montserrat, St. Lucia, St Vincent & Grenadines, Trinidad & Tobago, United States Virgin Islands **South**

America - Brazil, Chile, French Guiana

Oceania - Australia (Northern Territory, NSW, Queensland), French Polynesia, Guam, New Caledonia, Northern Mariana Islands, Tonga, Wallis and Futuna Islands

Host Range

Complete development is only achieved on mangoes. Oviposition was obtained in the laboratory on potatoes, peaches, *Litchi chinensis*, plums, *Phaseolus vulgaris* and several cultivars of apples, but none of the resulting larvae reached maturity.

Biology & Ecology:

Adult habits and behavior

- feed on the leaves and tender shoots of mangoes during March and April in India
- nocturnal, fly readily and usually feed, mate and oviposit at dusk
- enter a diapause after emergence, which varies in duration with the geographic range (ie. southern India, all adults emerging during June enter a diapause from July until late February of the following year)
- onset and termination of diapause appear to be associated with photoperiod
- diapause under loose bark on mango tree trunks and in branch terminals, or in crevices near mango trees
- few adults live through two seasons with a diapause period in between
- females began oviposition 3-4 days after mating, when the fruit was about marble-size
- oviposition period varies from 3 - 5 weeks
- lay eggs mostly on the sinus of the fruit or sometimes on the stems
- female makes a boat-shaped cavity in the skin (epicarp) into which an egg is deposited
- One female may lay 15 eggs per day, with a maximum of almost 300 over a 3-month period in the laboratory (Balock and Kozuma, 1964)
- usually remain in the vicinity of the parent tree until the following fruiting season

Life cycle:

- egg incubation requires 5-7 days
- newly emerged larva burrows through the flesh of the fruit and into the seed
- tunnel and seed entry are completely obliterated as fruit develops, so that in time it is impossible to distinguish infested from non-infested seeds, unless they are cut open
- larval development usually occurs within the maturing seed, but also very occasionally within the flesh
- larval development takes about a month in S India, 22 days to 10 weeks in Hawaii
- five or seven larval instars
- pupates within the seed and rarely in the flesh
- pupal period lasts about a week
- one to six adults may develop per seed
- adults cut their way out of the naked seed, usually via a small circular hole made in the concave edge of the endocarp, generally 4-8 weeks after the fruit falls and decays
- move out of the seeds and seek hiding places by crawling, rather than flying.
- estimated time required for development from egg to adult is 35-54 days

Natural Enemy:

“There is no published information about the parasites and predators of *S. mangiferae*. The only published record of any pathogen is a baculovirus affecting the larvae of *S. mangiferae* (Shukla et al., 1984).”

***Sternochetus frigidus* (Fabricius, 1787)**

Other Scientific Names

Acryptorhynchus frigidus
Sternochetus gravis (Fabricius)
Cryptorrhynchus gravis Fabricius
Curculio frigidus Fabricius
Cryptorhynchus frigidus (Fabricius)
Acryptorrhynchus frigidus (Fabricius)

International Common Names

English: mango fruit weevil, mango weevil, mango pulp weevil, mango flesh weevil, northern mango weevil
 Spanish: picudo del mango
 French: charançon du manguier

EPPO code

CRYPGR (*Sternochetus frigidus*)

Distribution

Asia - Bangladesh, Brunei Darussalam, India, Indonesia, Malaysia, Philippines, Thailand, Myanmar (CABI as of August 2016) **Oceania** - Papua New Guinea

Host Range

Cultivated and wild *Mangifera* spp. (*Mangifera foetida* (bachang), *Mangifera odorata* (kurwini mango), *Mangifera sylvatica*), including all varieties of *M. indica* as well as *M. foetida*. However, oviposition on *M. foetida* is rare.

Biology & Ecology

Adult habits and behavior:

- In Tripura, India, reproductively immature adult *S. frigidus* overwinter inside seeds or other protective places from May until February. Mating occurs 10-15 days after termination of hibernation. Oviposition occurs from March to May on immature mango fruits.
- requires 6 weeks to be fully matured and able to mate
- hide during the day, for example in bark crevices or under epiphytic plants
- feed at night on the gum that exudes from puncture wounds on young fruits
- strong fliers but do not move far from their emergence sites
- mate repeatedly
- A female will often lay her total daily quota of eggs in one fruit.

- If suitable fruits are available, females lay about eight eggs daily, producing a total of 75 (maximum 180 eggs) in 3 weeks.
- If deprived of suitable fruits for 5 months, the egg production drops to three eggs per day; additional delays lead to further reduction in oviposition.
- Female dies soon after oviposition
- Large populations of *S. frigidus* are generally only found in sites with high humidity, containing large stands of mangga (*M. indica*), kekembem and kweni (*M. foetida*) trees which flower more than once a year.
- The weevils have only one generation during the fruit season Weevils that survive from the preceding year infest fruit formed after the earliest flowering.
- If early flowering does not occur, they attack the fruits formed after the main flowering period. An extended flowering period leads to the production of large numbers of weevils. In a dry year, or if only one flowering period occurs, the weevil population remains low (on average four eggs per fruit).

Life cycle:


- The eggs are laid on fruits that has a minimum diameter of 6 cm.
- Similar to *S. mangiferae*, the females remove a piece of peel, oviposit, and cover the eggs with a brown layer of gum and faeces creating a crescent-shaped incision around the oviposition site.
- Eggs hatch after 4-65 days depending on the temperature (De and Pande, 1988).
- Newly hatched larva tunnels directly through the fruit pulp to the kernel. The shell of the kernel is not attacked but while boring through the pulp, the larvae imbibe the exuding sap, so that the tunnel remains relatively dry. The granular faeces accumulate in the hole formed by feeding.
- Larva undergoes 5 larval stadium over approximately 20 days
- Pupation takes place in a brown cocoon, constructed of frass, within these chambers.
- In the Philippines, total development from egg to adult is 32 days but the adult remains inside the fruit for another 37 days. In Bogor, Indonesia, it is observed to cover 5-7 weeks.
- Survival of newly hatched larvae reveal that exuding gum is responsible for 30-50% mortality. Dry conditions are thought to affect young adults adversely.

Natural enemy:


- *Flavopimpla mangae* - ichneumonid parasitic wasp
- Formicidae (*Oecophylla smaragdina*, *Camponotus* sp. and *Monomorium* sp.)
- *Aspergillus* sp. - entomogenous fungus
- *Rhizoglyphus* sp. - ectoparasitic mite (of hibernating adult)

Diagnostic Protocol of Mango Weevils


based Key to the species of mango weevils (Oberprieler & Banks, 2008 unpublished)
 Source: European and Mediterranean Plant Protection Organization PM 7/106 (1)



Sternochetus mangiferae, mesosternum & mesothoracic leg (Photo courtesy of LSV)



Aedeagus, dorsal view. (A) *S. olivieri* (B) *S. mangiferae* (C) *S. frigidus* (Photos of CSIRO)



Sternochetus mangiferae habitus, dorsal view




Sternochetus mangiferae habitus, lateral view
Length: 7.5-10mm

Sternochetus spp. habitus (i) dorsal view; (ii) lateral view (Photo courtesy of LSV)

***Sternochetus mangiferae*:**

- Body length: 7.5-10mm
- Elytra with sides nearly parallel from base to beyond middle
- Interstriae flat to faintly but evenly costate (ridged)
- strial punctures rectangular to square
- whitish macula forming a more or less distinct V and transverse posterior band
- pronotum with erect black scales scattered over basal part of pronotal disk -aedeagus with pair of internal sclerites separate, not touching apically.



Sternochetus frigidus habitus, dorsal view



Sternochetus frigidus habitus, lateral view
Length: 3.8-5.9mm

***Sternochetus frigidus*:**

- Body length: 3.8-5.9mm
- Elytra narrowing from base to apex
- Odd interstriae except sutural one distinctly costate-tuberculate
- Strial punctures round
- Whitish macula fragmented but usually
 - forming a vague anterior inverted triangle inscribing a similar, smaller black median triangle and a broken posterior band on declivity
- Pronotum with erect black scales arranged in medial pair of loose clusters
- Aedeagus with pair of internal sclerites overlapping apically.

***Sternochetus olivieri*:**

- Body length: length 6–8 mm.
- Elytra with a large, whitish macula (patch) stretching from just behind humeri (shoulders) to top of declivity, inscribing a black, inverted medial triangle before
- middle length and sometimes posteriorly interrupted by a fainter, dark, transverse band above declivity;
- Pronotum medially with a conspicuous carina (keel) in basal 2 / 3 of length, which is flanked on either side by a line of white scale and at its anterior end (in middle pronotum) by a tuft of dense, erect black scales;
- Aedeagus with sides nearly parallel, apically broadly rounded and no internal sclerites.



Control of Mango Weevils

Cultural control

Good orchard sanitation involving the destruction of all the fallen fruit, stones and fruits with seed weevil damage during and immediately after mango harvest.

In nursery, the seed may be shelled and only sound kernels planted.

Open center- canopy pruning makes the micro climate unfavourable to *S. frigidus*.

Fruits of the size of a chicken egg can be bagged individually using durable paper bags to reduce damage of *S. frigidus*.

Biological control

The ant *Oecophylla smaragdina* is an effective biocontrol agent of *S. mangiferae* adults (Peng and Christian, 2004, 2007).

The only published record regarding pathogens is of a baculovirus affecting the larvae of *S. mangiferae* (Shukla et al., 1984).

Host plant resistance

In India, ten cultivars (out of 92 studied) were found to be free from *S. mangiferae* infestation, and these cultivars are Sindhu, Bombay Green, Firangi Ludua, Pulihora, Jahangir, Sabja, Salgadino, Hatizool, Dodamio and Fazri (Godse and Bhole, 2003). Larval penetration of the seed of the variety Itamaraca is reported to be impossible (Balock and Kozuma, 1964).

Chemical control

Chemical control has been used with some success and a wide range of insecticides have been recommended (see, for example, Shukla and Tandon, 1985; Villiers, 1987). The main strategy is to attack diapausing adults by trunk applications or to use foliar sprays at the time of oviposition.

Means of Movement & Dispersal

Longer-range dispersal occurs largely through transport of fruit and seeds containing the larvae, pupae or adults. The pest has been intercepted in mango fruits and seeds in international trade.

Phytosanitary Measures

Methyl bromide fumigation at the rate of 36 gm³ for 8 h at 21°C killed all stages but injured the fruit (Balock & Kozuma, 1964).

Balock & Kozuma (1964) found gamma radiation to be the most effective method for killing or sterilizing weevils within fruit.

In the Philippines, Obra et al (2014) At 150 Gy (measured doses 96.7–164.1 Gy) caused complete sterility of *S. frigidus*.

In Hawaii, Seo et al. (1974) treated packaged mangoes with cobalt-60 gamma irradiation. Minimum doses of 20.6 and 32.9 krad killed insects of all stages within the fruits; the few surviving adults were sterile and short-lived.

In South Africa, irradiation of ripe, marketable fruit protected it from damage and prevented adult emergence. The most effective dosages ranged from 0.5 to 0.85 kGy; higher rates tended to be phytotoxic (Kok, 1979).

Hot and cold treatment of fruit has also been tried but gave unreliable results and proved phytotoxic (Balock & Kozuma, 1964; Seo et al., 1970; Shukla & Tandon, 1985).

References:

CABI. (2007). Crop Protection Compendium, 2005 Edition. (c) CAB International Publishing. Wallingford, UK. www.cabi.org

Alam MZ, 1962. A list of insects and mites of East Pakistan. Dhaka, Bangladesh: Department of Agriculture.

Bagle BG, Prasad VG, 1985. Studies on varietal incidence and control of stone weevil, *Sternochetus* (=Cryptorrhynchus) *mangiferae* Fabricius (Coleoptera: Curculionidae). *Indian Journal of Entomology*, 47(3):362-364.

Balock JW, Kozuma TT, 1964. Notes on the biology and economic importance of the mango weevil, *Sternochetus mangiferae* (Fabricius), in Hawaii (Coleoptera: Curculionidae). *Proceedings of the Hawaiian Entomological Society*, 18:353-364.

Basio RG, Johnson PJ, Pua DR, Bergonia HT, Diloy CC, Villegas EL, 1994. Mango pulp weevil (*Sternochetus frigidus* (Fabr.)) (Curculionidae, Coleoptera) found in Palawan. *Philippine Entomologist*, 9(3):350-351.

De Jesus, L.R.A., and F. Cortez. 1998. Field infestation and chemical control of the mango pulp weevil, *Sternochetus frigidus* (Fabr.) in Palawan. *Philipp. Agric.* 81: 131–140.

De Jesus, L.R.A., and R. R. Gabo. 2000. Life history and host range of the mango pulp weevil, *Sternochetus frigidus* (Fabr.) in Palawan, Philippines. *Philipp. Agric. Sci.* 83: 145–150.

De Jesus, L.R.A., S.M.F. Calumpang, J. R. Medina, and K. Ohsawa. 2003. Feeding and reproductive behavior of the mango pulp weevil, *Sternochetus frigidus* (Fabr.), at different phenological stages of mango, *Mangifera indica* L. *Philipp. Agric. Sci.* 86: 282–289.

Dey K, Pande YD, 1987. Evaluation of certain non-insecticidal methods of reducing infestation of the mango nut weevil, *Sternochetus gravis* (F.) in India. *Tropical Pest Management*, 33(1): 27-28, 101, 105.

Dey K, Pande YD, 1987. Efficacy of insecticides in controlling *Sternochetus gravis* (Fabricius). *Indian Journal of Agricultural Sciences*, 57(9):678-680.

Dey K, Pande YD, 1988. Bionomics and some behavioural aspects of the mango stone weevil, *Sternochetus gravis* (Fabricius) (Coleoptera: Curculionidae). *Entomon*, 13(1):17-24.

Devi MB, Prasad B, 1992. A note on the hibernation of mango fruit weevils (Curculionidae: Coleoptera). *Bioved*, 3(2):235-236.

EPPO, 2006. PQR database (version 4.5). Paris, France: European and Mediterranean Plant Protection Organization. www.eppo.org. Firman ID, 1982. Plant protection news. Information Circular, South Pacific Commission, No. 90:8 pp.; [7 fig.].

Follett PA, Gabbard Z, 2000. Effect of mango weevil (Coleoptera: Curculionidae) damage on mango seed viability in Hawaii. *Journal of Economic Entomology*, 93(4):1237-1240.

Kalshoven LGE, 1951. The Pests of Cultivated Plants in Indonesia. Part II. The Hague, Netherlands: Van Hoeve. Kalshoven LGE, Laan P. A. van der (Reviser and translator), 1981. Pests of crops in Indonesia. Jakarta, Indonesia: Ichtisar Baru.

Kok IB, 1979. Control of the mango seed weevil by trapping and irradiation. *Citrus and Subtropical Fruit Journal*, No. 552:14-16; [1 fig.].

Medina CDR, Velasco LRI. 2001. Diagnostic characters to identify the sexes of the adult mango pulp weevil (*Sternochetus frigidus* (Fabr.) (Coleoptera: Curculionidae). *Philipp. Ent.* 15(2): 191-195.

Medina, J. R., O. S. Opina, L.R.A. De Jesus, and S.M.F. Calumpang. 2005. Development of an integrated pest management program for mango in Palawan, Philippines. *Philipp. Agric. Sci.* 88: 453–461.

Obra, G. B., S. S. Resilva, and L.R.A. Lorenzana. 2013. Irradiation as a potential phytosanitary treatment for the mango pulp weevil *Sternochetus frigidus* (Fabr.) (Coleoptera: Curculionidae) in Philippine super mango. *Philipp. Agric. Sci.* 96: 172–178.

Obra, GB, Resilva SS, Follett, PA, Lorenzana, LRJ. 2014. Large-Scale Confirmatory Tests of a Phytosanitary Irradiation Treatment Against *Sternochetus frigidus* (Coleoptera: Curculionidae) in Philippine Mango. *J. Econ. Entomol* 107(1)161-165.

Pande YD, Dey K, 1986. Outbreaks and new records. India. *Sternochetus gravis* (Col.: Curculionidae) causing serious damage to mangoes in northeast India. *FAO Plant Protection Bulletin*, 34(2):109.

Pinese, B., and Holmes, R. (2005). Managing mango seed weevil. Horticulture and Forestry Science Department of Primary Industries and Fisheries, The State of Queensland.

Rahman ML, Ahmad M, 1972. The immature stages of the mango fruit weevil, *Sternochetus frigidus* Fabricius (Coleoptera: Curculionidae). *Pakistan Journal of Scientific and Industrial Research*, 15:188-190.

Rahman MF, Khan S, 1989. Anatomy and histology of the alimentary canal of the adult mango weevil, *Sternochetus frigidus* (Fabricius) (Coleoptera: Curculionidae). *Bangladesh Journal of Zoology*, 17(2):189-195.

Seo, S. T., R. M. Kobayashi, D. L. Chambers, L. F. Steiner, C.Y.L. Lee, and M. Komura. 1974. Mango seed weevil: Cobalt-60 gamma irradiation of packaged mangoes. *J. Econ. Entomol.* 67: 504–505

Shukla RP, Tandon PL, Singh SJ, 1984. Baculovirus-a new pathogen of mango nut weevil, *Sternochetus mangiferae* (Fabricius) (Coleoptera: Curculionidae). *Entomon*, 10:215-218

Shukla RP, Tandon PL, 1985. Bio-ecology and management of the mango weevil, *Sternochetus mangiferae* (Fabricius) (Coleoptera: Curculionidae). *Current Science*, 53:593-594. SPC, 1989.

Shukla R.P. and Tandon P.L. (1985). Bio-ecology and control of the mango weevil *Sternochetus mangiferae* (Fabricius) (Coleoptera: Curculionidae) *International J. Trop. Agri.* Vol 3 No 4 pp 293 -303.

Silva, AC and Ricalde, MP. 2017. First occurrence of *Sternochetus mangiferae* (Fabricius) (Coleoptera: Curculionidae) in Brazil. *Neotrop. Entomol* May 24, 2017 doi: 10.1007/s13744-017-0523-1. [Epub ahead of print]

Van Dine DL, 1907. Notes on *Cryptorhynchus mangiferae* (Coleoptera). *Proceedings of the Hawaiian Entomological Society*, 1:79-82.

Villiers EA de, 1987. Mango weevil must be controlled. *Information Bulletin, Citrus and Subtropical Fruit Research Institute, South Africa*, 176:12-13.

Voute AD, 1935. *Cryptorhynchus gravis* F. and the causes of its multiplication in Java. *Archives Néerlandaises de Zoologie*, 2:112-142.

Waterhouse DF, 1993. *The major arthropod pests and weeds of agriculture in Southeast Asia*. Canberra, Australia: ACIAR.

Wheatley PE, 1961. Insect pests in the Coast Province of Kenya. 1-Mango. *East African Agriculture For. Journal*, 26:129-131.

Woodruff RE, 1970. The mango seed weevil, *Sternochetus mangiferae* (Fab.) (Coleoptera: Curculionidae). *Entomology Circular, Florida Department of Agriculture and Consumer Services, Division of Plant Industry*, 93:1-2.

Wongsiri S, 1983. *Insect pests of agricultural crops in Thailand*. Bangkok, Thailand: Odeon Store Publishing.

Weevils Attacking Banana

Cosmopolites sordidus (Germar), *Philicoptus iliganus* (Heller)

Cosmopolites sordidus (Germar, 1824)



Genus: Cosmopolites; Species: Banana Weevil Borer

Other Scientific Names

Sphenophorus sordidus Germar
Calandra sordida Germar
Sphenophorus striatus Fabricius
Sphenophorus musaecola
Sphenophorus cribricollis Walker
Sphenophorus pygidialis Chevrolat
Sphenophorus liratus
Metamasius sordidus (Germar)
Cosmopolites striatus (Fabricius)
Metamasius striatus (Fabricius)

International Common Names

English: banana weevil, banana weevil borer, banana root weevil, banana root borer, banana rhizome weevil, banana borer, plaintain weevil, corm weevil, banana beetle, cosmopolitan root borer, plantain root borer

Spanish: picudo negro, picudo negro del banano, gorgojo del tallo del banano, gorgojo negro del pl-tano, barrenillo del banano

French: charançon du bananier, charançon noir des bananiers, broca da bananeira, moleque da bananeira

EPPO code

COSMSO (*Cosmopolites sordidus*)

Distribution

Asia - Bangladesh, Brunei Darussalam, Cambodia, China, Hongkong, Taiwan, Christmas Is, Inida, Indonesia, Israel, Japan, S. Korea, Malaysia, Peninsular Malaysia, Maldives, Myanmar, Mepal, Pakistan, Philippines, Singapore, Sri Lanka, Thailand, Vietnam

Africa - Angola, Benin, Burkina Faso, Burundi, Cameroon, Cape Verde, Comoros, Congo, Gabon, Ghana, Guinea, Kenya, Madagascar, Malawi, Mauritania, Mauritius, Niger, Rwanda, Reunion, St Helena, Senegal, Sao Tome & Principe, Seychelles, Sierre Leone, Somalia, S. Africa, Tanzania, Togo, Uganda

North America - Florida (eradicated), Hawaii, Mexico

Central America & Caribbean - Bermuda, Costa Rica, Cuba, Dominica, El Salvador, Grenada, Guadeloupe, Guatemala, Haiti, Honduras, Jamaica, Martinique, Nicaragua, Panama, Puerto Rico , Saint Lucia , Saint Vincent and the Grenadines ,Trinidad &Tobago

Europe - Madeira, Canary Island

South America - Argentina, Bolivia, Brazil, Chile, Columbia, Ecuador, Easter Is, French Guiana, Guyana, Peru, Suriname, Venezuela

Oceania - American Samoa, Australia (NSW, Queensland, Western Australia), Belau, Cook Is, Fiji, French Polynesia, Guam, New Caledonia, Niue, Paps a New Guinea, Samoa, Solomon Is, Tonga, Vanatu, Wallis & Futuna Islands

Biology & Ecology

Life cycle:

- Wide variability in stage duration: 4-36 days for eggs, 12-165 days for larvae, 1-4 days for prepupae, 4-30 days for pupae and 24-220 days from egg to adult
- Under tropical conditions, the egg stage is normally 6-8 days, while the larval period is 20-25 days.
- After hatching the larvae tunnel into the host-plant tissues
- Larvae are variously reported have 5-8 instars
- Pupation is usually near the surface of the corm, inside an oval chamber
- Adults emerge from pupa in 5-8 days
- Adults are fully hardened in 5-7 days

Reproduction:

- Pre reproductive period is 5-20 days for females and for males at 18-31 days.
- Age at first oviposition is 27-41 days
- Egg production was influenced by temperature, but not rainfall or humidity.
- Eggs are laid throughout the year at a rate varying with temperature and up to 100/ female/year
- Under field conditions, egg density is greatest on flowering plants

- Eggs are placed singly in small cavities, chewed out by the female, with over 90% occurring in the base of the pseudostem, with the remaining eggs found in the corm and roots near the soil surface.
- In stands displaying high mat (a condition in which part of the corm appears above the soil surface), more eggs are found on the corm than on the pseudostem. The majority of eggs are normally placed below the soil surface.
- Crop residues also receive high levels of oviposition with as many as 200 eggs recorded from a single stump.

Adult survivorship:

- can survive extended periods without food
- sensitive to desiccation and will die within 48 hours if kept in a dry substrate
- Preference for moist environments continues to breed throughout the year, provided that food supply is plentiful
- live 1 year or less but some adults may live as long as 2 years

Adult habits:

- gregarious and are found in clusters in cavities and depressions in the outer sheaths of the banana close to the ground surface and also below the surface
- found either in the base of leaf sheaths or in the soil around the mat
- some adults are found associated with cut crop residues
- sedentary for extended periods of time
- nocturnal active between 18:00 and 06:00 with greatest activity from 21:00-04:00
- rarely fly, but commonly walk over the soil surface and vegetation

Adult trivial movement:

- not strong flyers and only cover short distances
- unusually sedentary: of 400 marked weevils released at one point in a Ugandan plantation, 35% were recovered over an 8-month period within a radius of 10 yards of the original release point

Population build up

- In a field trial, weevils were released 11 months after planting. Four months later, weevil numbers remained stable in mulched plots, while declining by 38-53% in controls and intercrops
- Overall, the field population of weevils remained below release levels until 39 months after planting and peaked at 47 months at 40% higher than release levels.

Natural Enemies

- Beetle predators of egg and larva.: *Plaesius javanus* , *P. laevigatus*; *Hololepta quadridentata*, *Belonuchus ferrugatus*, *Priochirus unicolor*, *Thyreocephalus interocularis*, *Dactylosternum hydrophiloides*, *D. abdominale* and *Cathartus* sp.
- In Cuba, the ant *Tetramorium bicarinatum* is reported to keep *C. sordidus* but not *T. guineense* .
- No hymenopterous or dipterous parasitoids.
- Strains of *Beauveria bassiana* in laboratory tests.

Control of Banana Corm Weevil

Simon (1994) produced a bibliographical synthesis reviewing integrated control of *C. sordidus*.

Cultural Control

- Use clean planting materials in the planting of new stands
- 'Paring and Sterilization' - peel the rhizomes free of lesions and immerse in hot water at 54°C for 10 minutes
- Dig out and remove old corms, trash and other materials in which weevils may breed
- Desucker and remove water suckers regularly and keep the plantation free of weeds at all times.
- 'Count, Cut and Dry' - Cut down the spent pseudostem close to the ground as soon as bunch is harvested. Cut it into lengths of up to 60 cm and split each along its length.
- Good husbandry practices, such as clean weeding, manuring and mulching produce vigorous banana plants which have improved weevil tolerance

Biological Control

Approaches include the use of endemics, exotics (classical biological control), secondary host association and microbial control.

- Fiji - *Plaesius javanus* and *P. laevigatus* in 1918
- Queensland - *P. javanus* from 1921 to 1928; *Dactylosternum hydrophiloides* in 1939
- Cook Island - *P. javanus* and *P. laevigatus* from 1937 to 1940
- Chiapas, Mexico - *P. javanus* in 1933
- Jamaica - *Dactylosternum hydrophiloides* and *D. abdominale* in 1942
- St. Vincent - *Hololepta quadridentata* in 1976
- New South Wales - Nematodes *Steinernema carpocapsae*, *S. glaseri* and *S. bibionis* in 1991
- Cuba - the fungus *Beauveria bassiana* & ants *Tetramorium guineense* and *Pheidole megacephala*

Host-Plant Resistance

Early researches showed potential for development of resistant varieties. The most important resistance mechanism appears to be antibiosis.

Chemical Control

- Apply insecticides to the bases of the plants, including the suckers, and the surrounding soil surface for a distance of 30 cm around the plants, using knapsack sprayers, high volume spraying units or by sprinkling as dusts
- In Tonga, planting material is dipped in suitable insecticides or granules 2-3 times a year
- In Puerto Rico two nematicides controlled the borer in plantains

Trapping and Pheromonal Control

- Disk-on-stump traps- corm slices or leaves placed on top of harvested plants cut at the rhizome
- Pheromone traps “sordidin”; 4 traps/ha, changed and moved every month

References

- Batra HN, 1952. Occurrence of three banana pests at Delhi. *Indian Journal of Entomology*, 14:60.
- Beauhaire J, Ducrot PH, Malosse C, Rochat D, Ndiege IO, Otieno DO, 1995. Identification and synthesis of sordidin, a male pheromone emitted by *Cosmopolites sordidus*. *Tetrahedron Letters*, 36(7):1043-1046.
- Bennett FD, Rosen D, Cochereau P, Wood BJ, 1976. Biological control of pests of tropical fruits and nuts. In: Huffaker CB, Messenger PS, eds, *Theory and Practice of Biological Control*. New York, USA: Academic Press, 359-395.
- Braithwaite BM, 1958. Ground spray treatments for control of banana beetle borer (*Cosmopolites sordidus* Germ.). *Journal of the Australian Institute of Agricultural Science*, 24:27-34.
- Budenberg WJ, Ndiege IO, Karago FW, 1993. Evidence for volatile male-produced pheromone in banana weevil *Cosmopolites sordidus*. *Journal of Chemical Ecology*, 19(9):1905-1916.
- Cheng KH, 1957. Banana corm weevil and its control experiments. *Agric. Res.*, 7:45-54.
- Clausen CP, 1978. Introduced parasites and predators of arthropod pests and weeds: a world review. *Agriculture Handbook*, United States Department of Agriculture.
- Collin PJ, Treverrow NL, Lambkin TM, 1991. Organophosphorus insecticide resistance and its management in the banana weevil borer, *Cosmopolites sordidus* (Germar) (Coleoptera:Curculionidae) in Australia. *Crop Protection*, 10(3):215-221.
- Deang RT, Caburubias RB, Quero EM, 1969. Insecticide test for the control of abaca corm weevil. *Philippine Journal of Plant Industry*, 34(1/2):79-87.
- Delattre P, 1980. Investigations into a method for estimating populations of the banana weevil, *Cosmopolites sordidus* Germar (Col., Curculionidae). *Acta Oecologica, Oecologia Applicata*, 1(1):83-92; [6 fig.].
- Fogain R, Price NS, 1994. Varietal screening of some *Musa* cultivars for susceptibility to the banana borer weevil. *Fruits (Paris)*, 49(4):247-251.
- Franzmann BA, 1972. Banana weevil borer in north Queensland. *Queensland Agricultural Journal*, 98:319-321.
- Gettman AD, Mitchell WC, Li P, Mau RFL, 1992. A hot water treatment for control of the banana root borer, *Cosmopolites sordidus* (Germar) (Coleoptera:Curculionidae) in banana planting stock. *Proceedings of the Hawaiian Entomological Society*, 31:59-63.

- Gold CS, Speijer PR, Karamura EB, Tushemereirwe WK, Kashaija IN, 1994. Survey methodologies for pest and disease assessment in Uganda. *African Crop Science Journal*.
- Gold CS, Night G, Abera A, Speijer PR, 1998. Hot-water treatment for the control of the banana weevil, *Cosmopolites sordidus* Germar (Coleoptera: Curculionidae), in Uganda. *African Entomology*, 6(2):215-221.
- Gold CS, Night G, Speijer PR, Abera AMK, Rukazambuga NDTM, 1998. Infestation levels of banana weevil, *Cosmopolites sordidus* Germar (Coleoptera: Curculionidae), in banana plants established from treated propagules in Uganda. *African Entomology*, 6(2):253-263.
- Greathead DJ, 1986. Opportunities for biological control of insect pests in tropical Africa. *Revue de Zoologie Africaine*, 100(1):85-96.
- Greathead DJ, 1993. Report on miscellaneous observations made during a visit to Cuba for FAO to assist in setting up a programme for biological control of water hyacinth. 20 May - 4 July 1993. (Unpublished IIBC report) Wallingford, UK: CAB International.
- Kalshoven LGE, 1951. *The Pests of Cultivated Plants in Indonesia. Part II.* The Hague, Netherlands: Van Hoeve. Kalshoven LGE, Laan PAVan der (Reviser and translator), 1981. *Pests of crops in Indonesia.* Jakarta, Indonesia: Ichtar Baru.
- Koppenh fer AM, 1993. Observations on egg-laying behaviour of the banana weevil, *Cosmopolites sordidus* (Germar). *Entomologia Experimentalis et Applicata*, 68(2):187-192.
- Koppenh fer AM, 1993. Egg predators of the banana weevil, *Cosmopolites sordidus* (Germar) (Col., Curculionidae) in western Kenya. *Journal of Applied Entomology*, 116(4):352-357.
- Koppenh fer AM, Reddy KVS, Madel G, Lubega MC, 1992. Predators of the banana weevil, *Cosmopolites sordidus* (Germar) (Col., Curculionidae) in western Kenya. *Journal of Applied Entomology*, 114(5):530-533.
- Koppenhofer AM, Seshu Reddy KV, 1994. A comparison of rearing methods for the banana weevil, *Cosmopolites sordidus* (Germar) (Coleoptera:Curculionidae) on its natural host. *Insect Sci. Appl.*, 15:191-195.
- Mesquita ALM, Caldas RC, 1986. Effect of the age and of the cultivar of banana on the biology and preference of *Cosmopolites sordidus* (Germar, 1824) (Coleoptera, Curculionidae). *Fruits*, 41(4):245-249.
- Ogenga-Latigo MW, Bakyalire R, 1993. Use of pseudostem traps and coefficient of infestation (PCI) for assessing banana infestation and damage by *Cosmopolites sordidus* Germar. *African Crop Science Journal*, 1(1):31-37.
- Ortiz R, Vuylsteke D, Dumpe B, Ferris RSB, 1995. Banana weevil resistance and corm hardness in *Musa* germplasm. *Euphytica*, 86(2):95-102.

Ostmark HE, 1974. Economic insect pests of bananas. Annual review of entomology. Volume 19., 161-176. View Abstract

Peasley D, Treverrow N, 1984. Count, cut, dry. A new approach to banana weevil borer control in bananas. The Banana Bulletin, 12:13.

Roth L, Willis E, 1963. The humidity behavior of *Cosmopolites sordidus* Germar (Coleoptera:Curculionidae). Annals of the Entomological Society of America, 56:41-42.

Sein F Jr, 1934. Paring and heat sterilization of the corms to eliminate the banana root weevil *Cosmopolites sordidus* Germar. Journal of Agriculture. University of Puerto Rico, 18:411-416.

Simon S, 1994. Integrated control of the black banana weevil *Cosmopolites sordidus*. Fruits (Paris), 49(2):151-162.

Speijer PR, Budenberg WJ, Sikora RA, 1993. Relationships between nematodes, weevils, banana plantain cultivars and damage. Annals of Applied Biology, 123(3):517-525. View Abstract

Traore L, Gold CS, Pilon JG, Boivin G, 1993. Effects of temperature on embryonic development of banana weevil, *Cosmopolites sordidus* Germar. African Crop Science Journal, 1:111-116.

Treverrow N, Bedding RA, Dettmann EB, Maddox C, 1991. Evaluation of entomopathogenic nematodes for control of *Cosmopolites sordidus* Germar (Coleoptera: Curculionidae), a pest of bananas in Australia. Annals of Applied Biology, 119(1):139-145.

Treverrow N, 1983. Banana weevil borer. Agfacts, No. H6.AE. 1:2 pp.; [3 fig.].

Viswanath B, 1978. Studies on the biology, varietal response and control of banana rhizome weevil, *Cosmopolites sordidus* (Germar) (Coleoptera:Curculionidae). Mysore Journal of Agricultural Sciences, 11:597-598.

Waterhouse DF, 1993. The major arthropod pests and weeds of agriculture in Southeast Asia. Canberra, Australia: ACIAR.

Waterhouse DF, Norris KR, 1987. Biological control: Pacific prospects., viii + 454pp.; [many fig., 26x18 cm].

Woodruff RE, 1969. The banana root borer (*Cosmopolites sordidus* Germar) in Florida (Coleoptera: Curculionidae). Florida Department of Agriculture, Entomology Circular No. 88, 1-2.

Zimmerman EC, 1968a. Rhynchophorinae of southeastern Polynesia (Coleoptera: Curculionidae). Pacific Insects, 10:47-77.

Zimmerman EC, 1968b. The *Cosmopolites* banana weevils (Coleoptera: Curculionidae: Rhynchophorinae). Pacific Insects, 10:295-299.

Philicoptus iliganus (Heller, 1912)



Other Scientific Names

Coptorhynchus iliganus (Heller)
Philicoptus waltoni Boheman

International Common Names

English: Philippine banana peel scarring weevil

EPPO code

not listed

Distribution

ASIA - Philippines (Davao, General Santos and South Cotabato)

Host Plants

Main host plant: banana (*Musa* spp)

Other minor host plants: *Artocarpus integrifolia* (jackfruit); *Citrus* sp.; *Coffea* spp. (coffee); *Dureo zibethinus* (durian); *Garcinia mangostana* (mangosteen); *Gliricidia sepium* (madre de cacao); *Lansium domesticum* (lansones); *Nephelium lappaceus* (rambutan); *Persea americana* (avocado); *Phaseolus aureus* (mungbeans); and *Theobroma cacao* (cacao) *Crassocephalum crepidioides* (Benth)

Biology & Ecology

Life cycle

- egg to adult development ranges from 111-176 days
- eggs singly or in a mass in the soil
- hatch in 10 days
- larva feeds on the corm
- total larval period from 102–174 days on banana suckers
- pupate in a chamber in the soil and this pupal period lasts 10–23 days

Adult behaviour & habits

- relatively immobile, unable to move long distances and cannot fly no hind wings and the elytra (first pair of wings) are firmly united at the suture
- hides in leaf axils, between touching leaves and concealed among fruit
- has periods of inactivity interspersed with periods of active crawling
- feeds on the surface of young fingers, youngest leaf of banana plants which have not yet fruited, lower bracts before young banana fingers are exposed
- tend to remain within the same area for several years without spreading

References:

Dawi, NM. 1982. Biology, Ecology and Control of Banana Scarring Weevil, *Philicoptus iliganus* (Heller) (Coleoptera: Curculionidae). Unpublished PhD Thesis. Department of Entomology, UP Los Banos. 123pp.

Heller, KM. 1912. Philippine Russelkafer. *Phil. J. Sci* 7(5):295-415.

Ostmark, HE. 1974. Economic insects of banana. *Ann. Rev. Entomol.* 19:161-176.

Schultze, W. 1923. Eleventh contribution to Coleoptera fauna of the Philippines. *Philipp. J. Sci* 23(6): 595-607.

Schultze, W. 1924. A monograph of the pachyrrynchid group of Brachyderinae, Curculionidae: Part I. *Philipp. J. Sci* 24(3): 309-360.

Stephen, CS. 1976. Philippine banana peel scarring weevils in Mindanao farms, Part II.

Unpublished report. Mindanao Fruits Co., Tagum, Davao. p.2.

Weevils Attacking Coconut

Rynchophorus ferrugineus, *R. palmarum* & *R. vulneratus*



Rynchophorus ferrugineus (Olivier, 1790)

Other Scientific Names

Calandra ferruginea Fabricius, 1801

Curculio ferrugineus Olivier, 1790

Rynchophorus signaticollis Chevrolat, 1882

International Common Names

English: Asiatic palm weevil; coconut weevil; red stripe weevil, red palm weevil

Spanish: picudo asiático de la palma

French: charançon asiatique du palmier

EPPO code

RHYCFE (*Rynchophorus ferrugineus*)

Distribution

Asia - Bahrain, Bangladesh, China, Georgia, India, Indonesia, Iran, Israel, Japan, Jordan, Kuwait, Laos, Lebanon, Malaysia, Myanmar, Oman, Pakistan, Philippines, Qatar, Saudi Arabia, Singapore, Taiwan, Thailand, Turkey, United Arab Emirates, Yemen

Africa - Egypt, Libya, Morocco, Spain, Tunisia

North America - California (eradicated)

Central America & Caribbean - Aruba, Curacao, Netherlands Antilles

Europe - Albania, Croatia, Cyprus, France, Greece, Italy, Malta, Portugal, Slovenia, Spain

Oceania - Australia, Papua New Guinea, Samoa, Solomon Is, Vanatu

Host Plants

Main host plant: palms (Arecaceae) - *Areca catechu* (betelnut palm), *Arenga pinnata* (sugar palm), *Borassus flabellifer* (toddy palm), *Cocos nucifera* (coconut), *Elaeis guineensis* (African oil palm), *Metroxylon sagu* (sago palm), *Phoenix canariensis* (Canary Island date palm), *Phoenix dactylifera* (date-palm), *Washingtonia* (Washington-palm)

& non-palm: *Calamus merillii* (rattan) Philippines (Braza, 1988)

***Rhynchophorus palmarum* (Linnaeus, 1758)**

Other Scientific Names

Calandra palmarum (Linnaeus) 1801
Cordyle barbirostris Thunberg, 1797
Cordyle palmarum (Linnaeus) 1797
Curculio palmarum Linnaeus, 1758
Rhynchophorus barbirostris (Thunberg)
Rhynchophorus cycadis Erichson, 1847
Rhynchophorus depressus Chevrolat, 1880
Rhynchophorus languinosus Chevrolat, 1880

International Common Names

English: South American palm weevil, palm weevil; palm-marrow weevil
Spanish: casanga; gorgojo cigarrón; gorgojo cigarrón del cocotero; gorgojo prieto de la palma; gualpa; mayate prieto del cocotero; picudo de la palma de coco; picudo del cocotero; picudo negro de la palma
French: charançon du palmier

EPPO code

RHYCPA (*Rhynchophorus palmarum*)

Distribution

North America - Mexico, USA - Arizona, California, Texas
Central America & Caribbean - Barbados, Belize, Costa Rica, Cuba, Dominica, Dominican Republic, El Salvador, Grenada, Guadeloupe, Guatemala, Honduras, Martinique, Nicaragua, Panama, Puerto Rico, St. Lucia, St Vincent and the Grenadines, Trinidad and Tobago
South America - Argentina, Bolivia, Brazil, Columbia, Ecuador, French Guiana, Guyana Paraguay, Peru, Suriname, Uruguay, Venezuela

Host Plants

Main host plant: *Cocos nucifera* (coconut), *Elaeis guineensis* (African oil palm), *Euterpe edulis*, *Metroxylon sagu* (sago palm), *Phoenix canariensis* (Canary Island date palm), *Phoenix dactylifera* (date-palm), *Saccharum officinarum* (sugarcane)

Non-significant hosts (adult feeding only) are *Ananas comosus* (pineapple), *Annona reticulata* (atis), *Artocarpus altilis* (breadfruit), *Carica papaya* (papaya), *Citrus spp.*, *Mangifera indica* (mango), *Musa spp.*, *Persea americana* (avocado), *Psidium guajava* (guava), *Theobroma cacao* (cacao).

Biology & Ecology

Life cycle

- Eggs hatch after 3.2 ± 0.93 days.
- Larvae have between six and 10 instars over a period of 52.0 ± 10.0 days.
- Prepupal stage lasts 4–17 days, during which the larvae make a cocoon using vegetative fibres.
- Pupal metamorphosis period lasts for 8 –23 days.
- Newly enclosed adults remain in the cocoon for 7.8 ± 3.4 days

Adult habits & behaviour

- Fecundity average 245 ± 155 eggs during a period of 30.7 ± 14.3 days; max- 697-718 eggs
- post-oviposition period of 10 days before the weevil dies.
- longevity : male 44.7 ± 17.2 days and females 40.7 ± 15.5 days
- In Egypt, has three generations per year, the shortest generation (first) of 100.5 days and the longest (third) of 127.8 days

Natural enemy

- Nematode parasite of larvae - *Billaea brasiliensis*, *Paratheresia menezesi*, *Paratheresia rhynch*

***Rynchophorus vulneratus* (Panzer, 1798)**

Other Scientific Names

Curculio vulneratus Panzer, 1798

Calandra schach Fabricius, 1801

Rynchophorus pascha Boheman in Schönherr, 1845

Rynchophorus ferrugineus v. *tenuirostris* Chevrolat, 1882

Rynchophorus glabrostris Schaufuss, 1885

Hallett et al (2004) - synonym of *R. ferrugineus*

Rugman et al (2013) - *R. vulneratus* is valid species

International Common Names

English: Asiatic palm weevil; coconut weevil; red stripe weevil, red palm weevil

Spanish: picudo asiático de la palma

French: charançon asiatique du palmier

EPPO code

Distribution

Asia - Indonesia, Malaysia, Singapore, Thailand, Philippines (unconfirmed)
North America - California (eradicated)

Host Plants

Main host plant: palms (Arecaceae) - *Cocos nucifera* (coconut), *Elaeis guineensis* (African oil palm), *Metroxylon sagu* (sago palm)

Control of Palm Weevils

“Integrated pest management for *R. ferrugineus* has been developed and tested in coconut palms in India (Kurian et al., 1976; Sathiamma et al., 1982, Abraham et al., 1999). Included in the IPM programme were cultural measures such as plant and field sanitation; physical methods by preventing entry of weevils through cut ends of petioles and wounds; and use of attractants and other chemicals (including filling of leaf axils with gamma BHC and sand as a preventive measure). Abraham et al. (1989) found the IPM approach very effective in reducing the number of infested palms in Kerala, India. Abraham et al. (1998) suggested that the major components of the IPM strategy for *R. ferrugineus* are surveillance, trapping the weevil using pheromones lures, detecting infestation by examination of palms, eliminating hidden breeding sites, clearing abandoned gardens, maintaining crop and field sanitation, using preventive chemical treatments, curative chemical control, implementing quarantine measures, training and education. In the Al Qatif region of Saudi Arabia, Vidyasagar et al. (2000a) successfully developed an IPM programme which, in addition to mass pheromone trapping, included a survey of all the cultivated gardens, systematic checking of all palms for infestation, periodic soaking of palms, and mass removal of neglected farms. A review of control strategies and IPM for the weevil were also presented by various other authors (Ramachandran, 1998; Nair et al., 1998; Murphy and Brisco, 1999). Faleiro (2006) has reviewed the issues and management of *R. ferrugineus* in coconut and date palm over the past 100 years.” (CABI, 2010)

Sterile Backcrosses /Sterile Insect Technique /Chemosterilization

“Ramachandran (1991) reported the effects of gamma radiation on *R. ferrugineus* whereby production of viable eggs decreased with increasing radiation dose, although there was no apparent effect on the F2 generation. Rahalker et al. (1973) reported that treatment of 1-2-day-old males of the weevil at a dose of 1.5 krad (15 Gy) resulted in 90% sterility with no adverse effect on survival. Treatment of higher doses increased sterility but reduced survival. A ratio of ten treated males to one normal one was needed for appreciable suppression of progeny production. Using chemosterilants Rahalkar et al. (1975) reported that treatment of male weevils with metepa or hempa did not result in a satisfactory level of sterility without adversely affecting their survival. However, metepa was more toxic than hempa.” (CABI, 2010)

References

- Abe F, Hata K, Sone K, 2009. Life history of the red palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Dryophthoridae), in southern Japan. *Florida Entomologist*, 92(3):421-425.
- Abraham VA, 1971. Note on an effective method of preventing entry by red-weevil, *Rhynchophorus ferrugineus* Fabricius (Curculionidae: Coleoptera), into the stem of coconut palm through cut petioles. *Indian Journal of Agricultural Sciences*, 41(12):1130-1131
- Abraham VA, Koya KMA, Kurian C, 1989. Integrated management of red palm weevil (*Rhynchophorus ferrugineus* F.) in coconut gardens. *Journal of Plantation Crops*, 16(Supplement):159-162

- Abraham VA, Nair SS, Nair CPR, 1999. A comparative study on the efficacy of pheromone lures in trapping red palm weevil, *Rhynchophorus ferrugineus* Oliv. (Coleoptera : Curculionidae) in coconut gardens. *Indian Coconut Journal (Cochin)*, 30(7):1-2; 6 ref.
- Braza RD, 1988. Asiatic palm weevil destroys rattan too. *Canopy International*. 13(6):6.
- CABI/EPPO, 2010. *Rhynchophorus ferrugineus*. [Distribution map]. *Distribution Maps of Plant Pests*, No. June. Wallingford, UK: CABI, Map 258 (3rd revision).
- Copeland EB, 1931. *The coconut*. London, UK: Macmillan and Co., Ltd.
- Dangar TK, 1997. Infection of red palm weevil, *Rhynchophorus ferrugineus*, by a yeast. *Journal of Plantation Crops*, 25(2):193-196
- Dhileepan K, 1991. Insects associated with oil palm in India. *FAO Plant Protection Bulletin*, 39(2-3):94-99.
- El-Ezaby FA, 1997. Injection as a method to control the Red Indian date palm weevil *Rhynchophorus ferrugineus*. *Arab Journal of Plant Protection*, 15(1):31-38.
- El-Garhy ME, 1996. Field evaluation of the aggregation pheromone of the red palm weevil, *Rhynchophorus ferrugineus*, in Egypt. *Brighton Crop Protection Conference: Pests & Diseases -1996: Volume 3: Proceedings of an International Conference*, Brighton, UK, 18-21 November 1996., 1059-1064
- EPPO, 2007. *Rhynchophorus ferrugineus* and *Rhynchophorus palmarum*. *Bulletin OEPP/*
- Faghih AA, 1996. The biology of red palm weevil, *Rhynchophorus ferrugineus* Oliv. (Coleoptera, Curculionidae) in Saravan region (Sistan & Balouchistan province, Iran). *Applied Entomology and Phytopathology*, 63(1/2):16-18
- Faleiro JR, 2006. A review of the issues and management of the red palm weevil *Rhynchophorus ferrugineus* (Coleoptera: Rhynchophoridae) in coconut and date palm during the last one hundred years. *International Journal of Tropical Insect Science*, 26(3):135-154.
- Faleiro JR, Abraham VA, Nabil Noudi, Al Shuaibi MA, Kumar TP, 2000. Field evaluation of red palm weevil, *Rhynchophorus ferrugineus* Oliv. pheromone (Ferrugineol) lures. *Indian Journal of Entomology*, 62(4):427-433.
- Hagley E (1965) On the life history and habits of *Rhynchophorus palmarum*. *Annals of the Entomological Society of America* 58, 22–28.
- Hallett, R.H., Crespi, B.J., Borden, J.H. 2004. Synonymy of *Rhynchophorus ferrugineus* (Olivier), 1790 and *R. vulneratus* (Panzer), 1798 (Coleoptera, Curculionidae, Rhynchophorinae). *J. Nat. Hist.* 38:2863-2882
- Hallett RH, Oehlschlager AC, Borden JH, 1999. Pheromone trapping protocols for the Asian palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae). *International Journal of Pest Management*, 45(3):231-237
- IPPC, 2007. Red Palm Weevil. IPPC Official Pest Report, No. TR-3/1. Rome, Italy: FAO. <https://www.ippc.int/IPP/En/default.jsp>

- Jaffé K, Sánchez P, Cerda H, Hernández JV, Jaffé R, Urdaneta N, Guerra G, Martínez R & Miras B (1993) Chemical ecology of *Rhynchophorus palmarum*: attraction to host plants and to a male-produced aggregation pheromone. *Journal of Chemical Ecology* 19, 1703–1720.
- Jaya S, Suresh T, Sobhitha-Rani RS, Sreekumar S, 2000. Evidence of seven larval instars in the red palm weevil, *Rhynchophorus ferrugineus* Oliv. reared on sugarcane. *Journal of Entomological Research*, 24(1):27-31.
- Justin CGL, Leelamathi M, Thangaselvabai T, Johnson SBN, 2008. Bioecology and management of the red palm weevil, *Rhynchophorus ferrugineus* Oliv. (Coleoptera: Curculionidae) on coconut - a review. *Agricultural Reviews*, 29(2):117-124.
- Kurian C, Antony J, Abraham VA, Natarajan P, 1976. Pest management in coconut gardens. An integrated approach. *Indian Farming*, 7(9):31-35.
- Kurian C, Mathen K, 1971. Red palm weevil - hidden enemy of coconut palm. *Indian Farming*, 21(1):29-31
- Li YueZhong, Zhu ZengRong, Ju RuiTing, Wang LianSheng, 2009. The red palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae), newly reported from Zhejiang, China and update of geographical distribution. *Florida Entomologist*, 92(2):386-387.
- Menon KPV, Pandalai KM, 1960. Pests. In: *The Coconut palm. A Monograph*. Inrankulam, South India: Indian Central Committee, 261-265.
- Misra RM, 1998. Insect pests of oil palm (*Elaeis guineensis* L.) from India and their management. *Indian Journal of Forestry*, 21(3):259-263.
- Muralidharan CM, Vaghasia UR, Sodagar NN, 1999. Population, food preference and trapping using aggregation pheromone (ferrugineol) on red palm weevil (*Rhynchophorus ferrugineus*). *Indian Journal of Agricultural Sciences*, 69(8):602-604.
- Murphy ST, Briscoe BR, 1999. The red palm weevil as an alien invasive: biology and the prospects for biological control as a component of IPM. *Biocontrol News and Information*, 20(1): 35-46
- Nair CPR, Sathiamma B, Chandrika Mohan, Murali Gopal, 1998. Newer approaches in the integrated pest management in coconut. *Indian Coconut Journal (Cochin)*, 29(4):99-103.
- Prabhu ST, Patil RS, 2009. Studies on the biological aspects of red palm weevil, *Rhynchophorus ferrugineus* (Oliv.). *Karnataka Journal of Agricultural Sciences*, 22(3):732-733.
- Rahalkar GW, Harwalkar MR, Rananavare HD, Shantaram K, Ayengar ARG, 1973. Laboratory studies on radiation sterilization of the red palm weevil (*Rhynchophorus ferrugineus* Oliv.) males. In: Nayar NM, ed. *Journal of Plantation Crops*, 1:141-145
- Rahalkar GW, Harwalkar MR, Rananavare HO, 1975. Laboratory studies on sterilization of the male red palm weevil, *Rhynchophorus ferrugineus* Oliv. International Atomic Energy Agency; Food and Agriculture Organization: Sterility principle for insect control 1974. Proceedings of the symposium on the sterility principle for insect control jointly organized by the IAEA and the FAO of the United Nations and held in Innsbruck, 22-26 July 1974. International Atomic Energy Agency. Vienna Austria, 261-267

- Ramachandran CP, 1991. Effects of gamma radiation on various stages of red palm weevil, *Rhynchophorus ferrugineus* F. *Journal of Nuclear Agriculture and Biology*, 20(3):218-221
- Ramachandran CP, 1998. The red palm weevil *Rhynchophorus ferrugineus* F.: a review and future strategies. *Indian Coconut Journal (Cochin)*, 29(4):104-106; 25 ref.
- Reginald C, 1973. *Principal Insect Pests*. In: *Coconuts. Tropical Agriculture Series*, London, UK:Longmans.
- Rugman-Jones PF, Hoddle CD, Hoddle MS, Stouthamer R, 2013. The lesser of two weevils: molecular-genetics of pest palm weevil populations confirm *Rhynchophorus vulneratus* (Panzer 1798) as a valid species distinct from *R. ferrugineus* (Olivier 1790), and reveal the global extent of both. *PLoS ONE*, 8(10):e78379. <http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0078379>
- Sacchetti P, Camèra A, Granchietti A, Rosi MC, Marzialetti P, 2006. Identification, biology and spread of the Red Palm Weevil *Rhynchophorus ferrugineus* (Olivier) in Italy. (Identificazione, biologia e diffusione del curculionide delle palme, *Rhynchophorus ferrugineus* (Olivier).) *Informatore Fitopatologico*, 56(6):35-40.
- Salama HS, Zaki FN, Abdel-Razek AS, 2009. Ecological and biological studies on the red palm weevil *Rhynchophorus ferrugineus* (Olivier). *Archives of Phytopathology and Plant Protection*, 42(4):392-399.
- Sathiamma B, Abraham VA, Kurian C, 1982. Integrated pest management of the major pests of coconut. *Indian Coconut Journal*, 12(6/9):27-29
- Vidyasagar PSPV, Al-Saihati AA, Al-Mohanna OE, Subbei AI, Abdul Mohsin AM, 2000. Management of red palm weevil *Rhynchophorus ferrugineus* Oliv., a serious pest of date palm in Al Qatif, Kingdom of Saudi Arabia. *Journal of Plantation Crops*, 28(1):35-43.
- Vidyasagar PSPV, Mohammed Hagi, Abozuhairah RA, Al-Mohanna OE, Al-Saihati AA, 2000. Impact of mass pheromone trapping on red palm weevil: adult population and infestation level in date palm gardens of Saudi Arabia. *Planter*, 76(891):347-355.
- Wang Feng, Ju RuiTing, Li YueZhong, Xu Ying, Du YuZhou, Zhou JianMing, 2009. Lab observation for biological characteristics and morphology of *Rhynchophorus ferrugineus*. *Chinese Bulletin of Entomology*, 46(4):556-560.
- Wattanapongsiri A, 1966. A revision of the Genera *Rhynchophorus* and *Dynamis*. *Department of Agricultural Science Bulletin*. Bangkok, Thailand.
- Wilson M (1963) Investigations into the development of the palm weevil, *Rhynchophorus palmarum*. *Tropical Agriculture Trinidad* 40, 185–196.

Weevils Attacking Sweet Potato



Cylas formicarius (Fabricius, 1798)

Other Scientific Names

Brentus formicarius Fabricius, 1798
Cylas elegantulus (Summers, 1875)
Cylas formicarius elegantulus Summers
Cylas formicarius formicarius
Cylas turcipennis Boheman
Otidocephalus elegantulus Summers, 1895

International Common Names

English: sweet potato borer, sweet potato weevil
 Spanish: gorgojo del camote; piogán de la batata (Dominican Republic);
 tetuán del boniato (Cuba)
 French: charançon de la patate douce; charançon faux-fourmi

EPPO code

CYLAFO (*Cylas formicarius*)

Distribution

Asia - Bangladesh, British Indian Ocean, Brunei Darussalam, Cambodia, China, Christmas Is, Cocos Is, India, Indonesia, Japan, Laos, Malaysia, Maldives, Myanmar, Pakistan, Philippines, Singapore, Sri Lanka, Taiwan, Thailand, Vietnam
Africa - Cameroon, Chad, Congo, Ethiopia, Ghana, Kenya, Liberia, Libya, Madagascar, Mauritius, Mozambique, Reunion, Senegal, Seychelles, Somalia, S Africa, Sudan, Swaziland, Tanzania, Uganda, Zimbabwe
North America - Mexico, USA - Alabama, Arkansas, Florida, Georgia, Hawaii, Louisiana, Mississippi, New Mexico, N Carolina, S. Carolina, Texas
Central America & Caribbean - Anguilla, Antigua & Barbuda, Bahamas, Belize, Cayman Is, Cuba, Dominican Republic, Guadeloupe, Guatemala, Haiti, Jamaica, Netherlands Antilles, Puerto Rico, St Lucia, Trinidad and Tobago, US Virgin Islands
South America - Guyana & Venezuela
Oceania - American Samoa, Australia, Cook Is, Fiji, French Polynesia, Guam, Kiribati, Marshall Is, Micronesia, New Caledonia, Palau, Papua New Guinea, Samoa, Solomon Is, Tonga, Tuvalu, Vanuatu, Wallis & Futuna Is

Host Plants

Main host plant: *Calystegia sepium* (great bindweed), *Colocasia esculenta* (taro), *Cuscuta* (dodder), *Ipomoea* (morning glory), *Ipomoea aquatica* (swamp morning-glory), *Ipomoea batatas* (sweet potato), *Ipomoea cairica* (five-fingered morningglory), *Ipomoea pes-caprae* (beach morning glory), *Ipomoea purpurea* (tall morning glory), *Ipomoea quamoclit* (Cupid's-flower), *Jacquemontia tamnifolia* (Smallflower morningglory), *Pharbitis nil* (Japanese morning glory)

Biology and Ecology

Life cycle

- Eggs are laid singly in cavities in the root or stems.
- Egg incubation period ranges from 4 days at 30°C to 7.9 days at 20°C or 4 to 56 days at mean temperatures of 20 and 10.5°C, respectively.
- Larvae feed inside roots or stems for 25-35 days during which they complete three larval instars. In USA, larval development in the field ranges 12-154 days.
- Pupation takes place within the sweet potato roots or stems where larvae feed.
- Pupal period lasts 4-8 days or 5 to 10.7 days at 25 and 20°C, respectively.
- Soon after emergence from the pupa, the adult stays in the pupal chamber and then cuts its way through the plant tissue.
- Adults mate soon after emergence but pre - oviposition period has a minimum of 4.5 days at 30°C or 7.7 days at 20°C.
- Egg to egg development time - 84.7, 33, and 33.7 days at 20, 27 and 30°C, respectively.
- Life cycle duration of 36-43 days under undefined conditions.

Adult behaviour & habits

- Longevity - maximum of 113 days (Indonesia), males 63-120 days and females 81-107 days (Philippines), males 94 days and 1females 09 days (India)
- Fecundity - 90-340 eggs per female, (Philippines), 185 eggs (Indonesia), 1-319 eggs (USA) with an average of 119 and 97-216 eggs (India).

Natural Enemy

- Fungal pathogen of adults - *Beauveria*, *Metarrhizium*
- Wasp parasitoid of larvae - *Bracon mellitor*, *B. cylasovoros*, *B. punctatus*, *Drapetis exilis*, *Euderus purpureas*
- Nematode parasite of larvae - *Heterorhabditis bacteriophora*, *Heterorhabditis heliothidis*, *Metapelma spectabile*, *Rhaconotus menippus*
- Ant Predator of adult - *Pheidole megacephala*

Prevention and control

Cultural Control - crop rotation, intercropping, mulching, sanitation, destruction of crop residues, flooding of infested fields, use clean cuttings as planting materials and removal of alternative hosts. Planting cuttings deep in the soil, use of deep-rooted cultivars, and harvesting the crop as soon as it has developed roots of acceptable size (Edwards, 1930; Holdaway, 1941; Sherman and Tamashiro, 1954; Sutherland, 1986a).

Host-Plant Resistance - no resistant variety

Chemical Control - pre-plant application to dip vines for planting; post-plant application is difficult with conventional spraying, dusting, fumigation or side-dressing of insecticide granules with presently available insecticides, once weevils are present within the crown or the tuberous root. Control achieved by post-plant applications appears to be due to mortality of weevil adults searching for feeding or oviposition sites.

Sex pheromone - female sex pheromone isolated, identified and synthesized the chemical (Z)-3-dodecen-1-ol(E)-2-butenate. This chemical has great potential for attracting male sweet potato weevils (Proshold et al., 1986; Jansson et al., 1992) and reducing the weevil populations in the field (Talekar and Lee, 1989).

Biological Control - *Beauveria bassiana* is produced in large quantities and used intensively for the control of the sweet potato weevil in Cuba. Sprays of the fungus have largely replaced the use of insecticides (Castellón et al., 1992).

References

Ahmad M, Hossain MB, 1979. Notes on weevils (Curculionidae: Coleoptera) of the Bangladesh Agricultural University area, Mymensingh. Bangladesh Journal of Zoology, 7(1):61-66

Alcazar J, Cisneros F, Morales A, 1997. Large-scale Implementation of the IPM for Sweetpotato Weevil in Cuba: A collaborative effort. Lima, Peru: International Potato Center, 185-190.

APPPC, 1987. Insect pests of economic significance affecting major crops of the countries in Asia and the Pacific region. Technical Document No. 135. Bangkok, Thailand: Regional Office for Asia and the Pacific region (RAPA).

Braun AR, Fliert Evan de, 1999. Evaluation of the impact of sweetpotato weevil (*Cylas formicarius*) and of the effectiveness of *Cylas* sex pheromone traps at the farm level in Indonesia. International Journal of Pest Management, 45(2):101-110; 26 ref.

CABI/EPPO, 2004. *Cylas formicarius*. Distribution Maps of Plant Pests, No. 278. Wallingford, UK: CAB International

Cockerham KL, 1943. The host preference of the sweetpotato weevil. Journal of Economic Entomology, 36:471-472.

Cockerham KL, Deen OT, Christian MB, Newsom LT, 1954. The biology of the sweet potato weevil. Technical Bulletin Louisiana Agricultural Experiment Station No. 483.

Coffelt JA, Vick KW, Sower LL, McClellan WTM, 1978. Sex pheromone of the sweetpotato weevil, *Cylas formicarius elegantulus* laboratory bioassay and evidence for a multiple component system. *Environmental Entomology*, 7(5):756-758

Gonzales SS, 1925. The sweet potato weevil (*Cylas formicarius*, Fabr). *Philippine Agriculturist*, 14:257-281.

Heath RR, Coffelt JA, Sonnet PE, Proshold FI, Dueben B, Tumlinson JH, 1986. Identification of sex pheromone produced by female sweetpotato weevil, *Cylas formicarius elegantulus* (Summers). *Journal of Chemical Ecology*, 12(6):1489-1503

Jansson RK, 1996. Infectivity and reproduction of three heterorhabditid nematodes (Rhabditida:

Heterorhabditidae) in two insect hosts. *Florida Entomologist*, 79(3):363-373; 18 ref.

Jansson RK, Bryan HH, Sorensen KA, 1987. Within-vine distribution and damage of sweetpotato weevil, *Cylas formicarius elegantulus* (Coleoptera: Curculionidae), on four cultivars of sweet potato in southern Florida. *Florida Entomologist*, 70(4):523-526

Jayaramaiah M, 1975. Bionomics of sweet potato weevil *Cylas formicarius* (Fabricius) (Coleoptera: Curculionidae). *Mysore Journal of Agricultural Sciences*, 9(1):99-109

Jayaramaiah M, 1975. Reaction of sweet potato varieties to the damage of the weevil, *Cylas formicarius* (Fab.) (Coleoptera: Curculionidae) and on the possibility of picking up of infestation by weevil. *Mysore Journal of Agricultural Sciences*, 9(3):418-421

Li ZZ, 1998. Control of *Cylas formicarius* Fabricius with pheromone. *Plant Protection*, 24(5): 28-30.

Miyatake T, Moriya S, Kohama T, Shimoji Y, 1997. Dispersal potential of male *Cylas formicarius* (Coleoptera: Brentidae) over land and water. *Environmental Entomology*, 26(2): 272-276; 23 ref.

Morales A, 1988. El cultivo del boniato (*Ipomoea batatas* (L.) Lam.) y su principal enemigo al *Cylas formicarius* var. *elegantulus*. In: *Proceedings of a Sweet Potato Workshop*, International Potato Center (CIP), Santo Domingo, Dominican Republic. Dominican Republic: CIP.

Moriya S, Hiroyoshi S, 1998. Flight and locomotion activity of the sweetpotato weevil (Coleoptera: Brentidae) in relation to adult age, mating status, and starvation. *Journal of Economic Entomology*, 91(2):439-443; 24 ref.

Mullen MA, 1981. Sweetpotato weevil, *Cylas formicarius elegantulus* (Summers): development, fecundity, and longevity. *Annals of the Entomological Society of America*, 74(5):478-481.

Nottingham SF, Son KC, Severson RF, Arrendale RF, Kays SJ, 1989. Attraction of adult sweet potato weevils, *Cylas formicarius elegantulus* (Summers), (Coleoptera: Curculionidae), to sweet potato leaf and root volatiles. *Journal of Chemical Ecology*, 15(3):1095-1106

- Palaniswami MS, Mohandas N, Saraswathy P, 1991. On farm survey on the incidence, extent and intensity of *Cylas formicarius* F. damage on sweet potato in Kerala. *Journal of Root Crops*, 17(Special issue):180-196; 8 ref.
- Proshold FI, Gonzalez JL, Asencio C, Heath RR, 1986. A trap for monitoring the sweetpotato weevil (Coleoptera: Curculionidae) using pheromone or live females as bait. *Journal of Economic Entomology*, 79(3):641-647
- Sato K, Uritani I, Saito T, 1982. Properties of terpene-inducing factor extracted from adults of the sweet potato weevil, *Cylas formicarius* Fabricius (Coleoptera: Brentidae). *Applied Entomology and Zoology*, 17(3):368-374
- Sherman M, Tamashiro M, 1954. The sweetpotato weevils in Hawaii, their biology and control. Hawaii Agricultural Experiment Station Technical Bulletin No. 23
- Shimizu T, Moriya S, 1996. Daily locomotor activity in the Indian sweet potato weevil, *Euscepes postfasciatus* (Fairmaire) (Coleoptera: Curculionidae) and sweet potato weevil, *Cylas formicarius* (Fabricius) (Coleoptera: Brentidae) monitored by an actograph system. *Applied Entomology and Zoology*, 31(4):626-628; 3 ref.
- Shimizu T, Moriya S, 1996. Flight time and flight age in the sweet potato weevil, *Cylas formicarius* (Fabricius) (Coleoptera: Brentidae). *Applied Entomology and Zoology*, 31(4):575-580; 13 ref.
- Singh B, Yazdani SS, Singh R, Hameed SF, 1984. Effect of intercropping on the incidence of sweet potato weevil, *Cylas formicarius* Fabr., in sweet potato (*Ipomoea batatas* Lam.). *Journal of Entomological Research*, 8(2):193-195
- Smit NEJM, 1997. The effect of the indigenous cultural practices of in-ground storage and piecemeal harvesting of sweetpotato on yield and quality losses caused by sweetpotato weevil in Uganda. *Agriculture, Ecosystems & Environment*, 64(3):191-200; 23 ref.
- Sorensen KA, 1987. Cultural, regulational and educational programmes on the sweetpotato weevil in the United States. *Insect Science and its Application*, 8(4-6):825-830
- Soumik Mondal, Tarafdar J, Sen H, 2006. Use of sex pheromone trap for monitoring and management of sweet potato weevil (*Cylas formicarius*, Fabricius). *Environment and Ecology*, 24S(Special 3):525-528.
- Suenaga T, Setokuchi O, Sakp K, 1987. Distribution of the sweet potato weevil *Cylas formicarius* Fabricius and the small sweet potato weevil, *Euscepes postfasciatus* Fairmaire in the Amami Islands. *Proceedings of the Association for Plant Protection of Kyushu*, No. 33:116-118
- Sugimoto T, Sakuratani Y, Fukui H, Kiritani K, Okada T, 1996. Estimating the reproductive properties of the sweet potato weevil, *Cylas formicarius* (Fabricius) (Coleoptera, Brentidae). *Applied Entomology and Zoology*, 31(3):357-367; 30 ref.
- Sutherland JA, 1985. The chemical control of *Cylas formicarius* (Fab.) using insecticides at low volume rates, with some considerations on the economics of control. *Journal of Plant Protection in the Tropics*, 2(2):97-103
- Sutherland JA, 1986a. A review of the biology and control of the sweet potato weevil, *Cylas formicarius* (Fab.). *Tropical Pest Management*, 32:304-315.
- Swindale A, 1992. Measurement of the economic importance of sweetpotato weevil.
- Unpublished discussion paper 13. Lima, Peru: International Potato Center.

Talekar NS, 1983. Infestation of a sweetpotato weevil (Coleoptera: Curculionidae) as influenced by pest management techniques. *Journal of Economic Entomology*, 76(2):342-344

Talekar NS, 1987. Feasibility of the use of resistant cultivar in sweetpotato weevil control. *Insect Science and its Application*, 8(4-6):815-817

Talekar NS, Lai RM, Cheng KW, 1989. Integrated control of sweetpotato weevil at Penghu Island Plant Protection Bulletin, Taiwan, 31(2):185-191.

Talekar NS, Lee ST, 1989. Studies on the utilization of a female sex pheromone for the management of sweetpotato weevil, *Cylas formicarius* (F.) (Coleoptera: Curculionidae). *Bulletin of the Institute of Zoology, Academia Sinica*, 28(4):281-288

Trehan KN, Bagal SR, 1958. Life-history, bionomics, and control of sweet potato weevil (*Cylas formicarius*) with short notes on some other pests of sweet potato in Bombay State. *Indian Journal of Entomology*, 19:245-252.

Waterhouse DF, 1993. The Major Arthropod Pests and Weeds of Agriculture in Southeast Asia. ACIAR Monograph No. 21. Canberra, Australia: Australian Centre for International Agricultural Research, 141 pp.

Waterhouse DF, Norris KR, 1987. Biological control: Pacific prospects. viii + 454pp
Wilson DD, Son KC, Nottingham SF, Severson RF, Kays SJ, 1989. Characterization of an oviposition stimulant from the surface of sweet potato *Ipomoea batatas* storage roots for the sweet potato weevil, *Cylas formicarius elegantulus*. *Entomologia Experimentalis et Applicata*, 51(1):71-75

Wilson DD, Son KC, Severson RF, Kays SJ, 1990. Effect of a pentacyclic triterpene from sweet potato storage roots on oviposition by the sweetpotato weevil (Coleoptera: Curculionidae). *Environmental Entomology*, 19(6):1663-1665

Wolfe GW, 1991. The origin and dispersal of the pest species of *Cylas* with key to the pest species groups of the world. In: Jansson RK, Raman, KV, eds. *Sweet Potato Pest Management: A Global Perspective*. Boulder, USA: Westview Press.

Yasuda K, 1989. Infestation of the sweetpotato weevils (*Euscepes postfasciatus* Fairmaire and *Cylas formicarius* Fabricius) in sweet potato fields. I. Invasion with cutting. *Proceedings of the Association for Plant Protection of Kyushu*, 35:120-122



ASEAN Regional Training Workshop on Diagnostics of Weevils of Quarantine Importance

July 10-22, 2017

**Institute of Weed Science, Entomology and Plant Pathology
College of Agriculture and Food Science
University of the Philippines Los Baños**

SESSION 6

WEEVILS AS STORAGE PESTS AND THEIR CONTROL

Lecture Notes by:

DR. PIO A. JAVIER

**Institute of Weed Science, Entomology and Plant Pathology College of Agriculture
and Food Science
University of the Philippines Los Baños**



COMMON STORAGE STORAGE PESTS



Pio A. Javier, Ph.D.
Research Professor (Adjunct)
Institute of Weed Science, Entomology and
Plant Pathology
College of Agriculture & Food Science
UPLB, College, Laguna
(pio.javier@yahoo.com.ph) 0927-329-3893

Economic Importance of SPPs

- Among the pests that degrade and destroy food crops, insects are the most important during storage.
- Stored commodities that are free from insect infestation & damage is of utmost importance. In grains alone, millions of dollars could be lost due to insect damage.
- Damage estimates caused storage pests ranged from 5-10% weight loss of all foodstuffs
- Losses due to storage pests in corn in the Philippines:
 - 17% weight loss for 6 months storage
 - 43% weight loss for 12 months storage
 - due to mixed pop of corn weevil & red flour beetle
 - 42% weight loss in 13 NFA warehouses

Economic Importance

- **Corn weevil consume about 10 – 20 mg food**
- **Extremely rapid population increase = 100X in each generation of 1 month**
- **Reduces seed viability and quality**
- **Contamination of grains (dead insects & body parts) caused spoilage & loss of cash value**

INTRODUCTION

- **Stored commodities that are free from insect infestation & damage is of utmost importance. In grains alone, millions of dollars could be lost due to insect damage in storage**
- **In the Philippines, storage losses in rice & corn amount to 15% & 12%, respectively. The country imports 5-10% of its total grain reqmts. Prevention of storage losses could be a better strategy (proactive) to attain food security to complement increased food production in the farm.**

PREDOMINANT STORAGE INSECT PESTS

40 coleopteran families are reported attacking stored products worldwide

23 are of importance in the tropics

14 are involved in the major & minor storage pests in Southeast Asia

Anobiidae	Cleridae	Dermestidae	Nitidulidae
Anthribidae	Cucujidae	Lophacateridae	Silvanidae
Bruchidae	Scolytidae	Trogositidae	Tenebrionidae
Bostrichidae	Curculionidae		

Major Stored Product Coleopteran Families

1. Anobiidae

- Adults are small, subcylindrical (wood-boring, oval or nearly globular (general feeders) w/ prothorax more or less covering the deflexed head.
- With 11-segmented antennae w/ a loose 3-segmented serrate club.
- Elytra completely cover the abdomen
- With 5 visible abdominal sternites
- All tarsi are 5-segmented w/ segments 1-4 decreasing in length.
- Two species are important in storage:

* *Lasioderma serricorne*


* *Stegobium paniceum*

Cigarette Beetle, *Lasioderma serricorne* (F.)

- Beetle is 2-4 mm long, oval, reddish to dark brown, covered w/ fine hairs.
- Antennae about half as long as the body and have 11 segments
- Elytra smooth with very short hairs but w/o striae.
- Common pest of tobacco & tobacco products; also common in many spices & animal feed concentrates.
- Adult lives for 2–6 weeks producing about 100 eggs.
- Found in temperate, sub-tropical & tropical regions
- When disturbed, adult conceals its head under the large pronotum

Cigarette Beetle, *Lasioderma serricorne*

- Eggs are laid in crevices or folds in the substrate.
- Hatch in 5-6 days at 35°C.
- Newly hatched larvae are very active & often move around & bore into packaged goods.
- Fleshy larvae are distinctly hairy, pass through 4-6 instars.
- Larval period at 32.5°C ~ 16 day
- Larva pupates within a cell among the substrate or within the grain.
- Pupal devt. lasts for 4 d at 30–37.5°C.



Drugstore Beetle, *Stegobium paniceum* (L.)



- Light brown beetle, 2-2.5 mm long & slightly hairy elytra with longitudinal ridges
- Last 3 antennal segments form a large loosely segmented club
- At 20-25°C & 6-80% RH, female lays ~ 75 eggs w/ life span of ~40-90 d.
- W/ 4-6 larval instars. Full grown larva constructs a cocoon where it pupates.
- Adult cannot fly & dispersal depends on passive distribution during movement of goods.
- Cold-hardy & can survive winter conditions in temperate regions.
- Cosmopolitan pest but is more temperate than tropical.
- Common pest of stored products, most serious on processed foods, spices and drugs.
- An occasional pest of stored grains.

2. Anthribidae

- **Antennae are 11-segmented, thread-like or filiform with a loose 3-segmented club.**
- **All tarsi are 5-segmented.**
- **Mostly of tropical distribution, associated with old wood, dead branches & fungi.**
- **Only one species, *Araecerus fasciculatus*, is of economic importance on stored products**

Coffee Bean Beetle

Araecerus fasciculatus




- Most imp. insect pest of stored coffee; attacks coffee beans
- Larvae bore into the beans & once damaged the bean may then be damaged by secondary insect pests
- Also attacks cocoa beans, cassava chips & occasionally corn
- Adult measures 3-5 mm long
- Prothorax & elytra bear many patches of light colored setae giving a mottled appearance
- The 3 terminal segments of antennae are longer than the other segments forming a loose club
- Life cycle ~ 46-66 days; female lays ~50 eggs; adults live for ~ 17 weeks

3. Bostrichidae

- **Body is cylindrical and prothorax coarsely tuberculate concealing the deflexed head.**
- **Antennae are straight & terminate in a loose**
- **3- or 4-segmented club**
- **Apically, the elytra are somewhat flattened & sloped ventrally or less steeply. This sloping region is referred to as declivity. This is marked with carinae, tubercles or hooks which are useful recognition features**
- **Tarsi are all 5-segmented**
- **Cosmopolitan, mainly wood-boring beetles; attack timber of dry wood and occasionally, unhealthy trees.**
- **Three species frequently infest stored grains:**
 - * *Rhizopertha dominida*
 - * *Dinoderus minutus*
 - * *Prostephanus truncatus*

Lesser Grain Borer

Rhizopertha dominica



- 2.5-3 mm long, dark brown, body is cylindrical & head is concealed ventrally by the prothorax, which is dotted with many small tubercles
- Pronotum is rounded & there is transverse rows of teeth
- Elytra w/ regular rows of punctures & short setae that curve posteriorly
- Cosmopolitan but of particular importance in tropical & subtropical regions
- Bores & feeds on a wide variety of foods, chiefly cereals, including grains of all kinds, dried cassava, flour, macaroni, beans, dried potato and many other materials. Adults are surface feeders while larvae are primarily internal feeders
- Attacks rough rice which is resistant to attack of most storage pests
- Female lays 200-600 eggs, which hatches in 7-8 days
- Larval development on whole meal at 28°C and 70% RH is 27-31 days.


Larger Grain Beetle

Prostephanus truncatus



- Serious pest of stored corn and dried cassava roots, & appears to attack corn in the field just before harvest. Adults bore into a wide range of foodstuffs & some other materials, e.g. wood
- Development from egg to adult in corn at 27°C & 70% RH is 32.5 days
- Introduced to Tanzania in late 1970 & has since spread into corn & cassava growing areas of tropical Africa
- Known to occur in South America, Mexico, Central America and the extreme south of USA
- Absent in Southeast Asia; important as a quarantine pest
- The declivity of elytra is flattened & steep & with many tubercles over its surface
- The limits of declivity, apically and laterally are marked by a carina
- Antennae are 10-segmented & have a loose 3-segmented club
- Body is 3-4.5 mm long

Powder Post Beetle, *Dinoderus minutus* (F.)




- Closely resembles *Rhizopertha* and *Prostephanus* except for the following:
 - there is a pair of shallow depressions at the base of the pronotum
 - scutellum is transverse, its width about twice its length. Elytral hairs are short & erect.
- Infests dried cassava, bamboo & wood. Can also attack cocoa beans, starch, sago and corn.

4. Bruchidae

- Clothed with hairs, w/c are usually colored forming distinct patterns
- Elytra do not cover the last abdominal segment
- Adults are small, strongly convex beetles.
- Antennae are relatively long; eyes are large (except genus *Caryedon*)
- Antennae are 11-segmented without a club, usually filiform w/ some segment which are compressed in the males of some *Callosobruchus*
- Tarsi are distinctly 4-segmented w/ the 3rd segment broadly dilated and deeply emarginate
- Hind femur are somewhat thickened or strongly enlarged and each bear one or more teeth
- Upon hatching, bruchid grubs have 3 thoracic legs but after the first molt they become apodous.
- Bean beetles do not possess the elongate rostrum of true weevils. All species that are impt as pest of stored products are legume feeders.
- Common species: *Acanthoscelides obtectus*, *Callosobruchus chinensis* and *C. maculatus*

Cowpea Bean Beetle


Callosobruchus chinensis



- **Adult are 2.5-3.5 mm in length with typical white markings on the scutellum.**
- **Has a pair of distinct ridges (inner & outer) on the ventral side of each hind femur. Each ridge has a tooth near the apical end. Inner tooth is slender, rather parallel-sided, & equal to the outer tooth.**
- **Antennae are pectinate in male & serrated in female.**
- **Elytra are pale brown w/ small median dark marks and larger posterior dark patches, w/c may merge to make the entire posterior part of the elytra dark**
- **Primary pest of legumes in tropical & subtropical climates, particularly chickpeas, Adzuki beans & cowpeas**

Southern Cowpea Bean Beetle

Callosobruchus maculatus



- Females have two large lateral dark patches midway along the elytra and smaller patches at the anterior and posterior ends, leaving a paler brown cross-shaped area covering the rest.
- Males are brown all over with small blackish markings.

Bean Beetle

Acanthoscelides obtectus



Hind
femur

- Cosmopolitan species that is a serious primary pest of common bean & lima beans. Also attacks other legumes.
- Beetle 3-4.5 mm long; elytra variegated w/ yellowish & dark brown patches of hairs; elytra do not completely cover the abdomen.
- Body colors are grey, brown or reddish brown w/o any distinctive pattern.
- Each hind femur has an inner ventral ridge near the apex, a large tooth followed by 2 or 3 smaller, sharp teeth
- Serrated antennae are dark grey except for segments 1-5 & 11, w/c are reddish.
- Legs are reddish except for the ventral half of hind & middle femur which is black.

Bean Beetle

Acanthoscelides obtectus



- Eggs are laid on ripening pods while in the field.
- Eggs are laid loosely w/in pod cavity or among stored seeds
- Newly hatched larva develops entirely w/in a single seed eventually forming a pupation chamber under the seed coat
- Adult that emerges from the chamber is short-lived and does not feed.


5. Cleridae

- Cleridae or “chequered beetles” are brightly colored pubescent insects of moderate size.
- Antennae is 11-segmented with the apical segment enlarged to form a distinct club.
- Elytra generally completely cover the abdomen
- There are 5 or 6 visible abdominal sternites.
- Tarsi are all 5-segmented w/ the 1st & 4th often small and indistinct
- Mostly predaceous feeding on wood borers

Red-legged Ham Beetle

(Copra Beetle)

Necrobia rufipes




- Beetle 4-5 mm long, metallic blue or bluish-green, red basal segment of antennae & legs
- Common insect pest of dried copra, dried fish & fish meal. Also attacks animal products such as ham & bacon, oil seeds & spices
- Larvae & adults are attracted to the steric & palmitic acid of rotting moldy copra
- Eggs are laid in batches of up to 30 in cracks & crevices
- With 3-4 larval instars, which are completed in ~17 days. Larvae are cannibalistic & predatory, attacking the eggs and larvae
- Female lives for up to 6 months & may produce as many 3000 eggs. Adults are active fliers
- Life cycle is completed in 42 days at 30°C and 65-70%RH.

6. Cucujidae

- Small beetles (1.5 to 2.5), flattened & reddish-brown
- Filiform antennae are usually long, up to half the body length to nearly as long as the body
- Prothorax possesses an entire, sub-lateral ridge (carina) on both sides w/c are more or less parallel
- In females, tarsi are 5-segmented while in males the front & mid-tarsi are 4-segmented
- Most species have been found under bark of trees, in damaged seed or in tunnels made by other beetles
- Some species are often predatory but *Cryptolestes* is of importance in stored products. It is both a predator and grain feeder
- The male and female closely resemble each other in appearance
- Determination of species is based on characteristics of the genitalia & antennae and the ratio of body length of male to that of female.

Rust Grain Beetle

Cryptolestes ferrugineus



- Smallest pest of stored grain, about 1/16 inch long
- Reddish-brown beetle with flattened body characterized by the presence of a very long antenna.
- Head & prothorax are very conspicuous & account for half of body length
- Cosmopolitan & is one of the most common insect pest of stored grains
- Adults are unable to survive in sound and uninjured grains.
- Follow-up the attack of the more vigorous grain insect pests

Flat Grain Beetle

Cryptolestes pusillus (Schoner)

Cosmopolitan; attack same range of commodities infested by *C. ferrugineus*

Unable to survive in sound, uninjured grains; also a scavenger, larva feeds on the remains of dead insects; often associated with grain & meat in poor conditions.

External appearance, life cycle & behavior is very similar to *C. ferrugineus* but not tolerant to low temperature and below 60% RH.

7. Curculionidae

- The species under this family is known as true weevils or snout beetles.
- Form the largest single family in Order Coleoptera that attack many growing crops of many kinds throughout the world.
- Approximately 30 species have been recorded in stored food products.
- Includes Genus *Sitophilus*, which contains 3 of the world's most destructive pests of stored grain.
- Adults may be distinguished from other stored product species by having the head protruded in front of the eyes to form a well defined snout (rostrum), antennae are elbowed and one-clubbed.
- All tarsi are 4-segmented.
- Larvae are legless in all species, stout, and slightly curved and are creamy white with a pale brown or yellowish head.

***Sitophilus* spp.- Rice weevil complex
(*Sitophilus zeamais* & *S. oryzae*)**



Sitophilus granarius

- Primary pest of all cereal grains but mainly found in corn, wheat, rice & sorghum
- Body color reddish-brown to black; elbow-like antennae

Pits on pronotum

Elytra

Flight

Distribution

Round

w/ 4 yellow spots

Good flyer

Tropical

Oblong

w/o yellow spots

Rarely fly

Temperate

Sitophilus spp.



Sitophilus granarius



- Primary pest of all cereal grains but mainly found in corn, wheat, rice & sorghum
- Head is prolonged into a snout; two light spots on each front wings
- Body color reddish-brown to black
- Small, round, closely compacted pits on pronotum
- Hind wings function as flight wings; infest corn in the field
- Larvae are white, wrinkled & legless; require 4 molts and growth period before transforming to adults
- Lives on an average of 4-5 months & each female lay 300 – 400 eggs
- Common in semi-tropical countries
- Can be identified by their genitalia

8. Dermestidae

- Adults are small to moderately large beetles (1.5 to 12 mm long), densely covered w/ hairs or colored scales
- Head is small & somewhat deflexed.
- Antennae are 5 to 11-segmented, w/ a fairly distinct 3-segmented club.
- Elytra completely cover the abdomen.
- All tarsi are 5-segmented.
- Larvae are characteristically very hairy. Their setae, which are often of taxonomic importance, may be detached from the shed larval skin.
- Live larvae can cause contamination or pose a health hazard when inhaled
- About 55 species have been reported as injurious to stored products.
- The most imp't species belong to the *Trogoderma* and *Dermestes*

Khapra Beetle

Trogoderma granarium



- Occurs in hot and dry conditions; in areas w/ a mean $T^{\circ} >20^{\circ}\text{C}$ & $<50\%$ RH; larva is highly resistant to starvation & may live for months or even years without food.
- One of the most damaging pests of stored products of agricultural origin throughout the world; most important quarantine pest.
- Wheat, barley, rice & spices seem to be the most attractive commodities but also feeds on almonds, walnuts, spaghetti & egg noodles, tapioca, dried beans, powdered skim milk, oilseed cake, corn meal, fishmeal, coconut and to a lesser extent, dried fruits.

Khapra Beetle

Trogoderma granarium



- Originated in India & had established in some Asian, Middle-East & African countries, being distributed mainly by shipping & international trade through the agency of man.
- Adults are dark brown mottled with black; body clothed with fine yellowish hairs; oblong oval approx. 1.75 – 3.5 mm long.
- Antennal segment is usually 11 but there can be as few as 9; antennal club consists at most of 4-5 segments for males, at least 3 segments for females.
- Life cycle varies from 4-6 months to several years depending upon the temperature and food supply

Hide Beetle, *Dermestes maculatus*

- The only beetle with enzymes necessary for, *Dermestes maculatus* breaking down keratin
- 5.5-10 mm long; head, pronotum and elytra dark gray to black
- Pronotum & elytra w/ uniformly-colored whitish or pale yellowish hairs
- Scutellum yellowish-brown to reddish-brown
- Antennae clubbed, basal segments reddish-orange, last 3 segments forming club
- Abdomen w/ dense covering of grayish-white hairs except for black spot in anterior-lateral corner of each abdominal segment, and an irregular-margined black band at base of abdomen
- Adults feed on dried or drying fish & lay up to 800 eggs. Rate of egg-laying is greatly increased if water is available for the female to drink.



9. LOPHOCATERIDAE

- Formerly under the Family Trogosidae.
- Only one species, *Lophocateres pusillus*, is known to occur in stored products.
- *Lophocateres pusillus* occurs widely in tropical & subtropical regions & has been recorded on many stored products such as cereals (particularly rice), pulses, cassava and groundnuts.
- Usually only a minor pest & is often found in association w/ the major primary pests.
- Common in South and Southeast Asia & its pest status may be higher in this situation.

Siamese Grain Beetle

Lophacateres pusillus King

- Occurs widely in tropical and subtropical regions.
- Found on many stored products such as cereals, legumes, cassava, & groundnuts; a minor pest occurring in large number on both paddy & milled rice in association with *S. oryzae* and *R. dominica* in South & Southeast Asia.
- Eggs are deposited in crevices; as many as 3-14 eggs may be laid per batch; at 30°C & 75%RH, egg hatches in about 7 days.
- Larval and pupal development are completed within 42 days.
- Found to develop adequately on a diet of frass from *S. oryzae*, but development was better on a medium of whole-wheat flour plus yeast.

10. Nitidulidae

- Large family of more than 2,000 species; extremely variable in form, structure & habits. Many inhabit flowers.
- Others are found in decaying animal & vegetable remains, in fungi, & in exuded sap.
- Some larvae are predatory
- Those important to stored products are attracted to moldy grains, cured meats, & dried or decaying fruits.
- Beetles are oblong with 11-segmented antennae terminating apically in a distinct 3-segmented club.
- Elytra are short: 2 or 3 abdominal sternites are always visible.
- All tarsi are 5-segmented, w/ the 4th segment shorter than the rest, while the 1st 3 segments are dilated & hairy beneath.

Corn Sap Beetle

Carpophilus dimidiatus



- Widespread in the tropical, subtropical & warm areas.
- Found in cereals and cereal products, peanuts copra, cotton seed & maize in cobs both in the field & storage; often inhabits rice mills breeding in accumulations of broken rice.
- Not attracted to clean, dry grains in good condition; presence of the insect is an indication of inadequate drying or of damp, even moldy conditions in storage

Corn Sap Beetle, *Carpophilus dimidiatus*



- 2-4 mm long, oblong, shining beetle w/ dark red to nearly black body.
- Elytra truncate, each usually bearing a large central dull orange-yellow spot with indefinite margins.
- Body cuticle is dull & moderately clothed w/ conspicuous pale yellow hairs.
- 11-segmented antennae w/ a compact 3-segmented oval or round club
- In most species, the elytra do not cover the apical two or three segments of the abdomen.
- Eggs are laid on or in decaying plant materials like moldy grain.
- Larva passes through 4 instars in the soft & moldy parts of a fruit
- Pupa is generally spent in the soil
- Female lays about 1,000 eggs over 3-4 months; adults fly readily, congregating at suitable breeding and oviposition sites.

11. Scolytidae

- Beetles of this family are mainly wood & bark borers. Only a few species are associated w/ stored products.
- Closely related to Fam. Curculionidae, & some taxonomist classify them as a subfamily of the latter.
- Many species are superficially similar to Bostrichidae but they are readily distinguished by antennal features: in Scolytidae, the first abdominal segment is about a third of the length of the antennae, w/c has a compact 3-segmented club; while in Curculionidae, antennae are elbowed & one-segmented club.
- *Hypothenemus hampei* is a serious pest of coffee berries and beans

Coffee Berry Borer (CBB)

Hypothenemus hampei (Ferr.)



- Endemic in Central Africa; introduced in the 1920s in Malaysia, Ceylon, India, & in the Phil (1960).
- CBB attacks coffee berries in the field; cause severe damage to beans if not treated before storage.
- An initial infestation of 20% of newly harvested berries could increase to 100% after 6 weeks of storage.
- Larva & adult CBB directly damage coffee by feeding on the tissues of the beans.
- Black beetle, subcylindrical and about 1.3 - 1.5 mm long.
- Mean life cycle of female is 86.5 d & longevity is 41.60 d while mean life cycle for male is 66.2 d and longevity is 25.83 d.
- Average longest female longevity is 156 d.

12. Silvanidae

- Small family closely related to the Cucujidae
- Beetles are generally parallel-sided, distinctly flattened, & rather short (2-4 mm).
- Most species have projections on the prothorax in the form of swellings or teeth in the anterior angle or several teeth along the lateral margin as in *Oryzaephilus*.
- *Tarsi are all 5-segmented.*
- Most species are probably predaceous, but two species of *Oryzaephilus* are of great importance as secondary pest attacking broken or milled cereals and oilseeds.
- Other species such as *Ahasverus advena* & *Cathartus quadricollis* are found in moldy foods (cocoa, grains and many others). is a serious pest of coffee berries and beans

SAW-TOOTHED BEETLE

Oryzaephilus surinamensis (L.)




- Slender, dark-brown beetle, about 0.5-3 mm long.
- Cosmopolitan secondary pest of cereals, mainly milled products like flour, breakfast foods, stock & poultry feeds, copra, desiccated coconut confectionary, and dried fruits.
- Larva enters damaged & broken kernel to feed especially on the germ.
- Adults & larvae are able to enter small cracks, so they can often attack packaged or nuts in shell.


SAW-TOOTHED BEETLE, *Oryzaephilus surinamensis*

- Prothorax has 6 distinctive tooth-like projections along each side
- Antennae are relatively short
- Length of temple (side of head behind the eye) is much shorter in *O. mercator* than in *O. surinamensis*
- Female lays some 375 eggs (at 30°C over a life span of 6-8 months; eggs are laid loosely among the substrate or tucked into crevices in the grain at the rate of 6-10 per day
- The pale yellow, elongate, somewhat flattened larva passes through two to four instars depending on feeding condition & freedom of movement.; full grown larva is about 4-5 mm long Full grown larva pupates within a cocoon-like structure of small grains or food particles
- Development is completed in 22 days at 30 °C & 68 days at 20 °C (70% RH).

O. mercator *O. surinamensis*



Relative lengths of eyes & temple



13. Tenebionidae

- Large family (>10,000) & ~100 species are associated w/ stored products.
- Several species are recognized as secondary pests.
- Adults vary in size from 3-10 mm long & more or less parallel-sided & nearly always uniformly black or brown
- Antennae of most Tenebrionids have 11 segments, of moderate length, generally thickened or clubbed towards the apices
- In most species, each compound eye is partly divided horizontally by a backward projection of the side of the head.
- Tarsi of hind leg are 4-segmented; front & middle legs = 5-segmented.
- Larvae are cylindrical, yellow to brown, generally well sclerotized, often w/ a distinctly banded appearance & w/ one or two pointed projections (urogomphi) at the end of the body.
- Many species =carnivorous, cannibalism is common, some are predators.
- Most important genera are *Tribolium*, & *Latheticus*; less imp: *Alphitobius*, *Gnatocerus* and *Palorus*.


Red Four Beetle, *Tribolium castaneum*

- Cosmopolitan but is more of a pest in tropical & warm temperate regions.
- Major pest of cereals, cereal products, legumes, oilseeds, nuts, spices & animal products.
- Benzoquinones secreted by the adults produce an unpleasant odor under high population.
- Reddish-brown beetle; 2.3-4.4 mm long.
- Appearance is similar to *T. confusum*, but antennae has distinct 3-segmented club.



Confused Flour Beetle

Tribolium confusum



T. confusum



T. castaneum




- Abundant in temperate regions & is less frequently found in the tropics
- Antennae have no distinct club, that expands gradually

Long-headed Flour Beetle


Latheticus oryzae

- Cosmopolitan secondary pest of some importance in the tropics & subtropics and especially common in South & Southeast Asia.
- Antennal club has 5 segments w/ terminal segment distinctly smaller than the others.
- Causes damage on whole grains & cereal products & causes same type of damage as that of *Tribolium* spp.



Depressed Flour Beetle


Palorus subdepressus



- Adult beetle is reddish-brown; 2.7 – 3 mm long.
- Head with sides flexed upwards & a distinct ridge above the eye covering the anterior portion (a).
- Eyes are large with vertical diameter exceeding the horizontal diameter and completely undivided
- Often found in most tropical and subtropical regions.
- Minor secondary pest of cereal & cereal products especially the residues of damp & moldy grain; often found in association w/ *Sitophilus* spp.

Small-eyed Flour Beetle

Palorus ratzeburgii



- Beetle is reddish-brown, flattened and somewhat oblong; 2- 3 mm long.
- Head with sides not strongly flexed upwards while the ridge above the eye is feeble and indistinct.
- Eyes are small and round
- Cosmopolitan but generally most common and more widely distributed than *P. subdepressus*.
- Appears to be of relatively minor economic importance.

14. Trogositidae

- **Members vary considerably in size, form and habits**
- **Adults are obovate (egg-shaped) to parallel without hairs or scales or at most with a very short, microscopic hairs**
- **Antennae are 11-segmented that terminate in a 3-segmented club**
- **Elytra completely cover the abdomen**
- **Tarsi are 5-segmented but the basal segment is sometimes so small that they appear 4-segmented**
- **Most tropical spp. are often associated w/ decaying wood preying on larvae of other insect pests. Some spp. are mycophagous (fungivorous).**

Cadelle Beetle

Tenebroides mauritanicus



- **Cosmopolitan; infests various commodities especially cereals & cereal products**
- **Excessive feeding by adults & larval feeding lowers germination**
- **Large (7-11 mm long), shining dark brown-black & dorsoventrally flattened**
- **Prothorax w/ base distinctly separated from abdomen by a narrow neck & w/ distinctive forwardly projecting front angles; elytra possess longitudinal ridges**
- **White eggs are laid in clusters in food material; hatch from 7-10 d**
- **Larva completes its growth from 2-14 mos then pupates in secluded place**
- **Female lays about 1,000 eggs under favorable conditions**
- **Both larvae & adults can live for considerable period w/o food**

Stored Product Lepidoptera

Moths, skippers and butterflies; few species of moths are known as storage pests.

Families Gelechiidae & Pyralidae: most important


Family Gelechiidae

Have long recurved labial pulps, a proboscis that is densely covered with scales and pointed wing tips.

***Sitotroga cereallela* and *Pthorimaea operculella* are important in Southeast Asia.**

Angoumois Grain Moth

Sitotroga cerealella (O.)



- Important primary pest of cereal grains in the tropics & in warm temperate regions
- Moth is a good flier attacking standing crop in the field & infestation is carried over to storage
- Moth is yellow-brown w/ a wing span of 10-18 mm
- Forewings have a small black spot in the distal half; hind wings have long fringe of hairs, longer than half of the wings & are sharply pointed at the tip
- When moth is at rest, wings are folded in sloping manner over the abdomen, antennae point backward & are slightly raised above the wings, while the long, slender & pointed labial palpi are raised upwards like two horns.



Stored Product Lepidoptera..... continuation

Family Pyralidae

- All have straight or recurved labial palps & rounded wing *tips*
- Include *Corcyra cephalonica*, *Doloesa viridis*, *Plodia interpunctella*, *Ephesia cautella* and *E. elutella*

Rice Moth

Corcyra cephalonica



- Found throughout the humid especially in South & SE Asia; widespread & common in Africa.
- Common pest of rice, corn, sorghum, millet & other cereals; other commodities such as cocoa, copra, legumes, dried fruits, nuts & spices are known to support infestation.
- Adult moth is nocturnal and short-lived.
- Webbing formed by larvae are tough & closely woven, later tightly matted together w/ the food particles, frass & cast skins; this white aggregation characterizes the infestation of this moth & most distinctive when attach to bag surfaces; the last instar larva pupates w/in this aggregation.

Rice Moth

Corcyra cephalonica (Stain.)



- Medium-sized w/ a wing span of 15-20 mm.
- Forewings are uniformly greyish-brown w/ thin vague lines of darker brown along the veins while the hind wings are pale buff; fringes along the wing margins are relatively short.
- Female labial palps long & pointed while short, blunt & inconspicuous in male.
- Larvae are generally creamish-white except the head capsule & prothoracic tergite w/c is brown;
- There are well-developed prolegs on abdominal segments 3-6 and 10.
- With a conspicuous seta above each spiracle; on the 8th abdo.segment, this seta arises from a clear patch of cuticle surrounded by a ring.
- Female lays an avg. of 154 eggs w/ a max. of 268 eggs; hatch in 3-9 d.
- w/ 7-8 larval instars; larval period is 29-50 d in female & 28-46 d in male.
- Pupal period is 6-11 days for both sexes.

Cacao Moth

Ephestia cautella



- Found throughout the tropics & subtropics but is rather less common in arid areas.
- Serious pest of wide range of commodities especially cereal flours & cereal products, nuts, cocoa, dried fruit, spices, etc; in the Phil, it is a common pest of cocoa beans, frequently found in milled rice and corn.
- Adult forewings are greyish-brown w/ indistinct pattern; wing span is 11-20 mm & both fore & hind wings have broadly rounded tips & only short fringes of hairs.
- Labial palps curve upward in front of the head, rather blunt at the tip

Cacao Moth

Ephestia cautella



- Female lays up to 300 eggs on stored commodity for the first 3-4 days & few in the remaining 4-5 days of life; hatch in ~3 d at 30°C.
- Normally undergoes 5 larval instars in about 22 days under optimum conditions of 32.5°C & 79% RH on the embryo of softer seeds.
- Larva spins silken threads while eating w/c pull the other grains and frass together (webbing) the characteristic infestation of storage moth.
- Before pupation, last instar larva spins a cocoon w/c is both thinner & looser than *C. cephalonica*; pupal period ~7-11 d.
- Under optimum conditions, devt from egg to adult ~29-31 d.

Indian Meal Moth

Plodia interpunctella

(Hubner)




- Cosmopolitan in warm climates but can survive in heated buildings in cool temperate countries.
- Although recorded in the Phil. In 1950s, the pest was rarely observed since the late 1970s.
- Serious pest of stored grains, legumes, oil & vegetable seeds and of processed and milled products, nuts, dried fruits and chocolates.

Indian Meal Moth

Plodia interpunctella

(Hubner)



- Moth wing span of 20 mm; forewing is cream colored in the basal 2/5; while the rest is red-brown w/ some grey markings.
- The slightly separated labial palps point directly forward like an open beak.
- Larva may be distinguished by the absence of dark spots at the bases of the setae together w/ the evenly thickened and poorly sclerotized rims in the spiracles.
- Female lays up to 400 eggs (average = 212 days).
- At 30°C & 70% RH, larva undergoes 4-7 molts in ~16 d.
- Pupal period is about 7 d.
- Total development is about 27 d.

Potato Tuber Moth, *Pthorimaea operculella*(Zeller)

- Cosmopolitan but apparently absent in West Africa; found in Africa, Asia, Australia, Central America & West Indies.
- First recorded in the Philippines in 1966 but could have existed earlier; could have been introduced thru importation of seed tubers prior to World War II from China, Germany, Japan & United States.
- Both a field & storage pest of potato; although it infests growing crops of many species of the family Solanaceae, but it is only a storage problem in potato tubers.
- The 7-9 mm long moth is generally grayish brown but mottled with dark brown.

Potato Tuber Moth, *Pthorimaea operculella* (Zeller)

- Hind wings are fringed w/ long hairs & wing span is about 12 mm long; w/ black dots on the forewings.
- Female lays ~100-300 eggs w/c hatch in 5-9 days at 20-25°C.
- Newly hatched larva is creamy & becomes brownish or yellow pale pink after feeding; undergoes 4 instars in 11-30 days in potato tubers.
- Under field conditions, pupal period is 5-12 d; on the growing plant larva passes the pupal stage in curl or rolled leaves.
- Life cycle is from 20-65 days in the laboratory
- Female life cycle ~41-68 days & the male from 37-53 days

Green Moth

Doloessa viridis (Zell.)



- Found in rice, maize, peanut & copra; reported as a potential pest of stored grains, shelled peanuts & dried fruits in Indonesia; & in copra in Malaysia & rice & corn in the Phil.
- In early 1960s, this pest was used for rearing of *Trichogramma* parasitoids in the Philippines;
- became scarce in 1970 & was replaced by *C. cephalonica*.
- Moth wing span of 13-25 mm; easily recognized by its emerald green forewings w/ 4 irregular black patches forming a discontinuous oblique bands across the middle to posterior part of wings; another set of 5 smaller irregular patches form an oblique band across the posterior area of the wings; & 10 irregular black spots form a discontinuous narrow band in the wings.

Green Moth

Doloessa viridis (Zell.)



- Female laid an avg. of 87 eggs w/c hatch in 6-7 days.
- Passed through six larval instars within 26 to 36 days; feeds on the embryo or softer parts of the seed until it matures, & silk thread spin (webbings).
- Infestation is characterized by the presence of clumps formed by the webbings to w/c the grains & frass adhere, that may ultimately contain the cocoon.
- Larva forms a pupation cell by spreading a mat of web to cover the whole body length; pupal period varies from 7-11 d.
- Adult longevity w/o food is 7 days; total life cycle in rice ranges from 43-54 d (avg = 47 d)



ASEAN Regional Training Workshop on Diagnostics of Weevils of Quarantine Importance

July 10-22, 2017

**Institute of Weed Science, Entomology and Plant Pathology
College of Agriculture and Food Science
University of the Philippines Los Baños**

SESSION 7

DISSECTION OF GENITALIA


Lecture Notes by:

DR. HIRAKU YOSHITAKE
Institute for Agro-Environmental Sciences
NARO, Tsukuba, Japan

DISSECTION OF GENITALIA

Procedure

- 1) To examine male and female genitalia, specimens are macerated in hot water (Fig. 1) for 10 minutes to 24 hours, depending on the condition of each specimen.
- 2) Each specimen is dissected under the stereoscopic microscope, using a pin and tweezers (Fig. 2).
- 3) The abdomen (or a part of abdomen) is removed from the body (Fig. 3).
- 4) The abdomen is cleaned in hot 10% KOH solution (Figs. 4, 5) for 2 to 15 minutes, depending on the condition of each specimen.
- 5) The male and female genitalia extracted from the abdominal segments are mounted on hole slides with glycerol (male; Fig. 6) or water (female; Fig. 7), and examined with stereoscopic or optical microscope, depending on the size of each specimen.
- 6) After examination, the male and female genitalia (as well as abdomen) are preserved in microtubes with glycerol (Fig. 8) or glued to cards (Fig. 9) and pinned with the body of each specimen (Figs. 8, 9).





ASEAN Regional Training Workshop on Diagnostics of Weevils of Quarantine Importance

July 10-22, 2017

**Institute of Weed Science, Entomology and Plant Pathology
College of Agriculture and Food Science
University of the Philippines Los Baños**

SESSION 8

PRESERVATION TECHNIQUES

Lecture Notes by:

DR. SHERYL A. YAP

Institute of Weed Science, Entomology and Plant Pathology College of Agriculture
and Food Science
University of the Philippines Los Baños



COLLECTION AND PRESERVATION METHOD

SHERYL A. YAP *Ph.D.*

Institute of Weed Science, Entomology and Plant Pathology
College of Agriculture and Food Science, UP Los Baños Los
Baños, Laguna, Philippines sayap3@up.edu.ph

TRAINING WORKSHOP ON DIAGNOSTICS OF WEEVILS OF QUARANTINE IMPORTANCE (Proj. No. AGF/CRO/11/007/REG)

10-22 July 2017, IWEP, CAFS, UPLB, PHILIPPINES

Funded by: Japan - ASEAN Integration Fund (JAIF)

Through: ASEAN Plant Health Cooperation Network of the ASEANET (APHCN-ASEANET)

Where Weevils can be found?



Locating insects:

Weevils can be found almost anywhere in the world.

Weevils are commonly found in:

- ***flower and vegetable garden***
- ***grasses in lawns and fields weeds***
- ***bushes***
- ***fruit and shade trees***
- ***storage facilities***
- ***homes***




COLEOPTERA

beetles

weevils



Pinterest



COLEOPTERA

- forewings: modified into hard plate-like coverings
- hindwings: membranous
- chewing mouthparts

Habitat: All habitats

Equipment to use: Sweep nets, light, traps

Collection method: Beating, sweep
vegetation, examine
trap samples, hand pick

Preparation: Pin adults, larvae 80-85% EtOH



COLLECTION OF SPECIMENS

The slide features a white background with several overlapping, semi-transparent green geometric shapes on the right side. These shapes include a large dark green triangle, a medium green triangle, and a light green triangle, all pointing towards the center of the slide. The text 'COLLECTION OF SPECIMENS' is centered horizontally and positioned in the lower half of the slide.

Beating



MAKE GIFS AT GIFSOUP.COM

Sweeping



Traps



Sex pheromone trap



Pecan weevil trap



PRESERVATION AND STORAGE OF SPECIMENS

PRESERVATION OF SPECIMENS

TEMPORARY STORAGE

- **Refrigeration**

- a. medium-sized to large specimens may be left in tightly closed bottles for several days in a refrigerator and still remain in good condition for pinning
- b. After refrigeration, moisture must be removed by drying or by placing an absorbent paper

TEMPORARY STORAGE

- **Dry preservation**
 - a. Place in small boxes, paper tubes, triangles, or envelopes for an indefinite period
 - b. To allow specimens to become dry
 - c. Under the sun or oven

STORAGE OF SPECIMENS

TEMPORARY STORAGE






- **Dry preservation**

- Small boxes, paper tubes or triangles, or envelopes
 - ▶ **NO** - Soft-bodied insects
 - ▶ **NO** - Tightly closed, impervious containers of metal, glass or plastic
 - ▶ **NO** - specimen collected at different times or places in the same container
 - ▶ **Can be layered in the same container** - each **layer** with data

TEMPORARY STORAGE

Papering

- ← Do not pack specimens together
- ← No - airtight containers or envelopes




MOUNTING/PERMANENT PRESERVATION

- a. so that they may be handled and examined with the greatest convenience
- b. with the least possible damage

“ The value of specimens may depend on how well they are preserved.”

MOUNTING/PERMANENT PRESERVATION

- Dry specimens
 - must first be relaxed so that body and appendages will not break
 - it can also be rearranged and repositioned







MOUNTING/PERMANENT PRESERVATION

- **Liquid-preserved specimens**
 - a. for soft-bodied specimens**
 - b. it can also be rearranged and repositioned**
 - c. stored in small containers or vials with ethyl alcohol or other preservatives**
- **Slide mounting**
 - a. for small and minute specimens such as mites**



PINNING (INSECTS)

- a. insect pins are available in different sizes
- b. made of stainless steel






Pinning Blocks

Rectangular Block



Stair-step Block







Large Coleoptera

Right forewing (wing cover or elytron) near the base.

Do not spread the wings of beetles.

Which is correctly pinned?




- A. correctly pinned specimen
- B. specimen mounted too low
- C. specimen not level

STORAGE OF SPECIMENS

Double mounts

- ▶ for insects that are too small to be pinned directly on standard pins - minuten pin or card points (acid-free)
- ▶ Adhesive - white glue or carpenter's glue - soluble in a solvent
- ▶ Right side of the specimen, with the left side and midventral area clear
- ▶ Card point - side of the thorax not on the wings



Double mount using card point method.

LABELING

- it is important so that specimens will have a scientific value
 - a. paper must be 100% rag content - not acidic
 - b. pens must have a permanent ink and will not “run” or blot
 - c. labels must be lettered carefully - readable
 - d. size must be relative to the specimens - not too small, not too large

Label data includes:

- Locality
- Date of collection
- Name of collector
- Other useful data such as host plants, etc

Philippines: Luzon Is. Brgy.
Pinaglubayan, Polillo, Quezon
March 12, 2001
SA Yap

LABELING

- ← Collection data should accompany specimens at all times during preparation
 - ← The label or the information is as important as the specimen
- No label, no SCIENTIFIC VALUE**

Paper

← An appropriate is important

- ▶ Acid free
- ▶ Made of cotton fibers
- ▶ 100% rag content
- ▶ Should be heavy
- ▶ Surface of the paper should be smooth

Lined ledger paper = 100% rag & 36 -pound weight

2-ply Bristol board

Herbarium sheets

Pens

- ← “rapidograph” or technical drawing pens
- ← 0.25 mm (no. 000) to 0.30 mm (no.00)



Ink

- ← India ink
- ← Permanent
- ← Will not “run” - placed in jars or vials with liquid preservatives
- ← Soft lead pencil

Lettered and Printed Labels

- ← Labels may be lettered carefully by
 - hand with a fine-pointed pen
 - printers - Arial, size 4 to 6

Size of Labels

- ↯ The maximum size of a label should be about **7 x 18 mm**
- ↯ **5 lines or about 13 capital letters**
- Use multiple labels - to accommodate **all** of the information that must be presented

Label Data

- accurate, concise and unambiguous

← Locality

- ▶ it can be found in any good map
- ▶ Coordinates of latitude and longitude may be given
- ▶ Countries: capital letters
- ▶ U.S.: capital letters = States

e.g. PHILIPPINES: Laguna, Los Baños,
64km, 14.1667° N, 121.2167° E

Label Data

← Date

- ▶ Avoid ambiguity
- ▶ Cite the day, month and year
- ▶ Day and year = Arabic numerals
- ▶ Month = Roman numerals
- ▶ Hyphens or periods can be place in between

e.g. 01-XII-2012

01.XII.2012

Label Data

← Collector

- ▶ Very useful
- ▶ Last name = spelled out
- ▶ Given and middle name/s = initials
- ▶ Many collectors = Leader's name followed by et al.

e.g. S.A. Yap


S.A. Yap et al.

Label Data


← Other data

- ▶ Very useful or relevant to the specimens or collector
- ▶ E.g.: hosts of parasites or host plants = as much as possible do not give vernacular or common names, at least give the genus or family name (Host: *Homo sapiens sapiens*)
- ▶ E.g.: details of the habitat (elevation, ecological types and conditions of collection)
(1500masl, collected by Malaise trap)

Mounted Insect and Label Placement



A small insect mounted on a point



A properly mounted beetle

Placing the Labels

- use pinning block - desired height

← Direct pinning:


- ▶ Centered under the specimen
- ▶ Long axis (length) = long axis of the specimen
- ▶ Should be aligned transversely at right angles to the axis of the body with the upper margin toward the head

Placing the Labels

- use pinning block - desired height

Double-mounted insects:

pin = center of the right side of the label



Placing the Labels

← Vials or jars:

- ← Large labels that includes all collection data
- ← Do not fold the label - minimize damage and loss of small specimens
- ← Place labels inside the vials or jars
- ← Pencil or ink can be use

Placing the Labels

← Microscope slides:

- ← With pressure-sensitive adhesive
- ← Left side = taxonomic information
- ← Right side = collection information
- ← Mounting medium must be included



Placing the Labels

← Identification labels:

Used when specimens are sent to an expert for identification:

- All collection data or information must accompany the specimens including field notes if available



ASEAN Regional Training Workshop on Diagnostics of Weevils of Quarantine Importance

July 10-22, 2017

**Institute of Weed Science, Entomology and Plant Pathology
College of Agriculture and Food Science
University of the Philippines Los Baños**

SESSION 9

DNA SEQUENCING

Lecture Notes by:

DR. BARBARA L. CAOILI

**Institute of Weed Science, Entomology and Plant Pathology College of Agriculture
and Food Science
University of the Philippines Los Baños**

DNA Barcodes for Insect Identification

Barcode of Life

Identifying Species with DNA Barcoding

[About](#) [Community](#) [Resources](#) [Events](#) [Partners](#) [News](#)



Barbara L. Caoili, Ph.D.

Institute of Weed Science, Entomology and Plant Pathology
College of Agriculture and Food Science
University of the Philippines Los Baños



**ASEAN Regional Training Workshop on
Diagnostics of Weevil of Quarantine Importance**


July 10 to 22, 2017
IWEP, CAFS, UPLB

Lecture Outline

- I. What is DNA?
- II. What is DNA barcode?
- III. Specimen Collection
- IV. DNA Extraction Methods
- V. Polymerase Chain Reaction
- VI. Agarose Gel Electrophoresis
- VII. DNA Sequencing
- VIII. Data Analyses




What is DNA?

- deoxyribonucleic acid
- genetic material



What is DNA?

DNA Structure



What is a DNA Barcode?

A unique pattern of DNA sequence that can identify a species like how the unique patterns in universal product codes identify products



What is DNA Barcode?

- Standardized short sequence of DNA between 400 to 800 bp long
- Can be easily isolated and characterized for all species on the planet
- Used to identify species through reference to DNA sequence libraries or databases

The Uses of DNA Barcodes

As a research tool to:

1. assist taxonomist in species identification
2. identify regulated species (invasive and endangered)
3. survey of arthropods for microbial pathogens
4. test the identity and purity of biological products
5. new and cryptic species discovery
6. address fundamental ecological and evolutionary questions

DNA Barcoding Methods in Brief

Basic Steps:



1. Building the barcode library of known species
2. Matching, or assigning the barcode sequence of unknown sample against the barcode library for identification


What it takes to be a DNA Barcode

A DNA barcode should:

1. contain significant species-level genetic variability and divergence
2. process conserved flanking sites for developing universal PCR primers for the widest taxonomic application
3. be of appropriate sequence length so as to facilitate current capabilities of DNA extraction and sequencing

Specimen Collection

For beginners:



Specimen Collection

Specimen collection information

Specimen/ Field ID	Collectors	Collection Date	Location (GPS)	Notes	Storage Condition
IPL-01	Gelyn D. Sapin	July 17, 2017	47.9053, -124.6261	Light trap	Collected into an individual tube of 99.9% EtOH which was transferred to the -20C freezer on return to lab

DNA Extraction Methods

Basic Steps:

1. Cell lysis
2. Lipid removal
3. Protein removal
4. DNA precipitation



DNA Extraction Methods

Museum Sample: Modified QIAGEN Animal Tissues Spin Column Protocol (Sutrisno, 2012 modified from DNeasy Blood and Tissue Handbook)

Remove abdomen from the specimen



Transfer into sterile microcentrifuge tube



Add 1% PK Buffer (20 ul of Proteinase K (20 mg/ml) in 180 ul ATL Buffer)



Incubate at 55°C for 2 to 4 hours

Add 0.1 ml 1% PK Buffer



Incubate overnight at 55°C

Remove abdomen and process following the manufacturer's protocol

DNA Extraction Methods

Fresh or Pickled Sample: Modified CTAB Method (Latina et al., 2017 modified from Clark et al., 2001)

Grind beetles in 500 μ l CTAB buffer + 5 μ l Proteinase K (20 mg/ml)

↓ vortex, 1 minute

Incubate samples at 65°C for 1 h. vortex every 20 mins

↓ cool down sample at room temperature

Add 10 μ l RNase A (8 mg/ml); Incubate at 37°C for 2 h; vortex every 30 mins

🌀 10,000 x g, 10 mins, room temperature

Transfer supernatant into a new microcentrifuge tube with 500 μ l chloroform: isoamyl alcohol (24: 1), vortex

🌀 10,000 x g, 15 mins, room temperature

Transfer supernatant into a new microcentrifuge tube with 500 μ l 100% isopropanol; mix by gentle inversion (30 x)

DNA Extraction Methods

Fresh or Pickled Sample: Modified CTAB Method
(Latina et al., 2017 modified from Clark et al., 2001)

Continuation ...

Incubate at 4°C, 2 h

 10,000 x g, 15 mins, 4°C

Remove the supernatant & wash DNA pellet with 700 ul 70% ethanol

 10,000 x g, 15 mins, 4°C

Remove the supernatant & wash DNA pellet with 700 ul 100% ethanol

 10,000 x g, 15 mins, 4°C

Remove ethanol and air-dry, 10 mins



Resuspend DNA pellet in 30 ul TE Buffer

Polymerase Chain Reaction

- *In vitro* technique for amplification of a specific DNA sequence
- Based on the ability of DNA polymerase to synthesize a new strand of DNA complementary to the template strand
- Target DNA sequence can be amplified a billion fold in several hours
- Requires a DNA template and the free 3'-OH


Polymerase Chain Reaction

Components:

1. DNA template
2. Oligonucleotide primers (forward and reverse)
3. Deoxynucleotide triphosphates (dNTPs)
4. Buffer
5. Divalent cations (Mg_{2+} or Mn_{2+})
6. DNA polymerase: *Taq* polymerase
7. Sterile deionized water

Polymerase Chain Reaction

The three-step reaction



Step-1
Initialization

Denaturation

Step-2

Annealing

Step-3


Extension

Final Elongation


Final Hold

Agarose Gel Electrophoresis

A separation technique based on the mobility of ions in electric field




Agarose Gel Electrophoresis




DNA Sequencing

Process of determining the precise sequence of nucleotides in a DNA fragment




DNA Sequencing

Components




DNA Sequencing

Difference between ddNTP and dNTP




Dideoxynucleotide (ddNTP)



Deoxynucleotide (dNTP)

DNA Sequencing



Data Analysis

Bioinformatics Softwares

- computer programs used to process data
 - BioEdit
 - BLAST
 - Clustal X
 - Sequin

References

- Clark, T.L., Meinke, L.J. and Foster, J.E. (2001). Molecular phylogeny of *Diabrotica* beetles (Coleoptera: Chrysomelidae) inferred from analysis of combined mitochondrial and nuclear DNA sequences. *Insect Molecular Biology*, 10, 303–314.
- Hebert PDN, Cywinska A, Ball SL, deWaard JR (2003) Biological identifications through DNA barcodes. *Proc R Soc B* 270:313–321. doi: 10.1098/rspb.2002.2218
- Savolainen V, Cowan RS, Vogler AP et al (2005) Towards writing the encyclopedia of life: an introduction to DNA barcoding. *Philos Trans Ser B* 360:1850–1811. doi: 10.1098/rstb.2005.1730
- Winter WD Jr (2000) Basic techniques for observing and studying moths & butterflies (Memoir No 5). The Lepidopterist's Society, Cambridge

MODIFIED CTAB DNA EXTRACTION PROTOCOL FOR RICE and SWEET POTATO WEEVIL

1. Grind fresh live beetles (or 95% Ethanol-preserved or frozen (-80)) in 500 ul CTAB buffer (100 mM Tris-HCl, 1.4 M NaCl, 0.02 M EDTA (pH 8.0), 2.0% CTAB, and 0.2% β -mercaptoethanol). Add 5 ul of 20 mg/ml of proteinase K to each sample and vortex for 1 minute.
2. Hold the samples at 65 °C for 1 h with vortexing at 20 min intervals.
3. Cool the samples to room temperature then add 10ul of 8mg/ul RNase A. Vortex and incubate at 37 °C for 2 h. (Vortex the samples at 30 min intervals).
4. Subject the samples to centrifugation at 10 000 x g for 10 min at room temperature. Transfer the supernatant to a fresh tube, with 500 ul of a chloroform: isoamyl alcohol (24:1). Vortex the mixture and centrifuge for 15 min at 10 000 x g.
5. Transfer the upper aqueous layer to a fresh tube with 500 ul of 100% isopropanol (-20 °C). Mix the tubes through gentle inversion for 30 times then place them at 4 °C for at least 2 h followed by centrifugation at 10 000 x g at 4 °C for 15 min.
6. Remove the supernatant and wash the DNA pellet with 700 ul of 70% and 100% ethanol (-20 °C), respectively. Centrifuge the samples at 10 000 x g at 4 °C for 15 min for every wash.
7. After the final wash and ethanol removal, air-dry the DNA pellet for 10 minutes and re-suspend it in 30ul of TE buffer (10 mM Tris-HCl, 1 mM EDTA (pH 7.6)) or nuclease-free water.

Reference:

Clark, T.L., Meinke, L.J. and Foster, J.E. (2001) Molecular phylogeny of *Diabrotica* beetles (Coleoptera: Chrysomelidae) inferred from analysis of combined mitochondrial and nuclear DNA sequences. *Insect Molecular Biology*, 10, 303–314.



ASEAN Regional Training Workshop on Diagnostics of Weevils of Quarantine Importance

July 10-22, 2017

**Institute of Weed Science, Entomology and Plant Pathology
College of Agriculture and Food Science
University of the Philippines Los Baños**

SESSION 10

QUARANTINE POLICIES

(The Philippine Plant Quarantine System)

Lecture Notes by:

Mr. Joselito Antioquia

**Bureau of Plant Industry
National Plant Quarantine Service Division
Department of Agriculture**



PHILIPPINE PLANT QUARANTINE SYSTEM


Bureau of Plant Industry

National Plant Quarantine Services Division



BUREAU OF PLANT INDUSTRY (BPI)

Organizational Structure






GOVERNING LAW



Philippine Plant Quarantine is being enforced by the Bureau of Plant Industry- National Plant Quarantine Services Division by virtue of **PRESIDENTIAL DECREE 1433**, as amended otherwise known as the **Plant Quarantine Law of 1978**.

Law of 1978.





Mandate

- prevent the entry (and establishment) of foreign pests into the country
- prevent spread of pests already existing in the country and;
- comply with international standards



NATIONAL PLANT QUARANTINE SERVICES
DIVISION

Chief,
NPQSD

Biotech Secretariat

Asst Chief, NPQS

Asst
&
Admin
Section

Quarantine
Policy &
Coordination
Section

Pest Risk
Analysis
Section

Market
Access
Section

Special
Programs
Section

Laboratory
Operation
and
Diagnostic
Section

Data Base
Management
System
Section

Information
and
Communication
Section

SPS Section

Capacity
Building and
Cooperation
Section

NPQS
Stations



NATIONAL PLANT QUARANTINE SERVICES DIVISION

**Chief,
NPQSD
SG-24**

**Asst Chief, NPQS
SG-22**

Biotech
Secretariat

Accreditation & Licensing Section

- Import
- Export
- (Exporter/ Facility Operator/ Farmer)
- Treatment Providers (Vapor Heat Treatment/ Extended Hot Water Treatment /Fumigators, Heat Treatment)
- Clearances

Quarantine Policy & Coordination Section

- Supports activities of the PQ Board
- Drafts Memo/ AO & National Guidelines in compliance with approved international standards
- Coordinates with other regulatory agencies of DA which has overlapping functions w/ the NPQS
- Identifies/ formulates/ reviews quarantine policies & issuances
- Coordinates w/ the PRA group on the formulation of AOs, agreements,

Pest Risk Analysis Section

PRA conducts Risk Analysis for Imported products (pest categorization, risk assessment and risk management)

- Conducts PRA in situations where there is pest outbreak in the country of import or in the Philippines
- Revisit the existing PRA's where there is high pest detection, changed in policies/regulations and new pest information in the country of import
- Establishment/ updating pest data base
- Review of PQ import conditions
- Communicate with stakeholders & trading partners on the result of the PRA
- Visit exporting country to determine the phytosanitary measures to be implemented

Market Access Section

- Develop a work programme for the market access proposals
- Gather information & compile dossier
- Prepare market access submissions
- Consider results of risk analysis by importing country
- Recommend policy & operation of regulatory requirements
- Identifies/ request/ maintain market for agricultural products
- Drafts Multi-lateral and Bilateral Agreements/ workplan based on results of PRA
- Engages in actual negotiations with other countries
- Consultation with stakeholders

Special Programs Section

- Low Monitoring Survey
- Mango Program
- Okra Program

Laboratory Operation and Diagnostic Section

- Oversee the implementation of all laboratories of the NPQS
- Function as a center for pests identification and standardization of diagnostic procedures
- Develop new procedures for diagnosis & control of pests
- Provide accurate & timely identification of pest to be used as basis for any quarantine actions or decisions
- Collaborate w/ experts/scientist in providing technical assistance on matters related to different plant pest groups
- Evaluate the needs & prepare action plan for development/ improvement of each of the

Data Base Management System Section

- Collects, organizes, stores, modifies, updates, extract and provides information on Plant Quarantine
- Consolidation of monthly regional report & annual reports
- Establish /maintain/ update/ manage export & import data for agricultural plants & plant products

Information and Communication Section

- IEC
- Press Release
- Maintenance/ updating of the PQS website

SPS Section


- Gather/ update/ submit/ upload required info in compliance w/ the Phil. National reporting obligations to the IPPC, APPC
- review comment, act on notification related to trade
- Review/ evaluate/ comment on the draft ISPM
- Address issues/concerns on the implementation /enforcement of the food safety act
- Negotiate dispute on any matter that may arise under the SPS Agreement
- Prepares position papers/briefer for delegations, attending international forums such as SOM-AMAF, CPM, WTO, BIMF, etc.

Capacity Building and Cooperation Section

- Identify possible source of fund for capacity/capability building for all aspects of quarantine
- Coordinates w/ funding agencies which will support the capacity enhancement programs
- formulate proposals to be submitted to possible funding agencies
- determine capacity/cap. Enhancement needs of the NPSQD
- Manage the financial & administrative aspect of quarantine

NPQS Stations

- Inspects targeted cargo for pests and diseases
- Direct the treatment, re-shipment, re-exportation, or destruction of consignment when harboring pests, pathogens are found
- Encode relevant data to the DA Trade System
- Organize and conduct diagnostic tests as required.






Where We Work

3. **23** Plant Quarantine stations
4. **5** Sub-stations
5. **143** Sub-ports
6. **38** major seaports
7. **50** sub-seaports
8. **56** airports/ sub-airports

main office is located at Bureau of Plant Industry, 692 San Andres St. Malate Manila.





Manpower Complement

PERMANENT

304 incumbents

51 vacant positions (for filling up)

JOB ORDER

171 incumbents





What We do

- Promulgation and Enforcement of Plant Quarantine Rules and Regulations
- Bilateral discussions / negotiations (Market Access)
- Conduct Plant Pest Risk Analysis (PRA)
- Conduct Laboratory Analysis
- Commodity Inspection and treatment
- Post Entry Quarantine





What We do...


- Registration/Accreditation of Importers, Exporters, Farmers/Growers, Treatment Facilities and Treatment Providers
- Pest Survey
- Establishment of Pest Free Area (PFA)
- Other relevant functions to achieve its mandate (e.g. formulate rules, guidelines, orders, etc)



Commodity Inspection






Forms and Requirements SPS Import Clearance

		Republic of the Philippines Department of Agriculture BUREAU OF PLANT INDUSTRY			SPS Import Clearance	
This SPS Import Clearance is good for a single shipment only						
1. SPS Import Clearance No.		2. Place Issued	3. Date Issued		4. Must Ship Out by Date	
5. Name of Importer			6. Business Address of Importer / Handler / Company			
7. TIN		8. Contact No.				
9. Name of Manufacturer / Producer / Plant			10. Business Address of Manufacturer / Producer / Plant			
11. Establishment No. of Manufacturer / Producer / Plant						
12. Name of Exporter / Supplier			13. Address of Exporter / Supplier			
14. Country of Source		15. Country of Origin		16. Place of Origin		
17. Purpose of Importation						
18. Tariff code		23. Description / Specification / Classification	24. Quantity & Unit of Measure	25. Allowable Tolerance (% of qty)	26. Total Value (FOB US\$)	
19. Product / Commodity Name						
20. Brand Name						
21. Generic Common Name						
22. Scientific / Chemical Name						
29. Quarantine Site for live plants / animals / fish		29. Final Destination / Warehouse / Cold Storage / Plant				
30. Recommending Approval			32. Authentication			
31. Approval						

		Republic of the Philippines Department of Agriculture BUREAU OF PLANT INDUSTRY		SPS Import Clearance	
33. Import Conditions					
1. This shipment shall be accompanied by a Phytosanitary Certificate Issued by a duly authorized Plant Quarantine Officer of the country of origin and must be absolutely free from soil.					
2. Immediately upon arrival of said plant materials at the port of entry they are to be turned over to the Plant Quarantine Service for inspection.					
3. None of the above -described plant materials may be released at the port of entry unless cleared by the Plant Quarantine Officer thereof.					
4. If the plant materials are found upon inspection to be infested with pests or infected with any plant disease so that they can no be phytosanitized by any treatment available, they may be destroyed, and such destruction can not be made the basis of a claim for damages against the Bureau of Plant Industry.					
5. Corresponding fees for inspection, treatment, etc., shall be paid by the importer prior to release of said plant materials.					
34. Other Conditions / Requirements					
35. Import Clearance Fee				36. OR No.	
37. Inspection fees (indicative - actual depending on actual volume imported and inspection)					
38. Conforms / Acceptance [Signature over Printed Name of Importer]				39. Date Signed	
To be Accomplished by the DA Border Inspector at the Port of Entry					
40. Signature over/ Printed Name of Inspecting Officer				41. Date Inspected	
				42. DA Border Inspector's Report No.	
43. Inspector's Stamp				44. Fees Collected	
				45. OR No.	

Forms and Requirements

Phytosanitary Certificate from the country of origin


Name and Address of exporter:		No. 7107580			
M-SQUARE TRADING PTY LTD 3 HAWKESWORTH PLACE CHERRYBROOK NSW 2126 AUSTRALIA		 Australian Government Department of Agriculture, Fisheries and Forestry PHYTOSANITARY CERTIFICATE Plant Protection Organisation of the Australian Government			
Declared name and address of consignee:		Place of Origin	Code		
M-SQUARE MERCHANDIZING GROUP INC. 50 G DE JESUS ST BAGONG BARRIO, BARANGAY 139, CALOOCAN CITY PHILIPPINES		NEW SOUTH WALES	2000		
Declared means of conveyance:		Country of Final Destination	Code		
AIR FREIGHT/PR212	Declared point of entry:	PHILIPPINES	PH		
MANILA		TO: Plant Protection Organisation of PHILIPPINES			
Import Permit Number		ICDABPI1414034171 9-OCT-2014			
Distinguishing marks and container numbers:	Number and Description of packages:	Name of produce/quantity declared:	Total net contents KGM	Botanical Name of plants	Commodity Code
	5 CARTONS	LEEKs	50.000	<i>Allium ampeloprasum porrum Brassica oleracea var. italica</i>	07039000
	20 CARTONS	BROCCOLI	160.000	<i>Lactuca sativa Brassica oleracea var botrytis</i>	07041000
	48 CARTONS	LETTUCES	576.000	<i>Daucus carota Psalliotia campestris</i>	07051900
	5 CARTONS	CAULIFLOWERS	120.000		07041000
	30 CARTONS	CARROTS	120.000		07061000
	30 CARTONS	MUSHROOMS	120.000		07095115
ADDITIONAL DECLARATIONS					
No Additional Declaration					
			Number of Packages (Total)	Mass (Total)	
			114	1126.000	KGM
DATE	DISINFESTATION AND/OR DISINFECTION TREATMENT				
NO TREATMENTS					
This is to certify that the plants, plant products or other regulated articles described herein have been inspected and/or tested according to appropriate official procedures and are considered to be free from the quarantine pests specified by the importing contracting party and to conform with the current phytosanitary requirements of the importing contracting party, including those for regulated non-quarantine pests.					
Place of issue		Code:		Seal	
NSW		2129			
Name of Inspector	Inspection Date		Code:		
MARK WILSON	3RD DECEMBER 2014		2129		
Name of Authorised Officer		Signature of Authorised Officer an Officer of the Dept			
Bev Beacham					

H16-1106

No. 4763395		
 Department of Agriculture Ministry of Agriculture and Cooperatives, Bangkok, Thailand Phytosanitary Certificate		
Plant Protection Organization of Thailand		
TO: Plant Protection Organization (s) of PHILIPPINES		No. 4763395
1. Name and address of exporter:	2. Declared name and address of consignee:	
LEE ORCHIDS CO., LTD. 34/19 MOO 1 PETCHKASEM RD., BANGPAI BANGKAE BANGKOK 10160 THAILAND	EXIMARK TRADING INC SUITE 416 BPI CONNDOMINIUM PLAZA BINONDO, MANILA PHILIPPINES	
3. Number and description of packages:	4. Distinguishing marks:	
2 CARTONS		
5. Place of origin:	6. Declared means of conveyance:	7. Declared point of entry:
THAILAND	AIR	MANILA, PHILIPPINES
8. Name of produce and quantity declared:	9. Botanical name of plants:	
DENDROBILUM HYBRIDS 300,000 STEM(S) MOKARA HYBRIDS 1,236,000 STEM(S) ARANTHERA HYBRIDS 400,000 STEMS(S) (2 CARTONS OF FRESH ORCHID CUT FLOWERS)	DENDROBILUM HYBRID, MOKARA HYBRID, ARANTHERA HYBRID	
This is to certify that the plants, plant products or other regulated articles described herein have been inspected and/or tested according to appropriate official procedures and are considered to be free from the quarantine pests specified by the importing contracting party and to conform with the current phytosanitary requirements of the importing contracting party, including those for regulated non-quarantine pests.		
Additional Declaration		
NOTHING TO DECLARE		
Disinfestation and/or Disinfection Treatment		
10. Date:	11. Treatment:	12. Chemical (active ingredient):
MARCH 11, 2015	FUMIGATION	METHYL BROMIDE (CH3 Br)
13. Duration and temperature:	14. Concentration:	15. Additional information:
1.5 HRS/21 DEGREES CELSIUS OR ABOVE	20.00 GMS./CU.M.	TREATED BY EXPORTER
16. Stamp of organization:	17. Place of issue:	19. Name and signature of authorized officer:
	SUARNABHUMI AIRPORT, SAMUTPRAKAN THAILAND	 MS. PISOM CHAROENSRI FOR CHIEF, SUARNABHUMI AIRPORT PLANT QUARANTINE STATION
	18. Date:	
	MARCH 11, 2015	
NOTE : No financial liability with respect to this certificate shall attach to the Ministry of Agriculture and Cooperatives, Thailand, or any of its officers or representatives of that Ministry.		

982809


Inspection of Passengers Luggage at the Arrival Area



Inspection of Passengers luggage at the arrival area




Some of the confiscated commodities brought in by passengers



Interception at the Cargo Area








**PQ INSPECTORS CHECK
CONTAINER NUMBER
AGAINST THE IMPORT
DOCUMENTS**



Confiscation Procedures



INSPECTION AND SEIZED COMMODITIES



CONFISCATED COMMODITIES



MISDECLARATION OF GOODS



DISPOSAL OF SEIZED COMMODITIES





Presentation of the 67 boxes fresh ginger



Pouring of diesel fuel by PQ Inspector



Initiating the burning of the ginger



Boxes of fresh ginger on fire

Laboratory Procedures






Laboratory Analysis of Intercepted commodities



Detection of *Colletotrichum trichellum* on dragon fruit

Laboratory Analysis of Intercepted Commodities



Detection of *Colletotrichum gloeosporioides* (Penz.) Penz. & Sacc. on capsicum at NAIA

Laboratory Analysis of Intercepted Commodities



Baermann funnel method done for confiscated olive plant



Penicillium italicum
(Blue mould) of citrus)



Penicillium expansum
(Blue mould /soft rot)




Brown rot – lychee with
Fusarium growth)



Brown rot



Bitter rot of Apple




Apple mealybug
(Planococcus acaris)



Dead larva on apple
(unidentified)



Black powdery mould



Penicillium sp.



Aspergillus sp.



Alternaria sp.






Fusarium sp.

Laboratory Examination



Mango varieties susceptible to seed weevil

Laboratory Analysis of Intercepted Commodities



Mango seed weevil was detected from intercepted mango from Thailand on 22 April 2014



Laboratory Examination





Domestic Quarantine





BPI Special Quarantine Administrative Order No. 20, series of 1987


Declaring Mango Pulp Weevil *Sternochetus frigidus* (Fabr.) a Dangerous Pest and injurious to Mangoes and Placing the Palawan Island Group Under Quarantine to Prevent the Spread of Said Pest



Inset
KALAYAAN
GROUP OF
ISLANDS



**MANGO PULP WEEVIL
FREE AREA**




**Southern Palawan infested of Mango
Pulp Weevil**




Movement, transfer or carrying of mango trees, fruits, or parts thereof capable of harboring mango pulp weevil from Palawan Group of Island is prohibited.

It may be allowed in limited quantity and for experimental purpose only subject to the conditions that a permit for such purpose is first secured from the Director of BPI.





- Mangoes determined to be detected with the presence of the MPW will automatically be confiscated and disposed accordingly to prevent the spread of MPW within the City.
- Proper disposal shall include soaking with insecticides and burning of mango fruits.
- All mangoes in Non-Commercial quantity, wherein each mango only weigh less than 125 grams, but has to be transported and brought to and within the City shall undergo internal examination.
- Only mangoes weighing 125 grams and above shall be tested for MPW by passing through the City MPW Detection machine
- Checkpoints manned by the City PNP and personnel of BPI shall be established in the Northern and Southern part of Puerto Princesa




Establishment of Pest Free Areas





DA Administrative Circular No. 06, series of 2014

- ❖ Declaring the Whole Philippines as Area Free from Mango Seed Weevil (MSW) *Sternochetus mangiferae* Fabr. And Except for Palawan, Area Free From Mango Pulp Weevil (MPW) *Sternochetus frigidus* (Fabr.) and Prohibiting Importation From Countries Where the Pests Exist.





DA Administrative Circular No. 06, series of 2014

Movement, transfer or carrying of mango trees, fruits, or parts thereof capable of harboring mango pulp weevil from Palawan Group of Island is prohibited.


Importation / entry of mango fruits/mango plants and parts thereof / planting material from areas, places or countries where MPW and MSW exist is likewise prohibited.





Recognition as a Pest Free Area


October 1, 2014 – USDA issued the Final Rule Declaring the Philippines as Area free from MPW (except Palawan) and MSW



Australian Government

**Department of Agriculture,
Fisheries and Forestry**

December 18, 2015 -
DAFF recognition of the
whole Philippines, except
Palawan, as MPW and the
whole Philippines as MSW
free area.



NATIONWIDE LOW MONITORING OF MANGO PULP AND SEED WEEVIL ON MANGO PRODUCTION REGIONS/ PROVINGES OF THE PHILIPPINES



FRUIT COLLECTION



FRUIT DISSECTION





ASEAN Regional Training Workshop on Diagnostics of Weevils of Quarantine Importance

July 10-22, 2017

**Institute of Weed Science, Entomology and Plant Pathology
College of Agriculture and Food Science
University of the Philippines Los Baños**

SESSION 11

RESOURCES OF WEEVILS ON THE WEB

Lecture Notes by:

DR. HIRAKU YOSHITAKE
Institute for Agro-Environmental Sciences
NARO, Tsukuba, Japan

RESOURCES OF WEEVILS ON THE WEB

Browsing of web sites concerning with weevils

International Weevil Community Website

Welcome to International Weevil Community Website

This is a website for weevil taxonomists and others interested in the beetle superfamily Curculionidae, with the aim of facilitating cooperation and information-sharing worldwide. It will enable weevil specialists to share information online and together develop a community resource with data on weevil hosts, morphology, systematics, taxonomic characters, literature and other areas of mutual interest. The amount of information available now vastly outstrips our ability to keep track of it. For several years weevil taxonomists have realised they need a web resource where they can discuss issues and share information, for example on taxonomic character states, to minimise confusion in the literature and enable acceleration of research. This site grew from that need.

While anyone can access the content of the site (other than private work areas set up for small teams) only those with an account can contribute to the site or edit content. If you wish to become a contributor please click on the 'Log In' button on the top right of the screen, and then select 'Create New account'. When you complete the on-line form please include your interests and affiliation.

The site is dedicated in the first place to weevil systematics and taxonomy, but is by its nature responsive to the way in which we, the users, choose to develop it.

The site is maintained by Chris Lylol on behalf of contributors, who retain copyright.

<http://weevil.info/>

A website of International Weevil Community containing many fundamental data of weevils.

Electronic Catalogue of Weevil names (Curculionoidea)

WTaxa Electronic Catalogue of Weevil names (Curculionoidea)

Web Version: 3.6 Database Version: 1.7 (31 Apr 2014)

Please allow pop-up windows for this site:

The Electronic Catalogue of Weevil Names is a project in progress.

Ultimately it will contain all family, genus and species-group names in the Curculionoidea, a group of beetles with some 60,000 described species which includes many important insects, including the bark beetles (Scolytinae) and powder-post beetles (Platypodinae).

Currently the database contains some 176,000 names, including all generic and family-group names, in their current taxonomic position, and based on Alonso-Zarazaga & Lylol's (1998, 2002) catalogue of weevil family and genus names. Species-group names are being added, but are not yet all present (or checked against original publication or correct placement).

NEWS! The gathering of names from secondary sources is finished and considered complete as of 31st December 2012. We will continue checking names from the original sources and updating the nomenclature, deleting duplicates and unnecessary records, and adding new features to the display and contents to the database. We will be very grateful if any mistake or absence is notified.

<http://wtaxa.csic.es/>

A database of scientific names of weevils, but is still under development.


Japanese Weevil Database



<http://de05.digitalasia.chubu.ac.jp/>

A database of Japanese weevils, containing images of many species in common to / related to those from the ASEAN region. All the navigation is in Japanese, but it is very easy to find your way around.


Integrated insect types database of Taiwanese species



<http://twinsecttype.nmns.edu.tw/>

A database of the type specimens preserved in various institutions throughout world, containing data of Taiwanese weevils, most of which are in common to / related to those from the ASEAN region.


New Zealand Weevil Images



<http://www.landcareresearch.co.nz/science/plants-animals-fungi/animals/invertebrates/systematics/weevils>

A database of New Zealand weevils, some of which are in common to / related to those from the ASEAN region, containing many habitus images.

Curculio Institute - Center for western Palearctic Curculionioidea



<http://www.curci.de/institute3/index.php>

The website of the Curculio Institute containing a series of “Digital-Weevil-Determination” for Curculionoidea of West Palaeartic on “SNUDEBILLERonline”. The annual membership fee of EUR 50,00 to log in SNUDEBILLERonline and view all current articles.

Potential Invasive Weevils of the World

<http://www.piweevils.com/?app.html>

A database of important weevils for plant quarantine due to the high invasion risk.